

Remote follow-up technologies in traumatic brain injury: a scoping review

Short running title: Follow-up technologies in TBI: a scoping review

Brandon G. Smith*^{1,2}, Stasa Tumpa³, Orla Mantle⁴, Charlotte J. Whiffin^{2,5}, Harry Mee^{1,6}, Davi J. Fontoura Solla^{2,7},
Wellingson S. Paiva^{2,7}, Virginia F J Newcombe⁸, Angelos G. Koliass^{1,2}, Peter J. Hutchinson^{1,2}

* Corresponding author

¹ Division of Neurosurgery, Department of Clinical Neurosciences, Addenbrooke's Hospital and University of Cambridge, Cambridge, U.K.

² NIHR Global Health Research Group on Neurotrauma, University of Cambridge, Cambridge, U.K.

³ School of Clinical Medicine, University of Cambridge, Cambridge, U.K.

⁴ GKT School of Medical Education, King's College London, London, U.K.

⁵ College of Health, Psychology and Social Care, University of Derby, Derby, U.K.

⁶ Division of Rehabilitation Medicine, Addenbrooke's Hospital and University of Cambridge, Cambridge, U.K.

⁷ Division of Neurosurgery, Department of Neurology, University of São Paulo, Brazil

⁸ University Division of Anaesthesia, University of Cambridge, Cambridge, U.K.

Keywords:

Traumatic brain injury, follow-up technology, patient-generated health data, innovation, outcome assessment, telemedicine

Corresponding author: Brandon G. Smith

Author details:

Brandon G. Smith (bgs30@cam.ac.uk / +44 07870 805420)	Davi J. Fontoura Solla (davisolla@hotmail.com / +55 11 94234-1989)
Stasa Tumpa (st731@cam.ac.uk / +44 07503 730996)	Wellington S. Paiva (wellingsonpaiva@hotmail.com / +55 11 2661 7226)
Orla Mantle (orla.mantle@kcl.ac.uk / +44 07823 887303)	Virginia F. J. Newcombe (vfjn2@cam.ac.uk / +44 01223 217889)
Charlotte J. Whiffin (c.whiffin@derby.ac.uk / +44 01332 590500)	Angelos G. Koliass (ak721@cam.ac.uk / +44 01223 336946)
Harry Mee (hwjm2@cam.ac.uk / +44 01223 336946)	Peter J. Hutchinson (pjah2@cam.ac.uk / +44 01223 336946)

Abstract

Traumatic brain injury (TBI) remains a leading cause of death and disability worldwide. Motivations for outcome data collection in TBI are threefold: to improve patient outcomes; facilitate research; and provide the means and methods for wider injury surveillance. Such data plays a pivotal role in population health, and ways to increase the reliability of data collection following TBI should be pursued. As a result, technology-aided follow-up of neurotrauma patients is on the rise; there is, therefore, a need to describe how such technologies have been used. A scoping review was conducted and reported using the PRISMA extension (PRISMA-ScR). Five electronic databases (Embase, MEDLINE, Global Health, PsycInfo, Scopus) were searched systematically using keywords derived from the concepts of ‘telemedicine’, ‘TBI’, ‘outcome assessment’, and ‘patient-generated health data’. Forty studies described follow-up technologies (FUTs) utilizing telephones (52.5%, n = 21), SMS (10%, n = 4), smartphones (22.5%, n = 9), videoconferencing (10%, n = 4), digital assistants (2.5%, n = 1), and custom devices (2.5%, n = 1) amongst TBI patient cohorts of varying injury severity. Where reported, clinical facilitators, remote follow-up timing and intervals between sessions, synchronicity of follow-up instances, proxy involvement, outcome measures utilized, and technology evaluation efforts are described. FUTs can aid more temporally-sensitive assessments and capture fluctuating sequelae, a benefit of particular relevance to TBI cohorts. However, the evidence base surrounding FUTs remains in its infancy, particularly with respect to large samples, low- and middle-income patient cohorts, and the validation of outcome measures for deployment via such remote technology.

Introduction

Traumatic brain injury (TBI) is a concern in both public and global health, and is a leading cause of death and disability worldwide.^{1,2} Secondary deficits from TBI manifest in multiple ways, often with long-term symptoms in physical, cognitive and emotional domains,³⁻⁵ which have a collateral impact, both direct and indirect, on patients, families and wider society.^{2,3,6,7} The purpose of outcome data collection is threefold: to improve patient outcomes, to facilitate research, and to provide the means and methods for wider injury surveillance.

Each purpose may vary in the fidelity of the data sought - injury surveillance efforts may typically aim to garner general morbidity or mortality on a wider scale, whereas research initiatives may use refined, detailed assessment batteries centered upon the phenomena or sequelae in question. In clinical practice, we may seek a balance of both; employing more detailed outcome measures and assessments as required, whilst also addressing any ongoing sequelae and determining the general status and wellbeing of the patient.

In 2004, the World Health Organization (WHO) released Guidelines for Essential Trauma Care emphasizing the importance of surveillance data to reduce the global burden of death and disability from injuries.⁸ In addition to enabling clinical teams to determine the full extent of physical, mental and socio-economic sequelae post-injury,^{5,9-11} accurate data also facilitates the evaluation of systems and services, including: the efficacy of patient treatment and management decisions,^{10,11} identifying targets of wider systems improvement in injury prevention,¹² enabling continuous quality improvement projects and trials, and the formation of registries that may themselves be incorporated into care pathways, injury prevention strategies and policies,^{13,14} and lastly, to facilitate rehabilitation of an individual and improve rehabilitation pathways and services.

Despite improvements in injury surveillance data, data on disability and long-term functional outcomes remain poorly recorded in both high-income countries (HICs) and low- and middle-income countries (LMICs).^{5,10,15,16} However, given the heterogeneity both within and between LMICs, the collection of outcome data is considered more complex^{1,10,17} and as such is often limited to collection at hospital discharge only.¹⁰ Despite these limitations very little has been published on the challenges faced in facilitating long-term follow-up and collection of outcome data in LMICs. Of the research

that does exist, efforts in data collection at the clinical level were found to be complicated by factors such as weak healthcare and long-term support infrastructure, resulting in a lack of regular follow-up of trauma patients.¹⁰

In 2007 the world's population became more urban than rural for the first time.¹⁸ However, in many countries, and in particular LMICs, vast numbers remain in rural settings, and with this, have limited access to general health services. In the context of specialist services such as neurosurgery, which remain heavily centralized to urban settings, clinicians often have limited or no regular access to patients after discharge.¹⁹⁻²¹ Patients that are able to access neurosurgical follow-up often have to travel extensive distances at great personal cost.^{22,23} Those unable to access neurosurgical follow-up become lost to follow-up.^{24,25}

Over the last few decades, there has been rapid advancements in technology, especially in regard to telecommunications and its widespread adoption. According to a United Nations (UN) International Telecommunication Union (ITU) 2020 report,²⁶ 47% of households worldwide are estimated to have access to a computer, with 57% of households perceived to have internet access. Mobile phone and cellular network technologies however are arguably the 'common-denominator' technology worldwide. An estimated 75 per 100 of the world's population are thought to have an active mobile broadband connection, and 105 mobile-cellular telephone subscriptions per 100 population.

Similarly, short message service (SMS) technologies, initially made possible by second-generation or '2G' cellular network technology, remains a valuable opportunity for outcome data collection owing to its worldwide penetration and strong coverage, even in an LMIC setting where it is estimated that over 95% of the population have 2G coverage.²⁷ The adoption of next-generation network architecture, such as fifth generation (5G) cellular technology, can be expected to lead to exciting new possibilities for mobile health assessment owing to its ability to provide accessible, high-speed streaming capabilities²⁸ for use in high-definition remote video assessments.

Technologies are becoming frequently adapted to harness unique opportunities to connect patients with their health providers upon discharge. These follow-up technologies (FUTs) may provide innovative solutions that mitigate those deemed 'lost to follow-up', fill quantitative gaps in TBI epidemiology, and enable those in the remotest corners of the world to access specialist care.

Prior engagement with the literature on FUTs revealed a complex and diverse evidence base, and no prior attempt to synthesize this substantial body of work. A scoping review was therefore deemed an appropriate methodology to describe the type, characteristics, and size of the evidence in this field. The objectives of this review were to identify follow-up technologies, such as telephone- and SMS-based service, videoconferencing, and smartphone applications, implemented across global settings in traumatic brain injury. Secondly, this review aimed to characterize the outcome measures administered and the data collected, amongst the communication modalities used, and briefly highlight the success of each in context with the patient populations and settings they have been implemented in.

Materials and Methods

The reporting of this study is in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for scoping reviews (PRISMA-ScR),²⁹ with additional study screening and selection flowchart (Figure 1) as recommended by the JBI and PRISMA-ScR reporting guidelines.²⁹⁻³¹ Unlike systematic reviews, scoping reviews do not require an antecedent protocol registration,³² notwithstanding, review objectives, eligibility criteria and preliminary study characteristics to be extracted were determined a priori.

Our scoping review was guided by the comprehensive scoping review framework by Arksey and O'Malley,³³ with minor refinements, including those suggested by Levac *et al.*³⁴ and the Joanna Briggs Institute (JBI).³⁰ This framework includes: (1) Identifying the research question, (2) Identifying relevant studies, (3) Study selection, (4) Charting the data, (5) Collating, summarizing, and reporting the results, and an optional (6) Consultation exercise. This review serves as part of a wider programme of work in which stakeholders (neurotrauma physicians) are formally engaged in a separate qualitative research process, examining their views, experiences and opinions of FUTs as applied to post-TBI care provision. Prior to commencing the review, we shared similar perceptions to Levac *et al.*³⁴ regarding the challenges of conducting and integrating the results of stakeholder consultation within a review's findings. In light of this, and given the potential of qualitative inquiry to empower rich & in-depth investigations of the human experience,³⁵ stage 6 was not carried out within this review. Notwithstanding, the lead author (BGS) discussed findings with senior co-authors, who by their nature as LMIC neurosurgeons are stakeholders within this context.

Research question

What research has been conducted to describe, examine, or assess the use of follow-up technologies in traumatic brain injury cohorts worldwide?

To answer this question, the following sub-questions were posed:

- 1) What technologies are being used as FUTs in a global setting?
- 2) In what settings is FUT research being conducted?
- 3) What patient cohorts (demographics, injury severity) have been included in FUT research?
- 4) What factors may constitute successful implementation of FUTs?
- 5) What, if any, validated outcome measures are being deployed via FUTs?

Search Strategy

The final search strategy was determined with the assistance of an academic medical librarian following a consultation to derive keywords based on the review objective and concepts of ‘telemedicine’, ‘traumatic brain injury’, ‘outcome assessment’ and ‘patient-generated health data’. The authors selected a number of ‘indicator papers’ - predefined articles that one would expect to appear in their final search results - to test the quality and robustness of our search strategy. Several pilot searches were attempted before a final strategy was established and translated across a number of databases. The search strategy was executed on the 1st October 2021 on the following electronic databases: OVID Embase, OVID MEDLINE, OVID Global Health, OVID PsycInfo and Scopus. These databases were selected owing to their sufficient coverage given the multi-disciplinary nature of TBI, outcomes, and their assessment. A limited manual search was conducted on Google Scholar and a number of technology- and head-injury focused journals (Journal of Neurotrauma, The Journal of Head Trauma Rehabilitation, World Neurosurgery, NEUROSURGERY, Journal of Medical Internet Research, Journal of Telemedicine and Telecare). Following a pre-protocol pilot gray literature search, it became apparent that, of the limited materials retrieved, most were unsuitable for a number of reasons, including: insufficient indication of external peer review (of particular importance in commercial reports); a lack of FUT description or elucidation of the methods in their use; description or evaluation of the FUT was not perceived to be the primary focus of the resource. Compounded by the resource- and time-intensive nature of conducting a gray literature search in this context, gray literature sources were omitted in the final search strategy. Finally, a backward citation searching was undertaken, whereby the reference lists of papers deemed eligible for inclusion, and review papers which were not

eligible for inclusion, were screened for relevant studies. Searching for additional sources was completed on 25th April 2022.

Eligibility Criteria

Scientific articles reporting original research of the application of technology written in the English were included. All databases were searched from inception to achieve the largest scope possible and detail early innovations in this field.

The Population/Participants, Concept & Context (PCC) framework^{36,37} was used to inform our inclusion criteria, search and data charting strategies.

Eligibility criteria for articles to be included in this review were: (1) Any published original research, including: primary studies, reports, editorials, opinion papers, letters, conference abstracts, theses, and book chapters; (2) Reports with a primary aim to describe, assess, or examine the use of FUTs to facilitate remote collection of patient outcome data; and (3) Adult and pediatric all-severity TBI patient cohorts. Articles were excluded if they were (1) Study protocols or secondary research (reviews); (2) Reports describing the collection of family or caregiver outcomes only; or (3) No TBI population or involved healthy volunteers only.

Population/Participants

Studies were only eligible for inclusion in this review if their primary aim related to the development, implementation or validation of technologies contributing to the provision of follow-up care of discharged patients following TBI of any severity, whether directly or via proxy (family members/relatives, carers and guardians). Studies of mixed-pathology cohorts were included.

Concept

In this review, we defined follow-up technology as any system, device, equipment, component, or machinery used to both transmit and receive digital information electronically between a remote outpatient or their proxy, and a member of their clinical team. The focus of this exchange is to attain data from the patient as to their current welfare status in the form of either patient-generated health data (PGHD), i.e., self-reported, or clinician-derived health data (CDHD), i.e., garnered through clinician-led assessments or interview, either from the patient or their nominated proxy. In addition, we defined 'remote' as, at the time of information exchange, the patient resided in a community-based setting

(home, regional care providers such as a general practitioner, regional hospital, or other public settings) at a distance from the clinical team managing their follow-up.

Herein, we refer to synchronicity as the temporal aspect of the encounter between clinician and patient. Synchronous FUTs function to collect data in real-time, often allowing direct contact between clinician and patient through sensor-, text-, voice-, or video-based technology. Asynchronous FUTs collect data by store-and-forward techniques - data are gathered, stored, and transmitted for later review by the clinician at two independent timepoints; i.e., they do not interact in real-time.³⁸

Finally, we define ‘follow-up’ as any attempt to monitor, assess, communicate, or liaise with a patient, or their proxy, from the point of hospital discharge, for the benefits of furthering their health and wellbeing, research, or injury surveillance.

Context

Follow-up technologies utilized in any global health setting. We utilize HIC and LMIC classifications as defined by the 2021-2022 World Bank list of economies.³⁹

Study Selection

A two-stage screening process was followed. All search results were initially imported into the Zotero (Corporation for Digital Scholarship, Virginia, USA) reference manager for title and abstract review. Where necessary, duplicates were removed manually. Two researchers (BGS and ST) independently screened all titles and abstracts, and potentially eligible studies were identified for full-text review.

Disagreements arising from the selection process were either resolved by consensus, or where this was not possible, a third reviewer (OM) was consulted for resolution. Following preliminary screening, the remaining articles were exported to Microsoft® Excel (Microsoft, Redmond, Washington, USA), where full texts were independently screened by two researchers (BGS and ST) for final eligibility; a third researcher (OM) was consulted for disagreements as required. Selected studies formed the final repository of evidence for subsequent data extraction (charting) and collation.

Data Charting and Synthesis of Results

Owing to the exploratory nature of a scoping review, a precursor proforma was developed to facilitate data extraction. Within this document, initial elements of interest that sought to answer our research question were informed by our PCC framework and agreed upon by researcher consensus (BGS, ST, CJW, AGK). The proforma was subdivided into five key sections, including: authorship and study characteristics, description of neurotrauma, characteristics of FUT(s) utilized, and major findings & challenges. This was iteratively updated and refined as the charting process progressed, adding to pre-identified elements of interest (see Supplementary Material). Data charting was conducted independently for all articles by two researchers (BGS and ST). Following the charting of the first five studies, in line with recommendations by Levac *et al.*,³⁴ the authors reconvened to ensure proforma suitability in addressing the research question, and to advance the proforma following familiarization with this initial subset of studies. Final completed proformas were cross-checked for conformity, and a third researcher (OM) was consulted as necessary in cases of dispute. No formal critical appraisal, nor quality of evidence assessment, was conducted as this falls beyond the remit of a scoping review.⁴⁰ Following data charting, a narrative summary of included articles was constructed in relation to the review's overarching question and sub-questions.

Results

Characteristics of Included Studies

Executing the search strategy across the five electronic databases yielded a total of 1562 potentially eligible citations. Following deduplication, 1168 unique articles remained, with a subsequent title and abstract review delineating a pool of 96 citations for further full-text review. This final stage of screening concluded with 40 articles for inclusion; inclusive of 11 citations discovered through manual and citation searches. A full PRISMA-ScR flowchart for the study search, selection and exclusion process is depicted in Figure 1.

Of the 40 articles retrieved, the plurality (n = 15) were reported or deemed to be descriptive in design,^{19,41–54} and included small, non-randomized pilot studies and secondary analyses of data, or in two studies, were conducted in or described the retrospective analysis of a quality improvement initiative format.^{52,53} Citations with an observational design formed the second most common type (n = 10),^{10,55–63} encompassing prospective^{55,56,58} and retrospective¹⁰ cohort studies and cross-sectional studies.^{57,60,61} Research of experimental (n = 9)^{64–72} and quasi-experimental (n = 6)^{73–78} design were similar in frequency. Experimental designs included single-centre^{66,68–71} and international multi-centre^{65,67} randomized

trial designs, amongst non-randomized, open-label trials^{64,72}. Results of these studies were published primarily as original research articles (n = 31),^{10,19,41–44,48–53,55,56,59,61–75,78} with a lesser number as conference abstracts or research posters (n = 8),^{45–47,57,58,60,76,77} and research summary letters (n = 1).⁵⁴

International Context

The adaption of technologies for follow-up delivery has been used in multiple settings throughout the world. The majority of papers reported on FUTs in HICs (n = 34, 85%), including the USA (n = 24),^{41–43,51–61,63,64,66,67,72–74,76–78} Australia (n = 3),^{70,71,75} Canada (n = 3),^{45,47,68} Netherlands (n = 1),⁶² Ireland (n = 2)^{46,50} and a joint endeavor between Italy, Spain and Belgium (n = 1).⁶⁵ Whereas only six (15%) reported findings from studies in LMICs including Uganda (n = 2),^{19,49} Ethiopia (n = 1),¹⁰ India (n = 1),⁴⁴ Iran (n = 1),⁶⁹ Indonesia (n = 1).^{48,79}

Patient Population Demographics & TBI Characteristics

The majority of articles (n = 27, 67.5%) described civilian adult population cohorts (>18 years),^{10,41,42,44–48,50–52,55,57–60,62,65,66,69–71,73–75,77,79} from sample statistics reported representing 3,442 patients. A further five studies described military or veteran cohorts^{43,61,64,67,76} representing an additional 207 adult patients. Five studies reported exclusively pediatric patient cohorts,^{19,54,63,68,72} representing 287 patients. Three studies investigated mixed adult and pediatric cohorts,^{49,56,78} adding a further 774 patients to previous approximations. Two studies did not report the demographics of the cohorts investigated.^{44,53}

With respect to TBI severity, FUTs were most frequently implemented in TBI patient cohorts of undefined severity (n = 14)^{10,44,45,49,52,53,55,58,62,65,67,70,71,77} - in some of these cases, patient cohorts were pooled with other diagnoses (trauma, spinal cord injury, stroke, orthopedic, acquired brain injury amongst other neurologic conditions). Where TBI severity was defined, seven studies (n = 7) explored the implementation of FUTs in all-severity TBI patient cohorts.^{19,41,48,51,73,74,76} In studies recruiting patients with particular injury severities, concussion or mild TBI formed the majority (n = 12),^{43,46,50,54,56,59,61,63,66,68,72,78} whilst only one study investigated FUTs as applied to a severe TBI cohort exclusively.⁷⁵ No studies reported moderate TBI cohorts exclusively. Of the remaining studies, two reported FUTs for mild to moderate TBI cohorts,^{47,69} and four for moderate to severe TBI cohorts.^{42,57,60,64} Further, two studies described their TBI cohort as chronic.^{51,58}

Almost all (n = 37) studies used follow-up technologies whilst the patient was at home or in another non-health setting in the community. The remaining studies (n = 3) used technology while the patient was at other clinical settings away from the team responsible for follow-up assessment, including a polytrauma rehabilitation center in one study,⁷⁶ and a hospital research laboratory 15 kilometers from the assessing clinician^{70,71} in two linked studies by the same author team.

Technology Modality

Remote FUTs were grouped by their underlying modality: telephone, SMS, smartphone (e.g., mobile application), videoconference, and ‘miscellaneous’ - namely a personal digital assistant (PDA) and custom touchscreen device, each demonstrated in one study respectively. Telephone-based FUTs (52.5%) were the first to appear for use with TBI patient cohorts in 1997,⁴¹ and remained the modal technology utilized at the time of search strategy execution. Videoconference-based technologies (10%) were next to appear a decade later in 2008 as part of a multi-center randomized trial.⁶⁵ A single study exploring PDA (2.5%) as a remote FUT for ecological momentary assessment (EMA) in a pediatric concussion cohort followed shortly after in 2009.⁶³ SMS-based FUTs (10%) followed in 2012, forming the asynchronous communication element of a pilot EMA study in the USA.⁴³ Smartphone-based FUTs (22.5%) were introduced in 2015. Lastly, a single study⁶² explored the use of a custom electronic touchscreen device, the ‘PsyMate’, in 2019 to investigate the feasibility of EMA to explore the interactions between person, environment and effect in an ABI cohort. A figure demonstrating the technology modalities implemented over time can be seen in Figure 2.

Telephone-based Follow-up

Telephone-based follow-up technology (Table 1) was used in 21 studies (52.5%).^{10,19,41,42,44–50,52,53,55,57,60,64,68,69,74,76} Telephones were primarily used as a synchronous means of two-way communication to conduct structured interviews with patients and their proxies. While almost all studies, where defined, used clinical or research staff to conduct follow-up, one used an external call center who had integrated their systems with the hospital’s electronic medical records⁴⁴ to complete the follow-up interview. Another study⁵⁵ had no human-facilitator in their administration of telephone-based follow-up, instead using an asynchronous and interactive voice response system (IVR), whereby pre-recorded questions are played to the patient, and either a voice or key-pad can be used to respond. One study using scripted, structured telephonic follow-up described the additional use of a secure web-based data capture platform (REDCap) with branching logic to conduct the interview.⁵² Lastly, one study reported the use of a computer-assisted telephone interviewing system,

whereby the assessor can follow a script in the collection of data, enabling an assessor without familiarity with TBI to conduct the interview.⁷⁴

SMS-based Follow-up

The studies included demonstrated a multimodal use of SMS-based technology (Table 2). Four studies demonstrated SMS exclusively as the vessel of data collection, often in an asynchronous and automated manner, through the delivery and eventual receipt of timed symptom assessments to patients in the community,^{43,56,59,66} whereby responses were often logged on a database for later review. Two studies, despite not using SMS directly in the collection of outcome data, utilized SMS to deliver prompts or updates. These text messages prompted the patient to log their current status and wellbeing on other systems such as a secure website or app-based portal.⁶⁷ In another study, SMS was used as a reminder system, informing the patients to expect to shortly receive a telephone call, with a focus on improving telephonic response and attrition rates.⁴⁵

Smartphone application-based Follow-up

The second technology most frequently used were mobile applications, or ‘apps’, installed on Apple and Android devices such as smartphones and tablets in the remote collection of outcome data (Table 3) (n = 9).^{51,54,58,61,67,72,73,77,78} Such implementations of mobile applications ranged from gamified symptom journals and social networking⁷² to delivering questionnaires for ecological momentary assessment (EMA).^{51,58,73} Three studies described the further use of the device’s onboard sensors and additional functionality in the delivery and collection of data, including native ‘push notifications’ to prompt patients to input data,⁷⁸ GPS tracking function in the collection of activity and community participation data,⁶¹ or in one study, using an Apple iPod Touch and its in-built accelerometer to capture objective measurements of physical activity.⁵⁴

Videoconferencing-based Follow-up

Three studies utilized videoconferencing exclusively for remote assessment (Table 4).^{70,71,75} Two studies, by the same authorship group, described their use of two robotically controlled web cameras controlled remotely by the assessing speech-language pathologist.^{70,71} Additionally, in their second paper with a system re-design, the authors describe concurrent automatic store-and-forward facilities integrated into their system, enabling video and audio data of higher quality than that streamed over the 128kbit/s videoconference connection to be sent to the assessing clinician for later

review.⁷¹ Further, one study reported the use of a novel, custom portable home care activity desk (HCAD) installed in the patient's home. Each unit consisted of sensorized tools and videoconferencing facilities, providing store-and-forward capability between the patient's home and hospital servers.⁶⁵ One study additionally described the use of telephone as a back-up option utilized in cases of videoconferencing technical difficulty.⁷⁵

Miscellaneous Technologies for Follow-up

Two studies, pooled together as 'miscellaneous' (Table 5), described the use of portable touchscreen electronic devices, one proprietary device 'PsyMate'⁶² and one use of a commercially available personal digital assistant (PDA) device.⁶³ Both studies described auditory prompt capabilities with their devices, enabling semi-randomly scheduled ecological momentary assessment throughout the day.^{62,63} One study previously noted described allowing the patient to choose between a secure internet website portal or telephonic IVR to submit self-reported outcome data.⁵⁵

Clinical Facilitators

Clinical facilitators were most often explicitly described in the paper as the research staff (i.e. research coordinators and care managers, research assistants, and clinical researchers),^{19,42,49,52,59,61,72,73,75} or hospital outcome management staff⁵⁵ and blinded examiners.⁶⁶ Nursing personnel^{41,52,69} formed the next largest pool of facilitators, of which included specialist psychiatric nurses,⁶⁴ rehab-based nurse practitioners,⁴⁷ or neurosurgical nurse personnel experienced in TBI and/or neurotrauma outcome measure administration.^{48,53,80} Undefined clinical staff,⁴³ speech-language pathologists,^{70,71} physicians,⁵² general therapists,⁶⁵ occupational therapists,⁶⁸ and care team members,⁶⁷ were described as facilitators in a smaller subset of studies. In three studies, facilitators were primarily external non-clinical staff without familiarity of TBI, such as data clerks,¹⁰ call center personnel,⁴⁴ or interviewers otherwise undefined.⁷⁴ However, 15 studies did not explicitly report on who facilitated the follow-up reported in the paper.^{45,46,50,51,54,56-58,60,62,63,76-78,81} While it may be assumed that the facilitators of technology in these studies were the author teams themselves, this cannot be confirmed.

Despite not being described as active facilitators, it was of note that in two studies, family members, friends and relative facilitators were incorporated in the delivery of the follow-up technology,^{43,72} and instead may be designated as 'passive facilitators'. One study with a pediatric population describes a 'social networking' function built in to the smartphone application, allowing friends and family to connect with and receive notifications of the patient's activities and progress.⁷² An additional study describes how designated friends or relatives could opt-in to receive notifications should

their injured relative fail to maintain contact with the FUT services, or if they returned a score below a pre-set threshold that warranted further one-to-one contact outside of the FUT.⁴³

Timing of Follow-up & Time Since Injury, Diagnosis or Discharge

The timing of remote follow-up delivery with respect to the patient's time since injury/diagnosis (TSI/D), or time since hospital discharge, could be established directly or approximated in 77.5% (n = 31) of articles retrieved. Studies were broadly grouped together by respective timeframes of: Less than 1 month, 1 - 3 months, 3 - 6 months, 6 - 12 months, and 12 months or more. Eight (20%) studies reported remote assessment within a month of discharge or injury;^{42,45,47,54,59,66,68,78} within this group, five studies^{54,59,66,68,78} described recruitment and assessment of participants from discharge up to two weeks post-discharge for the assessment of concussion or mild TBI. Three (7.5%) studies^{41,48,64} assessed patients between 1- and 3-months post injury or discharge, and all utilized telephone as the modality of choice. Five (12.5%) studies^{10,52,63,69,76} conducted remote assessment beyond 3 months and within 6 months of discharge or injury. Four studies (10%) described assessment between 6 and 12 months.^{46,58,70,72} Lastly, 27.5% (n = 11) of studies^{49-51,57,60,65,71,73-75,82} depicted an average time point of remote assessment of a year and beyond hospital discharge or injury. Nine studies (22.5%) did not formally define the time point at which remote assessment was attempted.^{43,44,53,55,56,61,62,67,77}

Intervals between sessions

The most frequently reported (n = 14) timeframe of outcome data collection by FUT was at one single time point following injury or discharge.^{10,19,44,46,47,49,50,52,55,60,70,71,74,75} Of these studies, eleven (n = 11, 78.6%) utilized telephone-based technology.^{10,19,44,46,47,49,50,52,55,60,74} One study used two modalities in a single instance of follow-up, namely a secure web-portal and telephone-based interactive voice response (IVR) system.⁵⁵ Three studies used videoconferencing in a single instance.^{70,71,75}

The remaining studies (n = 26, 65%) reported more than one outcome data collection point. One study reported follow-up at yearly intervals for 2 years post-injury.⁵⁷ Another study reported two follow-up points at quarterly intervals of 3- and 6-months post-injury.⁴¹ One study described data collection on a monthly basis by telephone up to 3 months following discharge.⁴⁸ Similarly, a further study described fortnightly instances of follow-up by telephone over the course of 4 weeks.⁴⁵ Three studies reported collecting outcomes on a weekly basis^{43,64,69} - two of which were by telephone^{64,69} - and the first use of SMS is seen at this weekly interval.⁴³ In those studies using follow-up technology on

a daily basis, eight studies utilized smartphone applications^{51,54,58,61,67,72,73} and one employed a videoconference-based, sensorized homecare activity desk (HCAD) previously described.⁶⁵ Seven studies reported collecting data multiple times throughout the day;^{56,59,62,63,66,77,78} this session interval also used the most diverse range of technologies of the intervals described thus far. Three studies explored the use of SMS up to five times daily,^{56,59,66} and two studies demonstrated the use of smartphone applications up to four times daily.^{77,78} The remaining studies used a PDA⁶³ and custom touchscreen device⁶² to examine outcomes up to 10 times daily. Lastly, three studies employing telephone as a FUT collected data at steadily increasing intervals up to 9 months post-injury.^{42,68,76} One additional study, owing to the nature of the telephone-based service (a neurotrauma hotline), was not able to define an interval between sessions however reported an average of 3.3 calls per day over a 12 month period.⁵³ A visual summary of the intervals between FUT sessions with respect to technology modality can be found in Figure 3.

Synchronicity

Half of the studies utilized synchronous or real-time assessment in the collection of outcome data (n = 20, 50%).^{10,19,41,42,44–50,52,57,60,64,68,69,74–76} Sixteen (40%) used an asynchronous or store-and-forward approach.^{43,51,54–56,58,59,61–63,66,67,72,73,77,78} Four studies (10%) demonstrated the use of mixed synchronicity,^{53,65,70,71} often utilizing different capabilities of the technology in its service delivery, such as voicemail (asynchronous) services built in to a neurotrauma hotline (synchronous).⁵³

Use of Proxies

In 29 studies, there was no reported use of proxies (72.5%).^{41,43–47,50,51,54–63,65–67,70–74,76–78} Eight studies (20%) were extended to both patients or their proxies in remote follow-up.^{10,42,48,49,52,53,64,75} Of these studies, two (5%) used a mixture of both patient and proxy data, usually collected in the same sitting, to assess the patient's wellbeing.^{42,75} A further two studies described the assistive role of proxies where the patient was unable to directly report their wellbeing themselves.^{10,52}

Two studies (5%) described the role of proxies in assisting initiating contact with the patient if they were initially unable to be reached, although did not further disclose whether they were used to seek information on behalf of the patient.^{48,49} One study, in addition to assisting the patient report their wellbeing, used proxies to report data pertaining to mortality as necessary.⁵² One study,⁴³ despite not using a proxy to ascertain data regarding the status of health of the patient on

their behalf, utilized a clinical staff or ‘buddy’ contact system of friends and family to facilitate further check-in with the patient, should they not respond to the asynchronous technology-based wellbeing report, or not meet a threshold score indicating they were otherwise well.

Three studies (7.5%)^{53,64,69} described telephone-based support made available for patients and their proxies,^{53,64} and their proxies exclusively,⁶⁹ to share symptoms, wellbeing status (on behalf of the patient where required), or seek support at their own convenience, in between scheduled requests or prompts for patient status reports.^{64,69}

Three studies (7.5%) described the use of proxies exclusively to assess patients in their FUT deployment, without directly communicating with the patients themselves.^{19,68,69} Pediatric patient cohorts formed the basis of two of these studies,^{19,68} whereby follow-up and outcome data collection was solely provided by the patient’s caregivers owing to the patient’s age.

Deployment of Outcome Measures

Several studies utilized one or more validated outcome measures in their technologies, whilst a small number of studies used internally developed measures, scores and scales in the remote assessment of patients. The Glasgow Outcome Scale Extended (GOS-E) was the most frequently used in the studies included (n = 6), followed by the Rivermead Post-concussion Questionnaire or its derivatives (n = 5), and the 5-Level EuroQol 5-Dimension instrument (n = 4). A full list of outcome measures deployed in FUTs for TBI, ordered by frequency, can be found in Table 6.

In addition, many studies used generalized questionnaires briefly ascertaining overall wellbeing^{43,52,62,67} and fatigue,^{51,58,62,73} presence of pain,⁶⁷ past and current medical concerns,⁴² physical deficit or symptom checklists,^{19,45,49,64} both generalized and specific (e.g., headache, irritability, depression, memory problems, medication compliance and other miscellaneous complications and symptoms). A handful of studies deployed similar generalized questionnaires, yet with a focus on reporting-by-proxy through the patient’s caregiver.⁶⁹

Where reported, these broader questions often existed either alongside or exclusive to generic questions encompassing: employment or return to work,^{41,42,49,60} household and leisure activities, or activities of daily living,^{19,41,49,54,64,68} mood,^{42,67} energy and sleep status,⁶⁷ personal finance management,⁴¹ relationship status,⁶⁷ subjective return to pre-injury or baseline

status,⁴⁹ travel and location,^{41,62} and lastly social context, social activities & community participation.^{41,62-64} In pediatric populations, some generalized questionnaires sought information regarding the impact of injury or its sequelae on schooling and education,^{49,68} and play activities.⁶⁸ One study with pediatric participants⁵⁴ described measurement of daily cognitive rest and exertion by calculating a composite score of number of text messages sent, minutes of screen time and gaming, and minutes of reading and schoolwork.

Other domains, inclusive of those aforementioned, were assessed by one study as part of a structured interview addressing 17 broad domains including personal care, ambulation, home management, leisure, alcohol and drugs use, legal issues and spirituality⁴². One study⁷⁶ differed from those previously reported in deploying a neuropsychological assessment battery (including standard verbally administered tests of attention, memory, working memory, processing speed, language, executive skills and auditory-verbal adaptations of trail-making). Another study⁷¹ utilized informal oromotor and perpetual speech assessments as part of a wider speech-language battery. Weight status was included in one study with a longer duration of 36 weeks⁶⁷.

One study, instead of directly asking for a subjective measure of travel, activities or social and community participation, collected GPS-based activity data ('MOVES Storylines') to quantify this measure objectively and indirectly⁶¹. Similarly, another study⁵⁴ utilized the device's onboard accelerometer to quantify step count as part of activity monitoring. A minority of studies sought to assess the access of further care as part of their technology-based follow-up assessment^{19,49,69} - studies seeking this information were conducted solely in LMICs. One article describing the use of a telephone hotline for neurotrauma, by the nature of the technology differed greatly from others included by not reporting use of any outcome measures or other proforma for data collection⁵³.

Technology Evaluation

Where technology was evaluated, most studies employed generalized questionnaires developed internally, and often used visual analogue or 5- or 7-point Likert scales to gauge overall user satisfaction^{61,62,65,69-73,77} asking questions about acceptability, user friendliness, aesthetic, task difficulty, task appropriateness and general impression of the technology. A smaller set of studies assessed technology feasibility^{48,72,79} (such as retrospectively analyzing those eligible for FUT enrolment, those consented, and those that completed all instances of assessment, amongst reasons for loss of contact where realized), and compliance,^{72,73,77} often which was calculated retrospectively rather than evaluated by the patient

cohorts themselves, such as the use of a technology with respect to the investigator's target dose⁷². Further, two studies reported anecdotal or qualitative feedback pertaining to the patient's experience in using the technology.^{48,79} One study asked participants to log daily unexpected, technology-related events such as errors as part of the evaluation process, and their perceived accuracy of the GPS-based activity data with respect to their actual activities.⁶¹ Only one study utilized an externally validated assessment of telehealth services, the Telehealth Usability Questionnaire (TUQ).⁷³

Discussion

The purpose of this review was to describe the breadth of technologies implemented for follow-up, and highlight the instruments deployed with respect to their successes for all-severity TBI in a global setting. Forty-two studies were retrieved that utilized FUTs for symptom surveillance and outcome data collection and described technologies that fell under broad categories of telephone-, SMS-, smartphone-, videoconference-based technologies amongst a small number of miscellaneous devices that may fit under a number of these categories.

Smartphones are widely recognized to be both well-positioned and well-suited for emotional,⁷³ behavioural,⁸³ and physical monitoring,⁸⁴ particularly when applied to an ecological momentary assessment (EMA) or experience sampling method (ESM) methodologies. Traditional, face-to-face assessments conducted at infrequent intervals along the patient's journey of recovery rely upon retrospective self-reports that themselves are predisposed to recall bias,⁸⁵ which TBI patients are believed to be more susceptible to.⁸⁶ This is further compounded by the numerous challenges TBI patients face with cognitive impairment,⁸⁷ poor memory,⁸⁸ and impaired self-awareness.^{89,90} Smartphone and other FUT-based remote assessment, as mirrored in this review, shows promise to quantify the symptoms more accurately, and with respect to their temporal variability; otherwise uncaptured at a single time-point, and further facilitated in one's own natural environment.

The United Nations General Assembly in 2015 highlighted the impact technology-enabled breakthroughs have had in the healthcare sector, enabling greater numbers of people to have access to services otherwise out of reach or unaffordable.⁹¹ Whilst the implementation of technology to deliver follow-up is better than a complete absence of services, an informed understanding of the capabilities and technological fluency of the target population will be imperative for comprehensive and proper integration within standard practice. Several barriers to successful access of remote tele-health assessments and consultations have been identified in literature and in recent WHO 2019 guidelines,⁹²

including: disability such as hearing or cognitive impairment, lack of equipment, poor networking access and speeds, lack of organizational support, difficulty using the systems, security or privacy concerns, unfamiliarity with technology.⁹²⁻⁹⁵ One qualitative study exploring non-participant views of a wider telehealth and telecare trial depicted that some patients were hesitant to adopt new approaches where existing face-to-face services were often highly valued.⁹⁴ Patients with stigmatized health conditions may also possess additional concerns about the privacy of their information⁹² when handled digitally. Developing an awareness of the challenges faced by populations, particularly those most vulnerable such as the elderly and those recovering from the sequelae of injury, will ensure progress toward digital equality of service access, and mitigate the risk of bias or inaccurate data being introduced into technology-enhanced trials, registries, and injury surveillance campaigns.

Although the studies included often quantify their successes by compliance, adherence, and response rates, amongst broad and crude evaluations of these novel services, it is worth highlighting that any communication fostered by these technologies outside of pre-defined (and often sparsely scheduled) outpatient clinic meetings were appreciated and welcomed by both pediatric and adult patients and their caregivers, reflected by satisfaction rates reported in the studies included, and in part by strong compliance and return rates in the majority of FUT implementations in TBI. To our knowledge, no studies reported the use of a requirements elicitation survey or similar in the design and development of FUTs, and thus this is welcomed in future research in this area when examining what factors should be addressed and built into these services to improve patient compliance and satisfaction.

Further, the ‘successes’ of technology-based follow-up delivery (of which one may consider to be derived from compliance, attrition, or response metrics) do not appear to be reported in a standardized format across the studies retrieved. As a result, conducting systematic comparisons between technology modalities, and across cohorts of varying severity and demographics, remains a challenge. Whilst the majority of studies attempted to quantify success of their technology (Tables 1-5), such as comparing response rates versus prompts delivered, or the number of patients reached after an arbitrary number of contact attempts, further research or initiatives addressing this gap would serve favorably.

The compliance and satisfaction of clinicians remains key in developing technologies for integration into standard practice. It is not a new notion that neurosurgical services across the globe encounter heavy workloads and large patient numbers, especially in LMIC settings of whom must handle a disproportionate volume of cases in the face of smaller

specialist workforces when compared with their HIC colleagues⁹⁶. The benefits of asynchronous technologies thus become clear; enabling the clinician to review and act upon the data of their own accord, amongst providing patients with the ability to self-report at their discretion, and at more frequent instances otherwise impossible to fulfil synchronously by clinical staff.

Wearable technologies such as smart watches, biometric monitors, and smart clothing may offer additional asynchronous sources of data and are becoming slowly introduced as medical technologies, though were not identified in our review. Across medical disciplines, wearables have been demonstrated to enable real-time monitoring of vital signs, physical status and physiological parameters as patients go about their daily lives.⁹⁷ Although the literature describing wearables for TBI-afflicted individuals remains sparse,⁹⁸ we clearly envision the role these technologies may play in enabling additional remote, data-driven approaches for post-TBI monitoring and early sequelae management.⁹⁹

Lastly, it is of note that due to the global prevalence and availability of mobile phone or telephone services, many studies screened often briefly stated the use telephone for follow-up practice their reports, however, excluded a deeper insight, evaluation, or formal assessment of the utility of the remote technology for conducting the act of said follow-up itself. In this respect, due diligence was exercised by the authors across the screening stages.

This scoping review serves as a foundation for the application of technology in follow-up and outcome data collection. Herein, we propose a number of recommendations for future research and practice.

Firstly, from this review, there appears to be grounds for further research exploring, and perhaps refining, what outcome measures are deployed, and at what intervals assessments are conducted. Table 6 revealed a broad range of outcome measures used in the assessment of TBI patients, some of which were deployed in multiples. To ensure technology functions optimally and simultaneously for both patients and clinicians, a balance must be struck between the richness of assessments, and the efforts required on the patient's behalf, in quantifying their health and wellbeing at a distance. Further, it would be beneficial for future research to assess the validity of outcome measures when delivered sequentially in a single instance, and namely whether delivering multiple outcome measures introduces confounding effects, including when delivered remotely and without the option of clinical assistance or clarification. Similarly, it would be beneficial to explore and address the validities of outcome measures of which were initially designed to be deployed on

a face-to-face in-person basis, for when they have been applied to FUT-led research. We feel that there may be assumptions to challenge regarding whether if a party perceived an outcome measure could be successfully delivered by a technological medium with ease, that the validities of such measure must transfer simultaneously. As such, we encourage further efforts into FUT-led research practices themselves.

Secondly, a wide variety of outcome measures and instruments have been deployed by technology for remote assessment. Although beyond the scope of this review, it may be useful for further research to retrospectively address the validity of measures deployed, often designed for in-person use, with respect to their adaption for FUT deployment. Lastly, further investigation may be warranted to assess the effects on outcome measure validity in studies deploying multiple patient-reported outcome measures at single or close-together timepoints. Furthermore, such research may wish to examine whether these instruments when delivered individually or as sets remotely, can continue to accurately quantify outcomes from the acute to long-term stages of all-severity TBI and where these technologies may assist in facilitating data collection for common data elements as part of large-scale research efforts.

Strengths and Limitations

To our knowledge, this review is among the first to map the current global evidence base of technologies deployed to augment traditional modes of in-person follow-up. A broad and comprehensive search across five electronic databases was conducted, and as such this review serves as a strong foundation for understanding the use-cases of technology-based follow-up for TBI in a global setting. Whilst the number of studies retrieved is relatively small considering our eligibility criteria (FUTs employed in a global context for all-severity TBI in pediatric and adult cohorts), we present a particular slice of the evidence base in which included articles have a primary focus of exploring FUT deployment. In the context of future research, we trust this review will serve further as a comprehensive frame of reference for those wishing to apply FUTs in clinical practice or research. However, owing to a sheer number of terms describing global health, technologies, and their implementation, outcome measures, and follow-up, some published articles that may have met the pre-defined inclusion criteria could have been inadvertently excluded from this review. Further, we acknowledge limitations of our retrieved articles with respect to our eligibility criteria; we are aware of a number of large, high-profile injury surveillance studies, such as those utilizing the U.S. TBI Model Systems (TBIMS) database,¹⁰⁰⁻¹⁰³ that were not included, despite using FUTs in their research, as a description or assessment of FUTs was not the primary focus of such works. Similarly, by the nature of global health and technological reports, of which may not

always be confined to health journals and/or databases, nor always depicted in the English language, we are aware of the limitations in the literature retrieved, and of the literature that may exist in alternate academic or commercial domains and mediums. Considering these limitations, and with an understanding of the parameters of FUTs afforded by this review, a further systematic review as the evidence base evolves, inclusive of gray-literature, non-English publications, and articles utilizing FUTs but not necessarily as their primary aim, is recommended.

Conclusions

Our review has demonstrated that the evidence-base surrounding follow-up technologies remains in its infancy, particularly with respect to recruiting large patient cohorts, conducting a formal technology assessment, and the representation of research outside of high-income settings respectively. Of the use-cases described, incorporating technologies, both asynchronous and synchronous in nature, may leverage a clinician's abilities in gaining insights of the patient's well-being from discharge and beyond between traditional and often sparsely scheduled face-to-face appointments. FUTs may additionally serve to provide a more precise picture of TBI patient status through their ability to collect data at timepoints closer in proximity to in-person follow-up, harmonious with the WHO's adjuration and promotion of outcome data collection and injury surveillance in the reduction of global TBI burden. Further systematic analyses may prove useful in empirically quantifying the utility, acceptability, feasibility, and costs of each technology modality in neurotrauma practice. Future research may wish to characterize the challenges of implementing, sustaining, and adhering to these novel systems from the perspectives of patients, their proxies, and physicians.

Authorship confirmation/contribution statement

Brandon G. Smith: Conceptualization, Methodology, Formal analysis, Data Curation, Writing - Original Draft; **Stasa Tumpa:** Investigation, Writing - Review & Editing; **Orla Mantle:** Investigation; **Charlotte J. Whiffin:** Methodology, Writing - Review & Editing, Supervision; **Harry Mee:** Writing - Review & Editing; **Davi J. Fontoura Solla:** Writing - Review & Editing; **Wellingson Silva Paiva:** Writing - Review & Editing; **Virginia F J Newcombe:** Writing - Review & Editing; **Angelos G. Kolias:** Conceptualization, Methodology, Writing - Review & Editing, Supervision; **Peter J. Hutchinson:** Conceptualization, Writing - Review & Editing, Supervision.

Funding

The research was supported by the National Institute for Health Research (NIHR) Brain Injury MedTech Co-operative based at Cambridge University Hospitals NHS Foundation Trust and University of Cambridge, and by the National Institute for Health Research (NIHR) Global Health Research Group on Neurotrauma (grant number 16/137/105) using UK aid from the UK government.

Disclaimer

The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Disclosures

AGK and PJH are supported by the National Institute for Health Research (NIHR) Cambridge Biomedical Research Centre, the NIHR Brain Injury MedTech Co-operative, and the NIHR Global Health Research Group on Neurotrauma. BGS and CJW are supported by the NIHR Global Health Research Group on Neurotrauma. PJH is also supported by an NIHR Research Professorship, and by the Royal College of Surgeons of England. The NIHR Global Health Research Group on Neurotrauma was commissioned by the UK NIHR using Official Development Assistance funding (project no. 16/137/105). VFJN is funded by an Academy of Medical Sciences / The Health Foundation Clinician Scientist Fellowship.

References

1. Dewan MC, Rattani A, Gupta S, et al. Estimating the Global Incidence of Traumatic Brain Injury. *J Neurosurg* 2019;130(4):1080–1097; doi: 10.3171/2017.10.JNS17352.
2. Hyder AA, Wunderlich CA, Puvanachandra P, et al. The Impact of Traumatic Brain Injuries: A Global Perspective. Neufeld JA. ed. *NeuroRehabilitation* 2007;22(5):341–353; doi: 10.3233/NRE-2007-22502.
3. Langlois JA, Rutland-Brown W and Wald MM. The Epidemiology and Impact of Traumatic Brain Injury. *J Head Trauma Rehabil* 2006;21(5):375–378; doi: 10.1097/00001199-200609000-00001.
4. Bryant RA, O'Donnell ML, Creamer M, et al. The Psychiatric Sequelae of Traumatic Injury. *Am J Psychiatry* 2010;167(3):312–320; doi: 10.1176/appi.ajp.2009.09050617.
5. Maas AIR, Menon DK, Adelson PD, et al. Traumatic Brain Injury: Integrated Approaches to Improve Prevention, Clinical Care, and Research. *Lancet Neurol* 2017;16(12):987–1048; doi: 10.1016/S1474-4422(17)30371-X.
6. Hofman K, Primack A, Keusch G, et al. Addressing the Growing Burden of Trauma and Injury in Low- and Middle-Income Countries. *Am J Public Health* 2005;95(1):13–17; doi: 10.2105/AJPH.2004.039354.
7. Wesson HKH, Boikhutso N, Bachani AM, et al. The Cost of Injury and Trauma Care in Low- and Middle-Income Countries: A Review of Economic Evidence. *Health Policy Plan* 2014;29(6):795–808; doi: 10.1093/heapol/czt064.

8. Mock C, Lormand J, Goosen J, et al. Guidelines for Essential Trauma Care. Geneva, World Health Organization; 2004.
9. Juillard CJ, Mock C, Goosen J, et al. Establishing the Evidence Base for Trauma Quality Improvement: A Collaborative WHO-IATSIC Review. *World J Surg* 2009;33(5):1075–1086; doi: 10.1007/s00268-009-9959-8.
10. Laytin AD, Seyoum N, Azazh A, et al. Feasibility of Telephone-Administered Interviews to Evaluate Long-Term Outcomes of Trauma Patients in Urban Ethiopia. *Trauma Surg Acute Care Open* 2018;3(1):e000256; doi: 10.1136/tsaco-2018-000256.
11. Laytin AD and Debebe F. The Burden of Injury in Low-Income and Middle-Income Countries: Knowing What We Know, Recognising What We Don't Know. *Emerg Med J* 2019;emermed-2019-208514; doi: 10.1136/emered-2019-208514.
12. Holder Y, Peden M and Krug E. Injury Surveillance Guidelines. World Health Organisation: Geneva; 2001.
13. Mowafi H, Ngaruiya C, O'Reilly G, et al. Emergency Care Surveillance and Emergency Care Registries in Low-Income and Middle-Income Countries: Conceptual Challenges and Future Directions for Research. *BMJ Glob Health* 2019;4(Suppl 6):e001442; doi: 10.1136/bmjgh-2019-001442.
14. Agrawal A, Munivenkatappa A, Pal R, et al. Traumatic Brain Injury Registry: Sharing the Pilot Study Experience to Foster a Multicenter Project on Traumatic Brain Injury Core Data. *Indian J Neurotrauma* 2016;13(02):081–087; doi: 10.1055/s-0036-1586486.
15. Rubiano AM, Carney N, Chesnut R, et al. Global Neurotrauma Research Challenges and Opportunities. *Nature* 2015;527(7578):S193–S197; doi: 10.1038/nature16035.
16. Reilly P. The Impact of Neurotrauma on Society: An International Perspective. In: *Progress in Brain Research* Elsevier; 2007; pp. 3–9; doi: 10.1016/S0079-6123(06)61001-7.
17. De Silva MJ, Roberts I, Perel P, et al. Patient Outcome after Traumatic Brain Injury in High-, Middle- and Low-Income Countries: Analysis of Data on 8927 Patients in 46 Countries. *Int J Epidemiol* 2009;38(2):452–458; doi: 10.1093/ije/dyn189.
18. United Nations, Department of Economic and Social Affairs, and Population Division. *World Urbanization Prospects: The 2018 Revision*. 2019.
19. Vaca SD, Xu LW, Nalwanga J, et al. Long-Term Follow-up of Pediatric Head Trauma Patients Treated at Mulago National Referral Hospital in Uganda. *J Neurosurg Pediatr* 2019;23(1):125–132; doi: 10.3171/2018.7.PEDS17601.
20. Ganapathy K. Telemedicine and Neurosciences in Developing Countries. *Surg Neurol* 2002;58(6):388–394; doi: 10.1016/S0090-3019(02)00924-2.
21. Bagchi S. Telemedicine in Rural India. *PLoS Med* 2006;3(3):e82; doi: 10.1371/journal.pmed.0030082.
22. Turner B, Fleming J, Cornwell P, et al. A Qualitative Study of the Transition from Hospital to Home for Individuals with Acquired Brain Injury and Their Family Caregivers. *Brain Inj* 2007;21(11):1119–1130; doi: 10.1080/02699050701651678.
23. Thakar S, Rajagopal N, Mani S, et al. Comparison of Telemedicine with In-Person Care for Follow-up after Elective Neurosurgery: Results of a Cost-Effectiveness Analysis of 1200 Patients Using Patient-Perceived Utility Scores. *Neurosurg Focus* 2018;44(5):E17; doi: 10.3171/2018.2.FOCUS17543.
24. Dadlani R, Mani S, A U JG, et al. The Impact of Telemedicine in the Postoperative Care of the Neurosurgery Patient in an Outpatient Clinic: A Unique Perspective of This Valuable Resource in the Developing World—an Experience of More than 3000 Teleconsultations. *World Neurosurg* 2014;82(3–4):270–83; doi: 10.1016/j.wneu.2014.05.027.

25. Latulippe K, Hamel C and Giroux D. Social Health Inequalities and EHealth: A Literature Review With Qualitative Synthesis of Theoretical and Empirical Studies. *J Med Internet Res* 2017;19(4):e136; doi: 10.2196/jmir.6731.
26. Anonymous. Measuring Digital Development: Facts and Figures 2020. International Telecommunication Union (ITU): Geneva; 2020.
27. Bahia K and Delaporte A. The State of Mobile Internet Connectivity 2020. GSM Association; n.d.
28. Penn JW, Marcus HJ and Uff CEG. Fifth Generation Cellular Networks and Neurosurgery: A Narrative Review. *World Neurosurg* 2021;156:96–102; doi: 10.1016/j.wneu.2021.09.054.
29. Tricco AC, Lillie E, Zarin W, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* 2018;169(7):467–473; doi: 10.7326/M18-0850.
30. Aromataris E. Joanna Briggs Institute Reviewer’s Manual. The Joanna Briggs Institute; 2017.
31. Moher D, Liberati A, Tetzlaff J, et al. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *BMJ* 2009;339(jul21 1):b2535–b2535; doi: 10.1136/bmj.b2535.
32. Munn Z, Peters MDJ, Stern C, et al. Systematic Review or Scoping Review? Guidance for Authors When Choosing between a Systematic or Scoping Review Approach. *BMC Med Res Methodol* 2018;18(1):143; doi: 10.1186/s12874-018-0611-x.
33. Arksey H and O’Malley L. Scoping Studies: Towards a Methodological Framework. *Int J Soc Res Methodol* 2005;8(1):19–32; doi: 10.1080/1364557032000119616.
34. Levac D, Colquhoun H and O’Brien KK. Scoping Studies: Advancing the Methodology. *Implement Sci* 2010;5(1):69; doi: 10.1186/1748-5908-5-69.
35. Whiffin CJ, Smith BG, Selveindran SM, et al. The Value and Potential of Qualitative Research Methods in Neurosurgery. *World Neurosurg* 2021;S1878875021018842; doi: 10.1016/j.wneu.2021.12.040.
36. Anonymous. Joanna Briggs Institute Reviewers’ Manual: 2015 Edition / Supplement. 2015.
37. Aromataris E and Munn Z, (eds). *JBIM Manual for Evidence Synthesis*. JBI; 2020.; doi: 10.46658/JBIMES-20-01.
38. Bashshur R, Shannon G, Krupinski E, et al. The Taxonomy of Telemedicine. *Telemed E-Health* 2011;17(6):484–494; doi: 10.1089/tmj.2011.0103.
39. The World Bank. World Bank Country and Lending Groups (2021-2022). n.d. Available from: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>.
40. Arksey H and O’Malley L. Scoping Studies: Towards a Methodological Framework. *Int J Soc Res Methodol* 2005;8(1):19–32; doi: 10.1080/1364557032000119616.
41. Dombovy ML and Olek AC. Recovery and Rehabilitation Following Traumatic Brain Injury. *Brain Inj* 1997;11(5):305–18; doi: 10.1080/026990597123467.
42. Bell KR, Hoffman JM, Doctor JN, et al. Development of a Telephone Follow-up Program for Individuals Following Traumatic Brain Injury. *J Head Trauma Rehabil* 2004;19(6):502–12.
43. Smith B, Harms WD, Burres S, et al. Enhancing Behavioral Health Treatment and Crisis Management through Mobile Ecological Momentary Assessment and SMS Messaging. *Health Informatics J* 2012;18(4):294–308; doi: 10.1177/1460458212445349.

44. Xavier T, Robin M and Agrawal D. Using a Call Centre for Data Mining and Data Capturing - Pilot Study Using Glasgow Coma Outcome Scale in Head Injured Patients. *Stud Health Technol Inform* 2016;225(ck1, 9214582):866–7.
45. Varner C, McLeod SL, Nahiddi N, et al. Text Messaging Research Participants as a Follow-up Strategy to Decrease Emergency Department Study Attrition. *Can J Emerg Med* 2016;18(Supplement 1):S59; doi: 10.1017/cem.2016.121.
46. Deasy C and Underwood S. Telephone Follow-up of Mild Traumatic Brain Injury. *Brain Inj* 2016;30(5–6):808; doi: 10.3109/02699052.2016.1162060.
47. Thibault-Halman G, Fenerty L, Taylor P, et al. Early Telephone Follow-up for Traumatic Brain Injury Patients Using the Rivermead Post-Concussion Symptoms Questionnaire. *Can J Neurol Sci* 2017;44(Supplement 2):S17–S18; doi: 10.1017/cjn.2017.101.
48. Sutiono AB, Faried A, McAllister S, et al. The Bandung Neurosurgery Patient Outcomes Project, Indonesia (Part II): Patient Pathways and Feasibility and Acceptability of Telephone Follow-Up. *Int J Health Plann Manage* 2017;33(1):e49–e56; doi: 10.1002/hpm.2406.
49. Xu LW, Vaca SD, Nalwanga J, et al. Life After the Neurosurgical Ward in Sub-Saharan Africa: Neurosurgical Treatment and Outpatient Outcomes in Uganda. *World Neurosurg* 2018;113(101528275):e153–e160; doi: 10.1016/j.wneu.2018.01.204.
50. Underwood S, Campbell S and Deasy C. Telephone Follow-Up of Mild Traumatic Brain Injury; A Feasibility Study. *Ir Med J* 2019;112(8):984.
51. Juengst SB, Terhorst L, Kew CL, et al. Variability in Daily Self-Reported Emotional Symptoms and Fatigue Measured over Eight Weeks in Community Dwelling Individuals with Traumatic Brain Injury. Bell B Brown, Bryant, Carvalho, Colantonio, Crawford, Dicianno, Donders, Donker, Fann, Farace, Frazier, Grace, Iverson, John, Juengst, Juengst, Juengst, Juengst, Juengst, Juengst, Klonoff, Kroenke, Kroenke, Kumar, Kurtz, Ladwig, Macintyre, Neuberger, Perrin, Riggio, Sabaz, Schonberger, Schonberger, Shiffman, Shum, Spitzer, Squire, Srivastava, Stroup, Suffoletto, Terhorst, Terhorst, Terhorst, van Wijk, Wagner, Wagner, White, Worthen-Chaudhari, Zeger. ed. *Brain Inj* 2019;33(5):567–573; doi: 10.1080/02699052.2019.1584333.
52. Schlichter E, Lopez O, Scott R, et al. Feasibility of Nurse-Led Multidimensional Outcome Assessments in the Neuroscience Intensive Care Unit. *Crit Care Nurse* 2020;40(3):e1–e8; doi: 10.4037/ccn2020681.
53. Rhame K, Le D, Horner A, et al. Implementation of a Neurotrauma Hotline for Post-hospital Continuity of Care. *Acta Neurol Scand* 2020;141(4):351–354; doi: 10.1111/ane.13199.
54. Wiebe DJ, Nance ML, Houseknecht E, et al. Ecologic Momentary Assessment to Accomplish Real-Time Capture of Symptom Progression and the Physical and Cognitive Activities of Patients Daily Following Concussion. *JAMA Pediatr* 2016;170(11):1108; doi: 10.1001/jamapediatrics.2016.1979.
55. Wong AWK, Heinemann AW, Miskovic A, et al. Feasibility of Computerized Adaptive Testing for Collection of Patient-Reported Outcomes after Inpatient Rehabilitation. *Arch Phys Med Rehabil* 2014;95(5):882–91; doi: 10.1016/j.apmr.2013.12.024.
56. Anthony CA and Peterson AR. Utilization of a Text-Messaging Robot to Assess Intraday Variation in Concussion Symptom Severity Scores. *Clin J Sport Med Off J Can Acad Sport Med* 2015;25(2):149–52; doi: 10.1097/JSM.000000000000115.
57. Sy KT, Landau A and Dams-O’connor K. Feasibility of a Telephone-Administered Cognitive Test in Individuals with Moderate-Severe Tbi: A Tbims Study. *Arch Phys Med Rehabil* 2017;98(10):e8–e9; doi: 10.1016/j.apmr.2017.08.024.

58. Juengst S, Terhorst L, Arenth P, et al. Variability in Emotional Symptoms and Fatigue Measured via Mobile Ecological Momentary Assessment after TBI. *Arch Phys Med Rehabil* 2017;98(10):e130; doi: 10.1016/j.apmr.2017.08.424.
59. Pacella M, Prabhu A, Morley J, et al. Postconcussive Symptoms over the First 14 Days after Mild Traumatic Brain Injury: An Experience Sampling Study. *J Head Trauma Rehabil* 2018;33(3):E31–E39; doi: 10.1097/HTR.0000000000000335.
60. Ketchum J, Hart T, O’Neil-Pirozzi T, et al. Telephone Assessment of Cognitive Function Adds to Return to Work Prediction Following Traumatic Brain Injury. *Arch Phys Med Rehabil* 2019;100(10):e28–e29; doi: 10.1016/j.apmr.2019.08.069.
61. Wen P-S, Mackey J, Rose D, et al. Smartphone Application for Measuring Community Participation in Veterans With Traumatic Brain Injury. *OTJR Occup Particip Health* 2021;41(3):196–205; doi: 10.1177/15394492211004159.
62. Lenaert B, Colombi M, van Heugten C, et al. Exploring the Feasibility and Usability of the Experience Sampling Method to Examine the Daily Lives of Patients with Acquired Brain Injury. *Neuropsychol Rehabil* 2019;29(5):754–766; doi: 10.1080/09602011.2017.1330214.
63. Lewandowski L, Rieger B, Smyth J, et al. Measuring Post-Concussion Symptoms in Adolescents: Feasibility of Ecological Momentary Assessment. *Arch Clin Neuropsychol* 2009;24(8):791–796; doi: 10.1093/arclin/acp087.
64. Warden DL, Salazar AM, Martin EM, et al. A Home Program of Rehabilitation for Moderately Severe Traumatic Brain Injury Patients. The DVHIP Study Group. *J Head Trauma Rehabil* 2000;15(5):1092–102.
65. Huijgen BCH, Vollenbroek-Hutten MMR, Zampolini M, et al. Feasibility of a Home-Based Telerehabilitation System Compared to Usual Care: Arm/Hand Function in Patients with Stroke, Traumatic Brain Injury and Multiple Sclerosis. *J Telemed Telecare* 2008;14(5):249–56; doi: 10.1258/jtt.2008.080104.
66. Suffoletto B, Wagner AK, Arenth PM, et al. Mobile Phone Text Messaging to Assess Symptoms after Mild Traumatic Brain Injury and Provide Self-Care Support: A Pilot Study. *J Head Trauma Rehabil* 2013;28(4):302–12; doi: 10.1097/HTR.0b013e3182847468.
67. Pavliscsak H, Little JR, Poropatich RK, et al. Assessment of Patient Engagement with a Mobile Application among Service Members in Transition. *J Am Med Inform Assoc JAMIA* 2016;23(1):110–8; doi: 10.1093/jamia/ocv121.
68. Mortenson P, Singhal A, Hengel AR, et al. Impact of Early Follow-up Intervention on Parent-Reported Postconcussion Pediatric Symptoms: A Feasibility Study. *J Head Trauma Rehabil* 2016;31(6):E23–E32; doi: 10.1097/HTR.0000000000000223.
69. Shahrokhi A, Azimian J, Amouzegar A, et al. The Effect of Telenursing on Referral Rates of Patients With Head Trauma and Their Family’s Satisfaction After Discharge. *J Trauma Nurs Off J Soc Trauma Nurses* 2018;25(4):248–253; doi: 10.1097/JTN.0000000000000382.
70. Hill AJ, Theodoros D, Russell T, et al. Using Telerehabilitation to Assess Apraxia of Speech in Adults. *Int J Lang Commun Disord* 2009;44(5):731–747; doi: 10.1080/13682820802350537.
71. Hill AJ, Theodoros DG, Russell TG, et al. The Redesign and Re-Evaluation of an Internet-Based Telerehabilitation System for the Assessment of Dysarthria in Adults. *Telemed E-Health* 2009;15(9):840–850; doi: 10.1089/tmj.2009.0015.
72. Worthen-Chaudhari L, McGonigal J, Logan K, et al. Reducing Concussion Symptoms among Teenage Youth: Evaluation of a Mobile Health App. *Brain Inj* 2017;31(10):1279–1286; doi: 10.1080/02699052.2017.1332388.

73. Juengst SB, Graham KM, Pulantara IW, et al. Pilot Feasibility of an MHealth System for Conducting Ecological Momentary Assessment of Mood-Related Symptoms Following Traumatic Brain Injury. *Brain Inj* 2015;29(11):1351–61; doi: 10.3109/02699052.2015.1045031.
74. Cuthbert JP, Whiteneck GG, Corrigan JD, et al. The Reliability of a Computer-Assisted Telephone Interview Version of the Ohio State University Traumatic Brain Injury Identification Method. *J Head Trauma Rehabil* 2016;31(1):E36–E42; doi: 10.1097/HTR.000000000000075.
75. Rietdijk R, Power E, Brunner M, et al. Reliability of Videoconferencing Administration of a Communication Questionnaire to People With Traumatic Brain Injury and Their Close Others. *J Head Trauma Rehabil* 2017;32(6):E38–E44; doi: 10.1097/HTR.000000000000282.
76. Licona N, Adamson M, Santiago R, et al. Reliability of Telephonic Neuropsychological Assessment Post Traumatic Brain Injury. *Arch Phys Med Rehabil* 2017;98(10):e102–e103; doi: 10.1016/j.apmr.2017.08.330.
77. Graham K, Terhorst L, Skidmore E, et al. Preliminary Feasibility of Smartphone Technology to Assess Participation Following Traumatic Brain Injury. *Brain Inj* 2017;31(6–7):846; doi: 10.1080/02699052.2017.1312145.
78. Sufrinko AM, Howie EK, Charek DB, et al. Mobile Ecological Momentary Assessment of Postconcussion Symptoms and Recovery Outcomes. *J Head Trauma Rehabil* 2019;34(6):E40–E48; doi: 10.1097/HTR.000000000000474.
79. McAllister S, Ganefianty A, Faried A, et al. The Bandung Neurosurgery Patient Outcomes Project, Indonesia (Part I): Methods, Participant Characteristics, and Pre-Discharge Outcomes. *Int J Health Plann Manage* 2017;33(1):e57–e66; doi: 10.1002/hpm.2408.
80. Wilson JTL, Edwards P, Fiddes H, et al. Reliability of Postal Questionnaires for the Glasgow Outcome Scale. *J Neurotrauma* 2002;19(9):999–1005.
81. Hellawell DJ, Signorini DF and Pentland B. Reliability of the Relative’s Questionnaire for Assessment of Outcome after Brain Injury. *Disabil Rehabil* 2000;22(10):446–50.
82. Vaca SD, Xu LW, Nalwanga J, et al. Long-Term Follow-up of Pediatric Head Trauma Patients Treated at Mulago National Referral Hospital in Uganda. *J Neurosurg Pediatr* 2019;23(1):125–132; doi: 10.3171/2018.7.PEDS17601.
83. Porras-Segovia A, Molina-Madueño RM, Berrouiguet S, et al. Smartphone-Based Ecological Momentary Assessment (EMA) in Psychiatric Patients and Student Controls: A Real-World Feasibility Study. *J Affect Disord* 2020;274:733–741; doi: 10.1016/j.jad.2020.05.067.
84. Nabasny A, Mishra S, Khetsavanh L, et al. Compliance Using Mobile Health Heart Rate Variability and Ecological Momentary Assessment Tools in a Chronic Traumatic Brain Injury Sample. *Arch Phys Med Rehabil* 2021;102(10):e70–e71; doi: 10.1016/j.apmr.2021.07.680.
85. Shiffman S, Stone AA and Hufford MR. Ecological Momentary Assessment. *Annu Rev Clin Psychol* 2008;4(1):1–32; doi: 10.1146/annurev.clinpsy.3.022806.091415.
86. Haagsma JA, Spronk I, de Jongh MAC, et al. Conventional and Retrospective Change in Health-Related Quality of Life of Trauma Patients: An Explorative Observational Follow-up Study. *Health Qual Life Outcomes* 2020;18(1):157; doi: 10.1186/s12955-020-01404-1.
87. McInnes K, Friesen CL, MacKenzie DE, et al. Mild Traumatic Brain Injury (MTBI) and Chronic Cognitive Impairment: A Scoping Review. Kobeissy FH. ed. *PLOS ONE* 2017;12(4):e0174847; doi: 10.1371/journal.pone.0174847.
88. Vakil E. The Effect of Moderate to Severe Traumatic Brain Injury (TBI) on Different Aspects of Memory: A Selective Review. *J Clin Exp Neuropsychol* 2005;27(8):977–1021; doi: 10.1080/13803390490919245.

89. Robertson K and Schmitter-Edgecombe M. Self-Awareness and Traumatic Brain Injury Outcome. *Brain Inj* 2015;29(7–8):848–858; doi: 10.3109/02699052.2015.1005135.
90. Prigatano GP and Sherer M. Impaired Self-Awareness and Denial During the Postacute Phases After Moderate to Severe Traumatic Brain Injury. *Front Psychol* 2020;11:1569; doi: 10.3389/fpsyg.2020.01569.
91. United Nations General Assembly. 70/125. Outcome Document of the High-Level Meeting of the General Assembly on the Overall Review of the Implementation of the Outcomes of the World Summit on the Information Society. United Nations; 2016.
92. World Health Organization. WHO Guideline :Recommendations on Digital Interventions for Health System Strengthening: Evidence and Recommendations. 2019.
93. Almathami HKY, Win KT and Vlahu-Gjorgievska E. Barriers and Facilitators That Influence Telemedicine-Based, Real-Time, Online Consultation at Patients’ Homes: Systematic Literature Review. *J Med Internet Res* 2020;22(2):e16407; doi: 10.2196/16407.
94. Sanders C, Rogers A, Bowen R, et al. Exploring Barriers to Participation and Adoption of Telehealth and Telecare within the Whole System Demonstrator Trial: A Qualitative Study. *BMC Health Serv Res* 2012;12(1):220; doi: 10.1186/1472-6963-12-220.
95. Blandford A, Wesson J, Amalberti R, et al. Opportunities and Challenges for Telehealth within, and beyond, a Pandemic. *Lancet Glob Health* 2020;8(11):e1364–e1365; doi: 10.1016/S2214-109X(20)30362-4.
96. Whiffin CJ, Smith BG, Esene IN, et al. Neurosurgeons’ Experiences of Conducting and Disseminating Clinical Research in Low-Income and Middle-Income Countries: A Reflexive Thematic Analysis. *BMJ Open* 2021;11(9):e051806; doi: 10.1136/bmjopen-2021-051806.
97. Yetisen AK, Martinez-Hurtado JL, Ünal B, et al. Wearables in Medicine. *Adv Mater* 2018;30(33):1706910; doi: 10.1002/adma.201706910.
98. Veerubhotla A, Krantz A, Ibironke O, et al. Wearable Devices for Tracking Physical Activity in the Community after an Acquired Brain Injury: A Systematic Review. *PM&R* 2021;pmrj.12725; doi: 10.1002/pmrj.12725.
99. Schmid W, Fan Y, Chi T, et al. Review of Wearable Technologies and Machine Learning Methodologies for Systematic Detection of Mild Traumatic Brain Injuries. *J Neural Eng* 2021;18(4):041006; doi: 10.1088/1741-2552/ac1982.
100. DiSanto D, Kumar RG, Juengst SB, et al. Employment Stability in the First 5 Years After Moderate-to-Severe Traumatic Brain Injury. *Arch Phys Med Rehabil* 2019;100(3):412–421; doi: 10.1016/j.apmr.2018.06.022.
101. Hammond FM, Perkins SM, Corrigan JD, et al. Functional Change from Five to Fifteen Years after Traumatic Brain Injury. *J Neurotrauma* 2021;38(7):858–869; doi: 10.1089/neu.2020.7287.
102. Marquez de la Plata CD, Hart T, Hammond FM, et al. Impact of Age on Long-Term Recovery From Traumatic Brain Injury. *Arch Phys Med Rehabil* 2008;89(5):896–903; doi: 10.1016/j.apmr.2007.12.030.
103. Hammond FM, Corrigan JD, Ketchum JM, et al. Prevalence of Medical and Psychiatric Comorbidities Following Traumatic Brain Injury. *J Head Trauma Rehabil* 2019;34(4):E1–E10; doi: 10.1097/HTR.0000000000000465.

Table 1 - Citations reporting the use of Telephone-based Follow-up

Author, Year (Country)	Study Design (Author definition)	Study Aim / Objective	Sample Population Demographics TBI Characteristics	Follow-up Technology (FUT) Description Clinical Facilitator Sessions & Instances count	Synchronicity Use of Proxy	Constructs & Outcome Measures Deployed	Response / Success / Compliance Rates
Dombovy et al., 1997 (USA) ⁴¹	Descriptive	To determine if functional, neuropsychological and social outcome at 3 and 6 months in patients hospitalized following TBI could be ascertained via telephone follow-up, and assess use of rehabilitation services in this population.	<p>n = 74 Adult TBI patients (all-severity) at home/community</p> <p>Average Age (SD) = 39.2y</p> <p>Sex (F) = 29.7%</p> <p>Mean Admission GCS = 11.5</p> <p>Mild or Moderate = 77%</p> <p>Severe = 23%</p> <p>TSI/D = 3 months +/- 2 weeks</p>	<p>Telephone-based assessment at 3- and 6-months post-injury</p> <p>Nurse practitioner</p> <p>2 sessions, 15-30 minutes in duration</p>	Synchronous No reported use of proxy	<p>Functional Independence Measure (FIM)</p> <p>Telephone Interview for Cognitive Status (TICS)</p> <p>Neurobehavioral Rating Scale (NRS)</p> <p>Generic questions regarding employment, household activities, personal finance management, travel & social activities</p>	Telephone deemed a cost-effective way to ascertain functional and neuropsychological outcomes in TBI survivors, and may identify those whom may benefit from additional rehabilitation
Warden et al., 2000 (USA) ⁶⁴	Experimental (Non-randomized, open label, controlled)	To compare home versus inpatient cognitive rehabilitation for patients with moderate to severe head injury.	<p>n = 53 Adult (military) TBI patients (moderate-severe) at home/community in</p>	Telephone-based support (information providing, problem solving, support and encouragement) and assessment	Synchronous Family members able to contact nurse as required	Weekly generalised overall wellbeing checklist (headache, irritability, fatigue, depression, memory problems,	n = 47 (88.7%) completed the telephone-based program

			<p>home-program arm of trial</p> <p>Average Age (SD) = 26y (6.22)</p> <p>Sex (F) = 4%</p> <p>Mean Admission GCS = 9.5 TSI/D (SD) = 39days (33.2)</p>	<p>Psychiatric Nurse</p> <p>Weekly sessions over 8 weeks</p>		<p>medication compliance, miscellaneous problems requiring intervention)</p> <p>Activities at week 2 and 7 (shopping, watching television, community activities and socializing with friends)</p>	
<p>Bell et al., 2004 (USA)⁴²</p>	<p>Descriptive</p>	<p>To describe the development of a telephone follow-up program that addresses the needs of survivors of traumatic brain injury (TBI) and their families in the year following injury.</p>	<p>n = 84 Adult (moderate-severe) TBI patients at home/community</p> <p>Average Age (SD) = 34.4 (13.6)</p> <p>Sex (F) = 18%</p> <p>Moderate to Severe TSI/D = 2 weeks following discharge</p>	<p>Telephone-based assessment interview and provision of support and information</p> <p>Research Care Manager</p> <p>7 planned contacts at 2wks, 4wks, 2mo, 3mo, 5mo, 7mo, 9mo</p>	<p>Synchronous</p> <p>Use of proxy at each contact (family member or significant other)</p>	<p>Non-specific review of past and current concerns with triage/referral as required.</p> <p>At 4 weeks and 9 months, structured interview addressing 17 domains: personal care, ambulation, travel, work, school, home management, leisure, social integration, cognitive and behavioural concerns, standard of living, financial independence, medical concerns, emotional function,</p>	<p>Median 4 contacts with patients and 4.5 with relatives over 9 month period.</p> <p>Median call duration 34 minutes with TBI patients, 30 minutes with relatives.</p>

						alcohol use, drug use, legal issues and spirituality.	
Wong et al., 2014 (USA) ⁵⁵	Observational (prospective cohort study)	To evaluate the feasibility of computer adaptive testing (CAT) using an Internet or telephone interface to collect patient-reported outcomes after inpatient rehabilitation and to examine patient characteristics associated with completion of the CAT-administered measure and mode of administration.	n = 674 (mixed diagnosis: spinal cord injury, stroke, orthopaedic, 'other neurologic' or other conditions), of which n = 40 Adult brain injury patients at home/community Total cohort: Average Age (SD) = 62.9 (15.7) Sex (F) = 59.8% Not defined	Telephone-based (interactive voice response) or secure internet website-based self-reported computerized adaptive testing Hospital outcome management department staff One instance of delivery a month following discharge	Asynchronous No reported use of proxy	Community Participation Indicators (CPI) modified for computer adaptive testing (CAT) delivery	Across the diagnosis cohorts, 61.0% chose telephone IVR versus 39.0% choosing internet-based assessment. Patients with brain injury had an 81% reduced likelihood of competing any form of assessment with respect to other patient cohorts.
Xavier et al., 2016 (India) ⁴⁴	Descriptive	To assess the Glasgow coma outcome scale via a call center for head injured patients who were discharged after head injury.	n = 484 TBI patients at home/community Not reported Not reported	External call centre with integration of electronic medical records system to administer structured telephonic questionnaire interview to discharged patients Call Centre Staff Single instance of administration following discharge	Synchronous No reported use of proxy	Glasgow Outcome Scale (GOS) in Hindi	84% (n = 406) patients could be reached by phone and GOS elicited. 63% of patients were from rural areas.
Mortenson et al., 2016 (Canada) ⁶⁸	Experimental (Pilot RCT)	To investigate the effectiveness and feasibility of early	n = 66 paediatric patients with concussion injury	Structured telephone-based follow-up and symptom counselling	Synchronous	Intervention arm at 1 wk/1mo:	Across both groups, n = 8 children were identified to have

		intervention telephone counselling with parents in limiting post-concussion symptoms and impacts on children and youth.	and their parents at home/community (n = 32 intervention, n = 34 control) Average Age (range) = 11.9 (6.3-16.5)[I], 12.6 (5.2-16.8) [C] Sex (F) = 25% [I], 35% [C] Clinically diagnosed mild TBI or concussion, defined as GCS 13/14 at admission or LoC <30min or altered mental status at injury or post-traumatic amnesia <24h duration TSI/D = Within one week of injury	for parents with children with ongoing symptoms Occupational Therapist Two instances at 1 week and 1-month post-injury in intervention group. Intervention and control received telephonic PCSI at 3 months post-injury.	Full use of proxy. No patients directly communicated with.	Acute Concussion Evaluation (ACE) Protocol. Generalized interview regarding impact of symptoms on everyday function, home, educational and play activities. Both groups at 3mo: Post-Concussion Symptom Inventory (PCSI) Parent Assessment Form	ongoing symptoms not otherwise identified by regular care at 3mo post-injury. No significant difference of post-concussion symptoms between groups at 3 months post-injury. n = 8 lost to attrition (loss of contact) across both groups from initial randomization of n = 76 patients.
Cuthbert et al., 2016 (USA) ⁷⁴	Quasi-experimental (test-retest of stratified random sample from larger primary study)	To provide test-retest reliability (>5 months) of the Ohio State University Traumatic Brain Injury Identification Method modified for use as a computer-assisted telephone interview (CATI) to capture traumatic brain	n = 200 Adult patients (n = 50 TBI with LoC, n = 50 TBI w/o LoC, n = 50 major lifetime injury w/o TBI, n = 50 no major lifetime injury) at home/community n = 194: Median Age group (range) = 50-59 (18-89).	Standardized, computer-assisted telephone interview (CATI) Interviewer (undefined) without familiarity with TBI One instance of telephone delivery 6 to 18 months following initial interview	Synchronous No reported use of proxy	Ohio State University TBI Identification Method (OSU TBI-ID)	100% (n = 194) instances of CATI completed across 18 months (range 6 - 15mo). Mean time between initial and telephone follow-up interviews = 10.96mo (SD 2.37). High response and completion rate

		injury (TBI) and other substantial bodily injuries among a representative sample of noninstitutionalized adults living in Colorado.	<p>Sex (F) = 30.9%</p> <p>All-severity TBI TSI/D (n = 194) = <5yrs since injury n = 26, 5<yrs since injury n = 168.</p>				<p>reported, with 68.3% of patients reached within 5 contact attempts.</p> <p>Average delivery time = 5 minutes (range 1 - 18 minutes).</p>	
Varner et al., 2016 (Canada) ⁴⁵	Descriptive (secondary data analysis of RCT data)	To determine if text messaging study participants involved in an ongoing randomized trial resulted in a lower rate of attrition as compared to conventional telephone follow-up.	<p>n = 118 (n = 40 intervention, n = 78 control) Adult head injury patients at home/community</p> <p>Average Age (SD) = 35.2y (13.7)</p> <p>Sex (F) = 63.6%</p> <p>Presenting ED complaint of 'head injury' not otherwise quantified/defined TSI/D = 2-4wks post-ED discharge</p>	<p>Telephone-based symptom questionnaire with (intervention)/without (control) prior SMS reminder</p> <p>No defined facilitator</p> <p>Two instances of telephone follow-up at 2 and 4wks post-ED discharge</p>	Synchronous	No reported use of proxy	Unspecified general symptom questionnaire	<p>n = 3 withdrew from control.</p> <p>Of n = 75 receiving telephone follow-up w/o reminder, 32% at 2 wks and 42.7% at 4 wks were unable to be contacted.</p> <p>Of n = 40 with SMS reminder prior to telephone follow-up, 10% at 2wks and 25% at 4wks were unable to be contacted.</p>
Deasy et al., 2016 (Ireland) ⁴⁶	Descriptive	To investigate the prevalence of PCS and the quality-of-life of patients who were treated in the Clinical Decision Unit of Cork University Hospital in 2013.	<p>n = 112 Adult TBI patients at home/community</p> <p>Not reported</p> <p>Mild TBI confirmed by records and/or CT imaging review.</p>	<p>Structured telephone-based follow-up</p> <p>No defined facilitator</p> <p>One instance of follow-up within 1-year post-injury</p>	Synchronous	No reported use of proxy	<p>Rivermead Post-concussion Symptoms Questionnaire (RPQ)</p> <p>Quality of Life by 12-Item Short Form Survey (SF-12)</p> <p>Functional outcome (EQ5D5L)</p>	Significant loss to follow-up with only 50.9% able to be reached within 1 year post injury.

			TSI/D = Approximately within 1-year post- injury				
Thibault-Halman et al., 2017 (Canada) ⁴⁷	Descriptive	To examine the frequency and severity of common post-TBI symptoms, as assessed by the Rivermead Post-Concussion Symptoms Questionnaire (RPCQ).	n = 46 Adult TBI patients at home/community Not reported Mild or moderate TBI TSI/D = <2wks following hospital discharge	Telephone-based questionnaire administration Rehab-based nurse practitioner One instance of follow-up at 2 weeks following inpatient discharge	Synchronous No reported use of proxy	Rivermead Post-concussion Symptoms Questionnaire (RPQ)	100% of patients were reached, where post-TBI symptoms in at least one domain (emotional, somatic, cognitive) remained present in 100% of cases Referral for additional formal assessment, Symptom management, and advice was provided in 37% of cases
Sy et al., 2017 (USA) ⁵⁷	Observational (cross sectional and longitudinal analysis as part of wider cohort study)	To evaluate feasibility of a multidimensional telephone-administered cognitive test in individuals with moderate-severe traumatic brain injury (TBI).	n = 463 (1yr) and n = 386 (2yr) Adult TBI patients at home/community Not reported Moderate to severe TBI TSI/D = Up to 2 years post-injury	Telephone-based questionnaire administration No defined facilitator Two instances of delivery a year apart	Synchronous No reported use of proxy	Brief Test of Adult Cognition by Telephone (BTACTION)	Of the participants independently completing the questionnaire (Year 1 = 60%, Year 2 = 62%) completion rates ranged from 83% to 88%. Of the entire sample, completion rates ranged from 60% to 70% for Year 1, and 56% to 64% for year 2.

							Completion rates in lower in participants tested in Spanish (39% to 69%).
Licona et al., 2017 (USA) ⁷⁶	Quasi-experimental (test-retest)	To evaluate neuropsychological assessments by telephone on patients with mild-severe TBI to facilitate follow-up evaluations and research studies when in person assessment is not feasible.	<p>n = 21 Adult (military and veteran) TBI patients at a polytrauma rehabilitation center</p> <p>Average Age (range) = 49y (31-71)</p> <p>Sex (F) = 9%</p> <p>All-severity TBI (Mild n = 7, Moderate n = 5, Severe n = 11)</p> <p>TSI/D = Within 6mo post-injury</p>	<p>Telephone-based neuropsychological assessment</p> <p>No defined facilitator</p> <p>Two instances several weeks apart (median 15 days, range 7-62 days)</p>	<p>Synchronous</p> <p>No reported use of proxy</p>	<p>Neuropsychological assessment battery (including standard verbally administered tests of attention, working memory, processing speed, language, memory, executive skills, and auditory-verbal adaptations of trail-making)</p>	<p>43% (n = 10) completed all 17 tests.</p> <p>91% (n = 21) completed 15 tests.</p> <p>Telephone testing providing reliable scores across multiple domains even in patients with significant deficits, described as useful for those who find it difficult to travel.</p>
Sutiono et al. ♡, 2017 (Indonesia) ^{48,79}	Descriptive	To describe the patients' pathways into RSHS, the pathways following discharge, and the feasibility of following up this patient population by telecommunication.	<p>n = 178 Adult neurosurgical patients (n = 104 TBI patients) at home/community</p> <p>n = 217:</p> <p>Average Age (SD, range) = 41 (14.6, 18-84)</p> <p>Sex (F) = 41%</p> <p>All-severity</p>	<p>Telephone-based follow-up assessment</p> <p>Dedicated neurosurgery nurse</p> <p>Three instances at 1, 2 and 3mo following hospital discharge</p>	<p>Synchronous</p> <p>Use of Proxy (designated family member) in some instances to initiate contact with patient</p>	<p>Health-related quality of life (EQ5D5L) at each instance</p> <p>Glasgow Outcome Scale Extended (GOS-E) at 3mo</p> <p>Technology evaluation: Retrospective analysis for feasibility by recording numbers</p>	<p>Despite difficulty, all patients were able to be reached with no drop-outs.</p> <p>55% of patients answered the phone on first contact whilst 42% required between 2 and 5 attempts before contact was made.</p> <p>All but one patient owned regular cell</p>

			TSI/D = Within 3mo of discharge			admitted / eligible / consented / able to be followed up amongst reasons for loss of contact Anecdotal evaluation of telephonic follow-up experience	phones. One patient owned a smartphone allowing assessment by videoconference. For 3%, 5+ contact attempts were made. Patients were happy to be contacted by telephone due to support provided and opportunity to ask condition-related questions.
Shahrokhi et al., 2018 (Iran) ⁶⁹	Experimental (RCT)	To assess the effect of telenursing on referral rates of patients with head trauma and their family's satisfaction after discharge.	n = 72 (n = 35 intervention, n = 33 control following n = 4 exclusion) Adult TBI patients at home/community n = 68: Average Age (SD) = 34.11y (12.34)[I], 31.12 (10.83)[C]. Sex (F) = 26.5% Mild to moderate 'Head injury' (GCS 11 - 15 at admission) TSI/D = From 12wks of discharge	Intervention group: Telephone-based caregiver-reported patient status checklist, with telenurse available at any time Telenurse Intervention group: One instance per week for 12 weeks, with caregiver able to contact telenurse as desired. Control group: One instance at 12 weeks.	Synchronous Full use of proxy. No patients directly communicated with	Generalized patient status checklist for caregiver, including demographics and characteristics, outcomes of care (e.g. Readmission, referrals pressure ulcers). Cause of caregiver calls to telenursing also reported. Technology evaluation: Satisfaction with telenursing service.	n = 4 excluded (n = 1 intervention, n = 3 control) for consecutive 3-week non-response or where home nursing services used 53.8% of caregivers satisfied with telenursing program Telenursing program resulted in statistically significant less referrals to physicians (25.7%) versus control group (39.4%)

<p>Xu et al., 2018 (Uganda)⁴⁹</p>	<p>Descriptive</p>	<p>To describe the use of phone surveys developed and conducted in the 40 participant's language to assess mortality, neurological outcomes, and follow-up healthcare.</p>	<p>n = 1167 Adult and pediatric patients with mixed neurosurgical pathology (TBI, spina bifida, tumor, hydrocephalus and miscellaneous) at home/community</p> <p>Of those surveyed (n = 870), n = 740 (85%) with TBI</p> <p>n = 870: Median Age = 26y Sex (F) = 19%</p> <p>n = 596: Median GOS-E = 8 (GR+) Mean GOS-E = 6.83 (GR-) Median TSI/D = 1.53 years</p>	<p>Telephone-based survey administration with prospective record electronic database</p> <p>Research assistants in patient's language</p> <p>One instance of follow-up (average duration 20 minutes)</p>	<p>Synchronous</p> <p>Use of Proxy (designated family member) in some instances to initiate contact with patient or collection of demographics</p>	<p>Glasgow Outcome Scale Extended (GOS-E) or pediatric version (GOSE-peds)</p> <p>General survey items pertaining to: Quality of life (continuing physical deficits), Activities of Daily Living (ADLs), ability to speak and follow commands, perform household chores, school and work function, psychosocial function, and subjective return to baseline functional status.</p> <p>Additional follow-up healthcare accessed also assessed.</p>	<p>Utilizing telephone, there was a 74.5% response rate (n = 870).</p> <p>Of those reached, no patient refused telephone assessment.</p> <p>70% of those that survived pre-discharged (n = 1167) had a phone number on file.</p>
<p>Laytin et al., 2018 (Ethiopia)¹⁰</p>	<p>Observational (retrospective cohort study)</p>	<p>To assess the feasibility of telephone-administered interviews as means of collecting follow-up data in this context; to pilot a telephone-</p>	<p>n = 397 Adult mixed trauma patients (n = 111 neurologic injury) at home/community</p> <p>n = 397: Average Age (SD) = 32.8 (14.8)</p>	<p>Structured telephone-administered interviews</p> <p>Data clerk</p> <p>One instance of delivery at 6 months post-discharge</p>	<p>Synchronous</p> <p>Use of proxy ('surrogate' - relative, friend, caretaker) where patient unable to respond themselves</p>	<p>Glasgow Outcome Scale Extended (GOS-E)</p>	<p>Over half (n = 208) of initially identified patients (n = 397) were unable to be reached due to telephone contact details being emergency contacts and/or emergency bystanders, or were</p>

		administered interview tool for collecting data about long-term functional outcomes after injury; and to collect preliminary data about patients' long-term functional outcomes after hospital encounters due to injury.	<p>Sex (F) = 16.1%</p> <p>Not reported</p> <p>TSI/D = Within 6mo of discharge</p>				<p>believed to not answer due to not recognizing the phone number.</p> <p>Formal reasons recorded, where available in n = 131 (63%), included: n = 92 (72%) not having a valid number on file, n = 37 (28%) with a telephone line out of service or not answered on 3 attempts, and n = 2 (2%) of contacts made with someone who did not know the current condition of the subject or how to contact them.</p>
<p>Vaca et al., 2018 (Uganda)¹⁹</p>	Descriptive	To describe the use of a novel method of telephone surveys to conduct the first-ever long-term follow-up in Uganda to elucidate the outcomes of pediatric head trauma patients treated at the national referral hospital.	<p>n = 142 paediatric TBI patients and their caregivers at home/community</p> <p>Median Age (range) = 6y (0.17 - 17)</p> <p>Sex (F) = 29%</p> <p>All-severity TBI</p> <p>Admission GCS = Mild 54%,</p>	<p>Structured telephone survey</p> <p>Ugandan research collaborator</p> <p>One instance of follow-up at either 1y or 2y from discharge</p>	Synchronous	<p>Full use of proxy (pediatric patient's caregiver)</p> <p>Glasgow Outcome Scale Extended Pediatric Revision (GOSE-Peds)</p> <p>Quality of Life (physical and psychosocial deficits, ability to carry out ADLs)</p> <p>Further care sought since discharge, mortality</p>	<p>Average call duration 20 minutes</p> <p>With up to 5 contact attempts, achieved a 61% response rate (of initial n = 232 patients identified), representing 67% of patients receiving treatment and discharge in 12 months with a</p>

			Moderate 31%, Severe 15% TSI/D (median) = 1.48y				phone number on file Suggested as a suitable alternative for home visits for a large referral hospital
Underwood et al., 2019 (Ireland)⁵⁰	Descriptive	To investigate the prevalence of Post-Concussion Syndrome (PCS) one-year post-injury in patients that were treated for Mild Traumatic Brain Injury (mTBI) in the Clinical Decision Unit (CDU) of Cork University Hospital's (CUH) Emergency Department.	n = 57 Adult TBI patients at home/community Median Age (range) = 40y (27.5 - 57.5) Sex (F) = 42.1% Mild TBI TSI/D = 1-year post-discharge	Structured telephone assessment No defined facilitator One instance of follow-up at 1 year post-discharge	Synchronous No reported use of proxy	Rivermead Post-concussion Symptoms Questionnaire (RPQ) Short Form 12-item Health Survey (SF-12) EuroQol 5 Dimension 5 Level Outcome Assessment Tool (EQ5D5L)	51% response rate (n = 57) of initial n = 112 attempted after 4 attempts
Ketchum et al., 2019 (USA)⁶⁰	Observational (prospective cross-sectional cohort)	To assess the contribution of a brief telephone assessment of cognitive function on prediction of return to work at 1 year following moderate to severe traumatic brain injury (TBI).	n = 320 Adult TBI patients at home/community Age Range = 18 - 64y Sex (F) = Not reported Moderate to Severe TBI TSI/D = One year following injury	Structured, brief telephone assessment No defined facilitator One instance of assessment at 1-year post-injury	Synchronous No reported use of proxy	Brief Test of Adult Cognition by Telephone (BTACT) Return to Work & Employment Status	BTACT telephone assessment added significantly to predicting return to work following TBI

<p>Schlichter et al., 2020 (USA)⁵²</p>	<p>Descriptive (quality improvement initiative in observational cohort format)</p>	<p>To determine the feasibility of measurements of physical function, cognition, and quality of life in patients requiring neurocritical care.</p>	<p>n = 1324 Adult patients with mixed neurological diagnosis (n = 218 TBI) at home/community</p> <p>Average Age (SD) = 59.5y (17.6)</p> <p>Sex (F) = 45.3%</p> <p>Not reported</p>	<p>Structured telephone assessment utilizing secure web-based data capture platform (REDCap)</p> <p>Clinical nurse, physician, or dedicated research coordinator (all trained in assessment)</p> <p>One instance between 3- and 6-months following discharge</p>	<p>Synchronous</p> <p>Use of proxy (caregiver) for assistance of patient-report or reporting mortality</p>	<p>Modified Telephone Interview for Cognitive Status (mTICS)</p> <p>Patient-reported modified Rankin Scale (mRS)</p> <p>Patient-reported Glasgow Outcome Scale Extended (GOSE)</p> <p>Patient-reported EuroQol 5 Dimension 5 Level Outcome Assessment Tool (EQ5D5L)</p> <p>Overall ‘visual analog scale’ health question, on a scale of 0 to 100</p>	<p>Of all neurological diagnosis cohorts, overall loss to follow-up was 23.6% (n = 313 of 1324) at a mean (SD) time of 4.4 (0.8) months after initial admission.</p> <p>94% of patients or caregivers who answered calls did so by the second attempt.</p> <p>Of remaining TBI patients (n = 123), 24.4% did not answer (n = 30).</p> <p>On average, completed telephone assessments required 21.9 minutes to deliver across all diagnosis cohorts.</p>
<p>Rhame et al., 2020 (USA)⁵³</p>	<p>Descriptive (retrospective analysis of quality improvement initiative)</p>	<p>To describe the implementation and utilization of a Neurotrauma Hotline at a Level I trauma center.</p>	<p>n = 817 unique TBI patients at home/community represented by n = 1205 calls to the service</p>	<p>Neurotrauma telephonic hotline (serviced weekdays, 9am to 5pm) provided to patients upon discharge, with electronic record access for facilitator. Out of hours service provided by voicemail</p>	<p>Synchronous / Asynchronous</p> <p>Proxies able to access service</p>	<p>No reported use of outcome measures administered.</p>	<p>n = 1205 calls over 12 month period (n = 817 unique callers, n = 388 (28.2%) repeat callers) with an average of 3.3 calls/day</p>

			<p>No patient demographics reported.</p> <p>Caller demographics (approximate) = Patients = 29%, Family / Friend = 23%, Outside Provider / Staff = 18%, Internal Provider / Staff = 27%</p> <p>Not reported</p>	<p>messaging or access to on-call neurosurgeon</p> <p>Registered nurse with TBI expertise</p> <p>n = 1205 calls over 12 month period (n = 817 unique callers, n = 388 repeat callers) with an average of 3.3 calls/day</p>			<p>Calls from patients accounted for approximately 58.9% of system usage</p> <p>Calls were answered live 29.5% of the time. Those not answered live were answered at a median time of 3h 18min from initial call.</p> <p>Highest volume of calls received were in December (n = 132) and lowest in February (n = 68).</p> <p>Two highest reasons for calls were appointments (36.8%) or to seek advice (32.1%).</p>
--	--	--	--	---	--	--	---

Abbreviations: y = years. wks = weeks. h = hrs. min = minutes. (F) = female. (SD) = standard deviation. TSI/D = time since injury or diagnosis. GCS = Glasgow coma score. [I] = intervention. [C] = control. LoC = loss of consciousness. w/o = without. ED = emergency department. TBI = traumatic brain injury. CVA = cerebrovascular accident.

Notes: † denotes research published in two parts.

Table 2 - Citations reporting the use of SMS-based Follow-up

Author, Year (Country)	Study Design (Author definition)	Study Aim / Objective	Sample Population Demographics	Follow-up Technology (FUT) Description Clinical Facilitator	Synchronicity Use of Proxy	Constructs & Outcome Measures Deployed	Response / Success / Compliance Rates
------------------------	----------------------------------	-----------------------	--------------------------------	--	-------------------------------	--	---------------------------------------

			TBI Characteristics	Sessions & Instances count				
Smith et al., 2012 (USA) ⁴³	Descriptive (Pilot)	To assess the utility of mobile health (mHealth) technologies, including personal digital assistant-based ecological momentary assessment and two-way interactive text (SMS) messaging, for providing treatment feedback to clinicians, encouraging and motivating veterans throughout treatment, and monitoring participants for relapse after treatment discharge.	<p>n = 27 Adult (military veterans or active members) TBI patients with/without PTSD at home/mental health providers</p> <p>Sex (F) = 0%</p> <p>Mild TBI</p>	<p>In follow-up phase: SMS messaging (ecological momentary assessment) between patient and clinical staff/patient's identified 'buddies', and motivational reminder messages</p> <p>Clinical Staff, Patient 'Buddies' (relative / family member / friend)</p> <p>Approximately 10 'check-in' prompts per month up to 3 months</p>	Asynchronous	No reported use of proxy in response, although 'buddy' or clinical staff notified of contact lapse/below-threshold check-in response for 1-1 follow-up.	Generalized Likert scale 'Check-in' question "How are you doing overall" with (1 = 'Great' to 5 = 'Lousy') and unidirectional motivational messages.	<p>91% participants remained engaged (1 response/30days) at 90 days.</p> <p>Average days enrolled in messaging = 72.2 days</p> <p>Average check-in prompts delivered = 9.2 over 30 days.</p> <p>Average check-in responses = 8.2 over 30 days.</p>
Suffoletto et al., 2013 (USA) ⁶⁶	Experimental (randomised controlled trial)	To examine whether patients with mild traumatic brain injury (mTBI) receiving text	n = 43 (n = 18 intervention, n = 25 control) Adult TBI patients at home/community	<p>Timed, SMS-based symptom assessments with symptom-specific education and reassurance</p> <p>Blinded examiner</p>	Asynchronous	No reported use of proxy	Likert scale questions (0 = none to 4 = severe) across three domains (somatic: headaches, cognitive:	<p>84% (n = 36) completed 14-day SMS follow-up.</p> <p>93% felt that messaging system was useful to help</p>

		messaging-based education and behavioral support had fewer and less severe post-concussive symptoms than those not receiving text message support, and to determine the feasibility of using text messaging to assess daily symptoms and provide support to patients with mTBI.	<p>Average Age (SD) = 30 (9)</p> <p>Sex (F) = 56%</p> <p>Mild TBI</p> <p>TSI/D: Convenience sample from emergency department discharge</p>	3 timed questions (9am, 1pm, 5pm) per day over 14 days		concentration difficulties, emotional: anxiety or irritability), adapted from the Rivermead Post-concussion Symptoms Questionnaire (RPQ)	<p>them self-manage and understand symptoms.</p> <p>Over 14 days, 74% completed 9am headache assessment, 96% completed 1pm difficulty concentrating assessment, and 97% completed 5pm irritability/anxiety assessment.</p> <p>Amongst completed assessments, between 49% to 54% completed <1hr, and 29% to 54% completed <5 mins.</p>
Anthony et al., 2015 (USA) ⁵⁶	Observational (prospective cohort study)	To determine the amount of within-day variation of Concussion Symptom Severity Scores (CSSSs) in athletes with a clinically diagnosed concussion.	<p>n = 14 Youth (sports-related) concussion patients at home/community</p> <p>Age Range = 14 - 22y</p> <p>Clinically-diagnosed symptomatic concussion (CSSS score 10<)</p>	<p>Automated, timed SMS-based symptom checklist ('Text-messaging Robot') with scheduling database</p> <p>No defined facilitator</p> <p>Five scheduled assessments per day for 30 days or until CSSS score of 0 (asymptomatic) for 7 consecutive days</p>	Asynchronous	Concussion Symptom Severity Score (CSSS), calculated via SMS-delivered Post Concussion Symptom Score (PCSS)	<p>804 completed surveys (24180 messages).</p> <p>n = 3 subjects had inadequate response rates.</p> <p>Average follow-up duration 23.9 days.</p> <p>Time of day did not confound responses.</p>

<p>Pacella et al., 2018 (USA)⁵⁹</p>	<p>Observational</p>	<p>To examine changes in post concussive symptoms (PCS) over the acute postinjury recovery period, focusing on how daily PCSs differ between mild traumatic brain injury (mTBI) and other injury types.</p>	<p>n = 108 Adult mixed trauma patients (n = 39 mild TBI, n = 16 head injury w/o TBI, n = 53 non-head-injured trauma control) at home/community</p> <p>n = 39 TBI patients Average Age (SD) = 32 (12.1) Sex (F) = 49%</p> <p>Mild TBI TSI/D = From ED discharge</p>	<p>Automated SMS-based self-reported symptom assessment with response storage on electronic database</p> <p>Research team phone</p> <p>Three timed queries per day (9am, 1pm, 5pm) for 14 days</p>	<p>Asynchronous</p> <p>No reported use of proxy</p>	<p>Experience sampling method, using 3 symptom queries with a 5-point Likert scale to mirror the Rivermead Post concussion Questionnaire (RPQ), assessing: somatic (headaches), cognitive (difficulty concentrating) and emotional (anxiety or irritability) at 9am, 1pm and 5pm respectively</p>	<p>Of the 14 total queries, average of 11.4 completed for headaches, 11.9 for concentration and 11.6 for anxiety.</p> <p>Between 88% and 91% of subjected completed each PCS report on at least one day.</p> <p>Between 35% and 41% of subjects completed these reports every day.</p> <p>Low levels of education were the only variable associated with missing outcome reports (those with less than college education had higher odds of non-completion).</p>
---	----------------------	---	--	---	---	---	--

Abbreviations: y = years. wks = weeks. h = hrs. min = minutes. (F) = female. (SD) = standard deviation. TSI/D = time since injury or diagnosis. GCS = Glasgow coma score. [I] = intervention. [C] = control. LoC = loss of consciousness. w/o = without. ED = emergency department. TBI = traumatic brain injury. CVA = cerebrovascular accident.

Table 3 - Citations reporting the use of Smartphone-based Follow-up

<p>AS ut th hd oy</p>	<p>Study Aim / Objective</p>	<p>Sample Population Demographics TBI Characteristics</p>	<p>Follow-up Technology (FUT) Description Clinical Facilitator</p>	<p>Synchronicity Use of Proxy</p>	<p>Constructs & Outcome Measures Deployed</p>	<p>Response / Success / Compliance Rates</p>
---------------------------------------	------------------------------	---	---	--	---	--

r , Y e a r (C o u n t r y d e f i n i t i o n)			Sessions & Instances count			
J u e n s i - e x t p e r i m e n t	<p>To assess pilot feasibility and validity of a mobile health (mHealth) system for tracking mood-related symptoms after traumatic brain injury (TBI).</p>	<p>n = 20 Adult TBI patients at home/community</p> <p>Average Age (SD) = 36.7 (12.4) Sex (F) = 40%</p> <p>All-severity TBI, classed as initial GCS <12 or 13-15 with positive neuroradiologic findings consistent with TBI</p> <p>TSI/D (SD) = 5.2y (3.6)</p>	<p>Patient-facing smartphone application-based ecological momentary assessment, Personalized EMA Rehabilomics Forms for Rehabilitation Medicine ('iPerform') and clinician-facing web-based portal (iPerform Portal). App has additional communication functions allowing patients to send text messages and clinicians/researchers to send emails to patients.</p>	<p>Asynchronous</p> <p>No reported use of proxy</p>	<p>Daily: Patient Health Questionnaire 2 (PHQ-2)</p> <p>Generalized Anxiety Disorder 2 (GAD-2)</p> <p>General Fatigue statement agreement using 7-point Likert scale (1 =</p>	<p>73.4% assessments completed as scheduled. 79.8% completed as a whole.</p> <p>6.3 / 7 (SD 0.8) patient satisfaction with iPerform smartphone application assessment. From the TUQ, 6.2 / 7 (SD 0.8) reported ease of use, 4.3 / 7 (SD 1.7) for reliability and 5.5 / 7 (SD 1.1) for satisfaction with iPerform.</p> <p>Low reliability thought to be due to technical problems</p>

<p>0 1 5 (U S A) 7 3</p> <p>t a l (p i l o t s t u d y o f p r o s p e c t i v e r e p e a t e d m e a s</p>			<p>For comparison, traditional telephone-based interview</p> <p>Clinician / Researcher team</p> <p>One assessment per day during patient-identified preferred 3-hour window. Two week schedule of varying assessments. Schedule repeated 4 times for 8 weeks total.</p>		<p>strongly disagree, 7 = strongly agree)</p> <p>Positive and Negative Affect Schedule (PANAS)</p> <p>Biweekly: Patient Health Questionnaire 9 (PHQ-9)</p> <p>Generalized Anxiety Disorder 7 (GAD-7)</p> <p>Positive and Negative Affect Schedule (PANAS)</p> <p>Technology evaluation: Compliance (retrospectively calculated)</p> <p>Satisfaction: six 7-point Likert scale questions assessing usability and satisfaction completed bi-weekly via telephone</p>	<p>faced by participants throughout study (notification receipt errors or application crashing and/or not submitting assessment)</p> <p>High correlations with standard telephone-interview supporting validity of smartphone-based mood-related EMA in this population.</p>
---	--	--	--	--	--	--

<p>u r e s d e s i g n)</p>					<p>Usability: Telehealth Usability Questionnaire (TUQ) conducted during final week of assessment</p>	
<p>P a p e r i n s c e n t a k e t a l (s e c o n d i a r (U S A) 6 7 y s</p>	<p>To examine engagement with a mobile application (“mCare”) for wounded Service Members rehabilitating in their communities. Many had behavioral health problems, Traumatic Brain Injury (TBI), and/or post-traumatic stress disorder (PTS), and to examine associations between Service Members’ background characteristics and their engagement with mCare.</p>	<p>n = 95 Adult (military service members) of mixed diagnoses with behavioral health, PTSD and/or TBI at home/community</p> <p>n = 95: Average Age Range (SD) = 34.7 (10.3) to 39.7 (10.4) Sex (F) = 0% to 27.3%</p> <p>Not reported</p>	<p>Bi-directional mobile health smartphone application 'mCare' utilizing SMS updates/prompts, and secure encrypted website to deploy scheduled app-based status questionnaires</p> <p>Care Team Members</p> <p>Daily questionnaires (seven varieties delivered once/week) at 10am local time for up to 36 weeks</p>	<p>Asynchronous</p> <p>No reported use of proxy</p>	<p>Questionnaires included: General Status, Pain Status, Energy and Sleep Status, Anger Management, Relationship Status, Transition Goal Status, Mood Status.</p> <p>Weight status sent once per month.</p>	<p>Participants usually responded to 60% of the questionnaires weekly, generally in 10 h; however, participants with behavioural health problems had several weeks with <50% response/longest response times. Older age and higher general well-being schedule scores were associated with greater and faster responses.</p>

i s o f m u l t i s i t e p r o s p e c t i v e R C T)						
V i e s c r e i t p l i v e .	<p>To determine the feasibility of ecologic momentary assessment following youth concussion, gather real-time reports of cognitive and physical activity, and compare objective measures with</p>	<p>n = 34 pediatric concussion patients at home/community</p> <p>Median Age (range) = 15y (13 - 16)</p> <p>Sex (F) = 47%</p> <p>Concussion, diagnosis based on Zurich consensus diagnostic criteria</p>	<p>Apple iPod Touch with custom application and use of in-built accelerometer for administration of questionnaires following randomly timed prompts by ecological momentary assessment</p> <p>No defined facilitator</p>	<p>Asynchronous</p> <p>No reported use of proxy</p>	<p>Post-Concussion Symptom Scale (PCSS)</p> <p>Activity questionnaire of daily activities</p> <p>Step count</p>	<p>n = 28 (82%) responded to more than 80% of symptom questionnaire prompts.</p> <p>n = 34 were enrolled for a median of 6 days after injury (range 3 - 10)</p>

2016 (USA) 54	real-time reported symptoms among youths during recovery after concussion.	Median TSI/D (range) = 9 days (5 - 13)	Two weeks of daily symptom reports and physical activity monitoring		Daily cognitive rest and exertion (by calculating composite score of measurements in number of texts sent, minutes of screen time and gaming, and minutes of reading or schoolwork)	
VQRtshen-xCPheardndharrtieleatlw.	To evaluate whether the app would be feasible for use by youth with unresolved concussion symptoms as a complement to standard medical care (Phase I); and to assess whether recovery profiles differed between youth who augmented medical care with the app and those who received medical care alone (Phase II).	<p>n = 42 pediatric concussion patients at home/community (Phase I n = 20, Phase II n = 19)</p> <p>Phase I: Average Age (SD, range) = 15.6y (1.6, 13 - 18) Sex (F) = 70%</p> <p>Phase II: Average Age (SD, range) = 15.6 (1.7, 13 - 18) Sex (F) = 77%</p> <p>Physician-diagnosed concussion or Mild TBI (SCAT-3 score 4<) TSI/D = 3wks to 12mo post-injury</p>	<p>Smartphone application 'SuperBetter' encompassing a gamification-based symptoms journal, 'Battle Royal Power Pack', with personal social networking for in-app activity monitoring</p> <p>Research coordinator and friends/family</p> <p>One logged activity per day for 5 days over 3 weeks (target dose of 15 logged activities)</p>	Asynchronous No reported use of proxy	<p>Phase II: Concussion symptom severity on the SCAT-3 checklist score. Secondary: Optimism (measured by Life Orientation Test-Revised [LOTR]), Depression (measured by Center for Epidemiologica l Studies - Depression Child [CES-DC])</p>	<p>In phase I, n = 14 (70%) completed the intervention.</p> <p>Of the 14 participants in Phase, high satisfaction (median = 2, range 0) was reported.</p> <p>Application use was high in both phases (Phase I: median %Play = 110% +/- 22% of target dose, Phase II median 113% +/- 8%).</p> <p>In phase I, barriers to compliance (from remaining n = 6) included discontinuation of medical care (n = 3), unanticipated difficulty with home internet access (n = 1), concomitant illness (n = 1) and competing extracurricular schedules (n = 1).</p>

<p>2 0 1 7 (U S A) 7 2 p h a s , n o n - r a n d o m i z e d , o p e n l a b e l d e s i g n)</p>				<p>Technology evaluation: Phase I: Number of participants completing the intervention relative to all enrolled.</p> <p>Application use (%Play) expressed as % of target dose in first 3 weeks of intervention.</p> <p>Satisfaction with intervention (7-point Likert, 1 = high, 7 = low).</p>	<p>Symptoms and optimism improved more for the experimental than for the active control cohort.</p>
--	--	--	--	--	---

<p style="writing-mode: vertical-rl; transform: rotate(180deg);"> J u e n g s t o n a l (P r o s p e c t i v e s c o h o r t) </p>	<p>To investigate the within- and between-person variability in self-reported emotional symptoms and fatigue, measured through ecological momentary assessment (EMA), among individuals with chronic traumatic brain injury (TBI).</p>	<p>n = 21 Adult TBI patients at home/community</p> <p>Not reported</p> <p>Chronic TBI TSI/D = 6<mo post-injury</p>	<p>Smartphone-based ecological momentary assessment (EMA) on Apple and Android smartphones or tablets</p> <p>No defined facilitator</p> <p>Daily ecological momentary assessment for 8 weeks</p>	<p>Asynchronous</p> <p>No reported use of proxy</p>	<p>Alternating assessments between:</p> <p>Affect, assessed by Positive and Negative Affect Schedule (PANAS)</p> <p>Mood, assessed by Generalized Anxiety Disorder-2 (GAD2) and Patient Health Questionnaire-2 (PHQ2)</p> <p>General Fatigue item using 7-point Likert (“Fatigue interferes with my work, family or social life”)</p>	<p>Of n = 21 consented, n = 17 (81%) completed any daily assessments in the 8 week period.</p> <p>EMA demonstrated statistically significant fluctuations in affect and mood domains, demonstrating applicability of EMA in chronic TBI cohorts to adequately capture temporal symptoms over time.</p>
<p style="writing-mode: vertical-rl; transform: rotate(180deg);"> C r a s i - a n e </p>	<p>To assess the feasibility of using smartphone application technology to assess participation following TBI.</p>	<p>n = 10 Adult TBI patients at home/community</p> <p>Working-age adults</p> <p>Not reported</p>	<p>Smartphone application-based ecological momentary assessment (EMA)</p> <p>No defined facilitator</p>	<p>Asynchronous</p> <p>No reported use of proxy</p>	<p>Mobile Participation Assessment Tool (mPAT)</p> <p>Technology evaluation:</p>	<p>82.9% of all scheduled assessments were completed.</p> <p>Compliance varied by week (80.4%-90.6%) and time of day (79.4%-84.61%).</p>

<p>e x t p a e l r . i , n 2 e 0 n 1 t 7 a (l U (S p A r) o 7 s 7 p e c t i v e , r e p e a t e d m e a s u r e</p>			<p>4 times per day, daily for 4 weeks</p>		<p>Compliance, smartphone application ease of use, comfort using smartphone application to answer questions.</p>	<p>On average, on a scale of 1-5, patients reported mPAT was easy to use (mean = 4.5 SD 0.71), an acceptable way to measure their participation (mean = 4.3 SD 1.06) and satisfied with the mPAT as a measure of their participation (mean = 4.2 SD 1.03).</p> <p>Additionally participants reported feeling comfortable using the application (mean = 4.6 SD 0.52) and is an acceptable way to answer questions (mean = 4.6 SD 0.52)</p>
---	--	--	---	--	--	---

s d e s i g n)						
S u a s i - r i e x p e c t a l · 2 0 1 9 (U S A) 7 8	<p>To evaluate mobile ecological momentary assessment (mEMA) as an approach to measure sport-related concussion (SRC) symptoms, explore the relationships between clinical outcomes and mEMA, and determine whether mEMA was advantageous for predicting recovery outcomes compared to traditional symptom report.</p>	<p>n = 20 pediatric/adolescent concussion patients at home/community</p> <p>Average Age (SD, range) = 15.35 (1.98, 12 - 19)</p> <p>Sex (F) = 40%</p> <p>Diagnosed with isolated sports-related concussion</p> <p>TSI/D = Within 72 hours of injury</p>	<p>Specialized custom smartphone application (iOS and Android) for ecological momentary assessment, 'mEMA', with incorporated prompts (push notification)</p> <p>No defined facilitator</p> <p>Three instances of assessment at predetermined fixed time blocks (morning, afternoon, evening) daily until second follow-up or medical clearance (whichever first)</p>	<p>Asynchronous</p> <p>No reported use of proxy</p>	<p>Neurocognitive testing by Immediate Post-concussion Assessment and Cognitive Testing battery (ImPACT) with Post-concussion Symptom Scale (PCSS) embedded within</p> <p>Vestibular Ocular Motor Screening (VOMS)</p>	<p>90% of participants responded to mEMA prompts with an overall response rate of 52.4% (n = 1155 prompts)</p> <p>Average response rate of 50.4% (SD 29.3) per participant responded to throughout the study, with a range of 5.4% to 95.2%</p> <p>Average prompts received n = 64, range 19 - 173)</p> <p>There was no correlation between number of prompts received and the response rate (Spearman rho = 0.08, P = .77).</p> <p>Participants were less likely to respond as days since injury increase (OR = .91, 95% CI: 0.87-0.94, P< .001)</p> <p>Response rate differed by age with older participants less likely to respond (OR = 0.56, 95% CI: 0.34-0.93, P = .026).</p>

e p e a t e d m e a s u r e s)						<p>There was no association between response rate and time of day - (morning = 50.1%, afternoon = 52.9%, evening = 49.8% P = .411)</p> <p>There was no difference in response rate for initial symptom burden (OR = 0.97, 95% CI: 0.92-1.03, P = .354).</p>
J D u e s n c g r i s t p e t t i a v l e (b s , e 2 c 0 o 1 n 9 d (a U r S y A a) n	<p>To investigate within-person variability in daily self-reported emotional and fatigue symptoms and factors associated with high within-person variability among individuals with chronic traumatic brain injury (TBI).</p>	<p>n = 18 Adult chronic TBI patients at home/community</p> <p>Average Age (SD) = 38.3y (12.7)</p> <p>Sex (F) = 72%</p> <p>All-severity TBI</p> <p>TSI/D = 12y (67%), range 2-27 years post-injury</p>	<p>Smartphone-based ecological momentary assessment (EMA) on Apple and Android smartphones or tablets</p> <p>No defined facilitator</p> <p>Daily instance of assessment for 8 weeks (56 time points)</p>	<p>Asynchronous</p> <p>No reported use of proxy</p>	<p>At odd-numbered time points: Patient Health Questionnaire-2 (PHQ2), Generalized Anxiety Disorders-2 (GAD2), and general 7-point Likert scale question regarding impact of fatigue on daily life.</p> <p>On even-numbered time points: Positive and Negative Affect</p>	<p>Not reported.</p>

<p>5 1</p> <p>a l y s i s o f p r o s p e c t i v e d e s c r i p t i v e p i l o t s t u d</p>					<p>Schedule (PANAS)</p>	
---	--	--	--	--	-----------------------------	--

<p style="writing-mode: vertical-rl; transform: rotate(180deg);"> y) C b s e r v a t i o n a l (a p p l i c a t i o n s) 6 1 v a t i o n a l c o n f e r e n c e s (</p>	<p>To evaluate the feasibility of a smartphone application (app) called MOVES to objectively measure community participation; and compare MOVES with a self-report questionnaire, and differences between veterans with mTBI and civilians without TBI.</p>	<p>n = 16 (n = 11 veterans with TBI, n = 5 civilians with no TBI) at home/community</p> <p>Average Age (SD) = 36.14y (4.9) [veteran cohort], 33.00 (4.9) [civilian cohort]</p> <p>Sex (F) = 0% [both cohorts]</p> <p>Mild TBI TSI/D = Not reported</p>	<p>Smartphone application (MOVES) for iOS and Android, utilizing inbuilt phone GPS tracking. Store-and-forward of data by secure messaging email through a secured portal (MyHealtheVet)</p> <p>Research team</p> <p>Daily activity/GPS tracking for up to 6 weeks</p>	<p>Asynchronous</p> <p>No reported use of proxy`</p>	<p>GPS-based activity and location monitoring (MOVES Storylines)</p> <p>Technology evaluation: Satisfaction questionnaire at week 6 (ten 5-point Likert questions)</p> <p>Daily documentation of unexpected events (technology-related)</p> <p>Perceived accuracy of MOVES Storylines (Perceived Accuracy Daily Logs)</p>	<p>There was a 75% retention rate (n = 11).</p> <p>Participants reported an average of 90% accuracy between the MOVES Storylines and a self-reported questionnaire, Participation Assessment with Recombined Tools Objective (PART-O).</p> <p>Overall, all participants reported they were mostly satisfied (3.65 / 5) with a range of 2.1 to 4.8.</p> <p>The lowest satisfaction rating was received for the effort required with sending the application data to the research team with an average of 2.68 / 5 between veteran and civilian cohorts.</p> <p>The highest satisfaction rating for veterans was an item reporting the effort required to take the smartphone on their persons during the day with an average of 4.29 / 5.</p> <p>The highest satisfaction rating for civilians was an item reporting the ability of the MOVES application to capture their activities in the</p>
--	---	---	---	--	--	---

h o r t s t u d y						community with an average of 4.4 / 5. It took between 10 to 15 minutes to score each daily story line per person, with an estimated 4 to 5 hours required for 28 days of data per person.
---	--	--	--	--	--	--

Abbreviations: y = years. wks = weeks. h = hrs. min = minutes. (F) = female. (SD) = standard deviation. TSI/D = time since injury or diagnosis. GCS = Glasgow coma score. [I] = intervention. [C] = control. LoC = loss of consciousness. w/o = without. ED = emergency department. TBI = traumatic brain injury. CVA = cerebrovascular accident.

Notes: ^{b51} was a secondary analysis of a previous pilot feasibility study, ^{a58}, and included as a separate entry for totality.

Table 4 - Citations reporting the use of Videoconference-based Follow-up

Author, Year (Country)	Study Design (Author definition)	Study Aim / Objective	Sample Population Demographics TBI Characteristics	Follow-up Technology (FUT) Description Clinical Facilitator Sessions & Instances count	Synchronicity Use of Proxy	Constructs & Outcome Measures Deployed	Response / Success / Compliance Rates
Huijgen et al., 2008 (Italy, Spain, Belgium) ⁶⁵	Experimental (Randomized multicenter trial)	To investigate the feasibility of a telerehabilitation intervention for arm/hand function (the Home Care Activity Desk [HCAD] training) in a home setting.	TBI cohort: n = 30 (n = 20 intervention, n = 10 control) Adult TBI patients at home/community Average Age (SD) = 32 (13)[I], 38 (17)[C] Sex (F) = 20% [I], 30% [C]	Portable Home Care Activity Desk (HCAD) consisting of sensorized tools, videoconferencing facilities (2 x webcams) and remote data upload to hospital. Data reviewed in weekly	Synchronous / Asynchronous No reported use of proxy	Action Research Arm Test (ARAT) Nine Hole Peg Test (NHPT) Technology evaluation: General user satisfaction (acceptance, aesthetic, ease of	Average usage across pathologies (TBI, MS, Stroke) = 30 mins per day for 19 days (treatment time 9.5hrs/mo). Average treatment time similar to usual care. Overall compliance varied from 7 - 38

			<p>TSI/D (SD) = 7.5y (4.4) [I], 7.8y (2.9) [C]</p>	<p>patient-therapist videoconferencing</p> <p>Therapist</p> <p>30 minute sessions per day for 5 days per week (20 days total)</p>		<p>use, task difficulty, task appropriateness, general impression of HCAD)</p>	<p>days (recommended 30mins / 5 days per week totaling 20 days)</p> <p>As a majority, both patients and therapists satisfied with HCAD. Only aesthetic aspects of system and task difficulty resulted in slightly less satisfaction as a whole.</p>
<p>Hill et al.^c, 2009 (Australia)⁷⁰</p>	<p>Experimental (randomised controlled trial)</p>	<p>To determine if valid and reliable assessment of apraxia of speech using a standardized assessment tool was feasible via an internet-based telerehabilitation system.</p>	<p>n = 11 Adult patients with mixed diagnosis (n = 2 TBI, n = 9 CVA) at hospital research laboratory 15km from assessor</p> <p>TBI cohort, n = 2: Average Age = 20y Sex (F) = 50%</p> <p>Not reported TSI/D (TBI cohort, n = 2) = 6.5mo average post-injury</p>	<p>Custom real-time videoconference-based assessment using two web cameras mounted on robotic arm over 128kbit/s connection. Participant wore headset microphone and earphones. System incorporated concurrent automatic store-and-forward facilities integrated into software for high-resolution video and audio data sharing</p> <p>Speech-language pathologist</p> <p>Single instance of assessment</p>	<p>Synchronous / asynchronous</p> <p>No reported use of proxy</p>	<p>Apraxia Battery for Adults 2 (ABA-2)</p> <p>Technology evaluation: Participant Satisfaction Questionnaire, consisting of eight-items employing 5-point Likert scale</p>	<p>Across diagnosis cohorts, no significant differences were found between the telerehabilitation assessment versus in-person assessments, with moderate to very good agreement indicated.</p> <p>All participants eligible to complete the satisfaction questionnaire (n = 5) described the audio quality as good or excellent. n = 3 described the video quality as good or excellent, n = 2 adequate.</p>

							<p>n = 3 described the comfort level during the videoconference sessions as comfortable or very happy, with n = 1 having no feeling either way and n = 1 described it as uneasy.</p> <p>With regards to overall satisfaction, all were satisfied, with n = 3 describing the videoconference assessment as more than or very satisfied.</p> <p>All but one participant indicated they would be equally satisfied with services being delivered via videoconference versus in-person.</p>
<p>Hill et al.^d, 2009 (Australia)⁷¹</p>	<p>Experimental (randomised controlled trial)</p>	<p>To refine the telerehabilitation system used in the Hill et al.⁷⁰ study and re-evaluate this new system with a modified research design to</p>	<p>n = 24 Adult patients with mixed diagnosis (n = 11 TBI) at hospital research laboratory 15km from assessor</p> <p>n = 24:</p>	<p>Custom real-time videoconference-based assessment using two web cameras mounted on robotic arm over 128kbit/s connection. Participant wore headset microphone</p>	<p>Synchronous / Asynchronous</p> <p>No reported use of proxy</p>	<p>Assessment battery including:</p> <p>Informal oromotor assessment</p> <p>Informal perpetual speech assessment</p>	<p>Good strength of agreement was found between the FTF and telerehabilitation assessment methods.</p>

		determine validity and reliability of the assessment of acquired dysarthria in adults.	<p>Average Age (range) = 50.2y (16 - 78) Sex (F) = 37.5%</p> <p>Not reported TSI/D (range) (n = 24) = 42.6mo average (6mo - 11y)</p>	<p>and earphones. System incorporated concurrent automatic store-and-forward facilities integrated into software for high-resolution video and audio data sharing. Additional data-sharing capabilities that allowed instructional images and videos to be displayed locally versus transmitted allowing more streamlined and efficient assessment.</p> <p>Speech-language pathologist</p> <p>Single instance of assessment</p>		<p>Assessment of Intelligibility of Dysarthric Speech (ASSIDS)</p> <p>Technology evaluation: Participant Satisfaction Questionnaire, consisting of eight-items employing 5-point Likert scale</p>	<p>The majority of participants (n = 10 of n = 11) rated both the audio and visual quality as good or excellent.</p> <p>All (n = 11) participants were comfortable or very happy with the telerehabilitation assessment session, and all participants rated their overall satisfaction as more than satisfied or very satisfied.</p> <p>All (n = 11) participants reported being confident with the results gained via telerehabilitation assessment, and all were willing to participate in future telerehabilitation assessments.</p> <p>The majority of the participants (n = 8) stated that they would be equally satisfied with speech pathology services delivered</p>
--	--	--	---	--	--	---	--

							<p>via telerehabilitation methods.</p> <p>Only n = 4 thought it would be more convenient for them to access speech pathology services in this manner. Of the n = 7, n = 5 felt that telerehabilitation would not be more convenient for them, and n = 2 stated that it was not applicable as they did not have access to the Internet at home.</p>
<p>Rietdijk et al., 2017 (Australia)⁷⁵</p>	<p>Quasi-experimental (repeated measures design with randomized order)</p>	<p>To compare in-person with videoconferencing administration of a communication questionnaire for people with traumatic brain injury (TBI) and their close others.</p>	<p>n = 20 Adult TBI patients and their close others at home/community</p> <p>Sex (F) = 20% ‘Close others’ = n = 8 parents, n = 6 partners, n = 3 other family members, n = 3 friend of TBI patient</p> <p>Severe TBI (post-traumatic amnesia duration 21-180 days).</p>	<p>Videoconference - based (Skype) assessment with telephone fallback, with in-person comparison conducted in patient’s home</p> <p>Clinician researcher</p> <p>One instance of remote assessment paired with one instance in-person between 1 and 2 weeks apart</p>	<p>Synchronous</p> <p>Use of proxy, ‘close others’</p>	<p>La Trobe Communication Questionnaire (LCQ) for patient self-report (LCQ Form S) and their close other (LCQ Form O)</p>	<p>n = 1 excluded due to lack of comprehension of questionnaire in either in-person or videoconference setting.</p> <p>Of remainder, 89.5% (n = 17) received successful administration of videoconference outcome measure.</p> <p>Due to connection and quality difficulties, n = 2</p>

			TSI/D (range) = 6< months (8mo - 22y)				were administered by telephone. No significant differences between videoconferencing and in-person for LCQ score or administration time.
--	--	--	---------------------------------------	--	--	--	---

Abbreviations: y = years. wks = weeks. h = hrs. min = minutes. (F) = female. (SD) = standard deviation. TSI/D = time since injury or diagnosis. GCS = Glasgow coma score. [I] = intervention. [C] = control. LoC = loss of consciousness. w/o = without. ED = emergency department. TBI = traumatic brain injury. CVA = cerebrovascular accident.

Notes: Of studies ^{70c} and ^{71d} conducted by the same authors, ^d was included separately due to revised methods and a novel patient cohort.

Table 5 - Citations reporting the use of Miscellaneous Technology-based Follow-up

Author, Year (Country)	Study Design (Author definition)	Study Aim / Objective	Sample Population Demographics TBI Characteristics	Follow-up Technology (FUT) Description Clinical Facilitator Sessions & Instances count	Synchronicity Use of Proxy	Constructs & Outcome Measures Deployed	Response / Success / Compliance Rates
Lewandowski et al., 2009 (USA) ⁶³	Observational	To examine the feasibility of a momentary data-gathering method, as well as the sensitivity of the assessment to the subtle and dynamic changes in symptoms of concussion	n = 3 pediatric concussion patients and n = 3 healthy pediatric individuals at home/community Total: Age Range = 14 - 17y Sex (F) [concussion patients, n = 3] = 66.6%	Stylus-based touchscreen personal digital assistant (Palm Pilot 100) based ecological momentary assessment with auditory prompts No defined facilitator Five daily instances of delivery (between 9-10am, 11am-12pm, 2-	Asynchronous No reported use of proxy	Setting context Symptom Severity Scale (SSS), consisting of 13 symptom items and 7-point Likert scale Functional Status Scale (FSS), consisting of 15 functional impairment items	Each assessment took approximately 3-5min. There were no technical difficulties reported. Of a possible 75 assessments, 70 were complete, demonstrating a compliance rate of 93.3%.

			Symptomatic concussion/mTBI TSI/D = Average 117 days (range 78 - 165)	3pm, 5-6pm, 8-9pm) for five consecutive weekdays. Prompts received every 5 minutes until response received.		and 7-point Likert scale	The device was reported to be not disruptive to others in a school setting nor to the students using it.
Lenaert et al., 2019 (Netherlands) ⁶²	Observational (longitudinal observational study)	To investigate the feasibility of using experience sampling method (ESM) in individuals with acquired brain injury (ABI), to explore the usability of ESM data on a clinical level, by illustrating the interactions between person, environment, and affect	n = 17 Adult ABI patients (n = 8 TBI) at home/community n = 17: Average Age (SD, range) = 44.2y (14.5, 18 - 65) Sex (F) = 53% Not reported	Touchscreen electronic device, ‘PsyMate’, with semi-randomly scheduled auditory prompts for ecological momentary assessment No defined facilitator 10 instances delivered daily at semi-random scheduled times over 6 days	Asynchronous No reported use of proxy	Positive & Negative Affect Schedule (PANAS) Location and Social Context including appraisal using bipolar scale Activities and Physical well-being (including fatigue), including appraisal using bipolar scale Technology evaluation: Subjective experiences assessed by two debriefing questionnaires (18, 7-point Likert items in total) on user friendliness and general acceptability of methodology	Average response rate of 71.18% (n = 726) following delivery of 1020 prompts. 98.76% of prompted reports were completed, with an average of 42.7 questionnaires answered (range 28 - 57). There were no dropouts reported, with the method experienced as user-friendly. The device was reported to have little influence on their activities or social contacts (Average = 2.00 / 7, SD 1.16). There were little to no difficulties reported when using the device (Average

							<p>= 1.77 / 7, SD 1.36), with the device not experienced as burdensome (Average = 2.08 / 7, SD 1.19).</p> <p>The amount of beeps were not seen as much of a burden (Average = 2.08 / 7, SD 1.19)</p>
--	--	--	--	--	--	--	--

Abbreviations: y = years. wks = weeks. h = hrs. min = minutes. (F) = female. (SD) = standard deviation. TSI/D = time since injury or diagnosis. GCS = Glasgow coma score. [I] = intervention. [C] = control. LoC = loss of consciousness. w/o = without. ED = emergency department. TBI = traumatic brain injury. CVA = cerebrovascular accident.

Table 6 - Outcome measures deployed via follow-up technologies in TBI populations

Outcome measure (or derivative)	Abbreviated Outcome measure	Number of implementations	Citations
Glasgow Outcome Scale Extended <i>Glasgow Outcome Scale Extended: Pediatrics (GOSE-Peds)</i>	GOS-E GOSE-Peds	5 2	10,48,49,52,79 19,49
Rivermead Post-concussion Questionnaire <i>Short-form derivatives assessing key domain-symptom pairs (somatic: headaches, cognitive: concentration difficulty, emotional: anxiety or irritability)</i>	RPQ -	3 2	46,47,50 59,66
5-Level EuroQol 5-Dimension	EQ-5D-5L	4	46,48,50,52
Generalized Anxiety Disorder Assessment <i>Generalized Anxiety Disorder Assessment 2-item</i>	GAD-7 GAD-2	1 3	73 51,58,73
Public Health Questionnaire <i>Public Health Questionnaire 2-item</i> <i>Public Health Questionnaire 9-item</i>	PHQ PHQ-2 PHQ-9	0 3 1	- 51,58,73 73
Positive and Negative Affect Schedule	PANAS	4	51,58,62,73
Post-Concussion Symptom Score <i>Concussion Symptom Severity Score</i> <i>Immediate Post-concussion Assessment and Cognitive Testing Battery</i>	PCSS CSSS ImPACT	1 1 1	54 56 78
Brief Test of Adult Cognition by Telephone	BTACT	2	57,60
Glasgow Outcome Scale <i>Glasgow Outcome Scale: Hindi variation</i>	GOS GOS: Hindi	0 1	- 44
Medical Outcomes Study Short Form <i>12-Item Short Form Survey</i>	SF-36 SF-12	0 2	- 46,50
Telephone Interview for Cognitive Status <i>Modified Telephone Interview for Cognitive Status</i>	TICS TICSm	1 1	41 52
Action Research Arm Test	ARAT	1	65
Acute Concussion Evaluation	ACE	1	68
Apraxia Battery for Adults-2	ABA-2	1	70
Assessment of Intelligibility of Dysarthric Speech	ASSIDS	1	71
BSF/A: Functional Independence Measure	FIM	1	41
Center for Epidemiological Studies Depression Scale for Children	CES-DC	1	72
Community Participation Indicators <i>Computer Adaptive Testing Community Participation Indicators</i>	CPI CAT-CPI	0 1	- 55
Functional Status Scale	FSS	1	63
La Trobe Communication Questionnaire, self-reported (Form S) and proxy-reported (Form O)	LCQ	1	75
Mobile Participation Assessment Tool	mPAT	1	77
Modified Rankin Scale	mRS	1	52
Neurobehavioral Rating Scale	NRS	1	41
Nine Hole Peg Test	NHPT	1	65
Ohio State University TBI Identification Method	OSU TBI-ID	1	74
Post-Concussion Symptom Inventory: Parent Assessment Form	PCSI	1	68
Revised Life Orientation Test	LOT-R	1	72
Sports Concussion Assessment Tool 3	SCAT-3	1	72
Symptom Severity Scale (SSS)	SSS	1	63
Vestibular Oculomotor Screening	VOMS	1	78

