

Fractal dimension links responses to a visual scene to its biodiversity

Paul Stevens

Abstract

Humans appear to have an innate, beneficial response and preference for natural over urban scenes, yet "natural" is an ambiguous concept that varies from culture to culture. In looking for a commonality to natural scenes that tends to be lacking in built scenes, many researchers have turned to fractal geometry, finding that fractal dimension can predict preference. Here, I calculated the fractal dimension of the the dominant land-sky edge at a variety of sites having varying depths of water-table and levels of biodiversity (specifically, "species richness"). I then investigated changes in human physiological arousal (magnitude of skin conductance responses) in response to images of those scenes. Sites with high biodiversity were shown to have a significantly higher associated fractal dimension than low biodiversity sites, whereas shallow versus deep water-table sites showed no significant difference. When shown the images, the magnitude of skin conductance responses for human viewers showed a negative correlation with fractal dimension. Replicating earlier findings, ranked preference for a scene showed a positive correlation with fractal dimension. Taken together, these findings suggest an evolved response to stimuli associated with a healthy ecosystem: patterns of healthy vegetative growth determine visual fractal dimension, which reduces physiological arousal upon observation, this being experienced as a positive emotional state and expressed as a preference for that environment.

Keywords: fractal, natural environment, biodiversity, restorative.

1. Introduction

Research concerned with wider issues of how humans and the natural environment interact has demonstrated some interesting properties of human visual perception. It has long been known that people, irrespective of culture or education, tend to express a preference for natural scenes


over urban or built ones (Ulrich et al., 1991). It was later discovered that such preference appears to relate to some fundamental psychophysiological responses – arousal levels decrease, attentional capacity increases and emotional processing occurs faster (Ulrich, 1983; Parsons & Hartig, 2000) – suggesting that we might be preferentially responding to specific features related to the natural environments within which human sensory systems evolved (Balling & Falk, 1982). However, the concept of “natural” is still relatively ambiguous, covering a huge range of scenes from lush jungle vegetation to open savanna to barren, rocky mountains. To understand the empirical findings for human preference, there needs to be some commonality to natural scenes that tends to be lacking from artificial scenes. Many researchers have turned to fractal geometry for inspiration, with some promising results.

1.1 Fractal geometry

Since Mandelbrot (1983) published his mathematical description of the complex forms found in nature, research has shown that a wide range of natural forms exhibit repeating patterns when viewed at increasingly high magnifications i.e., they have a fractal geometry. This self-similarity of pattern at differing scales can be quantified by a parameter called the fractal dimension (D), essentially a non-integral quantity that relates to the number of self-similar pieces that an object can be “broken into” at different scales (e.g., see Glass & Mackey, 1988. p.53). So a simple line can be broken into as many self-similar pieces as you want at any magnification e.g., you can break it in half (a magnification of 2) to get 2 pieces, into 3 pieces at a magnification of 3, or N^1 pieces at a magnification of N . A square, however, can be broken into 4 self-similar pieces where each piece has sides that are half as long as the original, or 9 self-similar pieces with sides $1/3$ as long as the original i.e., you get 4 pieces at a magnification of 2, 9 pieces at a magnification of 3, or N^2 pieces at a magnification of N . This pattern thus gives us a simple power-law based definition of fractal dimension, $D = \log(\text{number of self-similar pieces}) / \log(\text{magnification factor})$. In application to two-dimensional images (which usually do not have obvious self-similarity), a commonly used technique for estimating D for that images is the box-counting technique

(Abarbanel, 1996). Essentially, this makes use of a computer algorithm which determines the number of “boxes” of linear size L needed to cover all the black points in a black and white image, for diminishing values of L . Plotting $N(L)$ against L will then have a line of best fit that has a gradient of $-D$.

Hagerhall et al. (2004) used this technique to explore whether fractal dimension may play a role in human preference for visual landscapes, based on previous cross-cultural findings (e.g., Ulrich, 1983) that natural scenes (i.e., the ones more likely to have a fractal geometry) tend to be preferred to urban ones. For their analysis, they focussed on the fractal dimension of the silhouette outline between the landscape and the sky, this representing the most striking feature upon which people fixate according to eye movement studies (e.g., Rayner & Pollatsek, 1992). To obtain this outline, the background area (i.e., the sky) for each image was selected using the automatic “area select” tool used by many graphical editing software. A “find edges” tool was then used to isolate the edges of the selected area, resulting in a single line tracing the contour between foreground objects and the background (see fig. 1). This line was then used to calculate D via the box-



counting method, thus giving a measure of visual complexity for each image.

Figure 1. Example of an extracted land-sky silhouette for fractal dimension calculation

For a sample of 52 images of natural landscapes with no visible water or dominant hills (the visual presence of either is a factor that appears to skew scene preference studies e.g., see Purcell & Lamb, 1984), Hagerhall et al (2004) found a significant positive correlation between the mean preference for an image and the fractal dimension of the extracted silhouette outline. A further study by Cheung and Wells (2004) also replicated the effect using just 12 images – six urban and six natural scenes – finding a significant correlation between mean preference rating

and both mean and maximum fractal dimension. Both of these studies confirmed an earlier exploratory study by Rogowitz and Voss (1990), which suggested that preference was based on the D value of the edge of dominant shapes in a scene. This ties in well with other research showing that, based on both empirical work and models of cortical neurons involved in subtle shape discrimination, human vision is optimised to power-law relationships of the optical environment (Párraga et al., 2000).

So why should this be? The key lies in some recent studies which found that the fractal dimension of landscape features also correlates to an interesting, non-visual property of those landscapes: they are “healthier” in terms of biodiversity (a measure of the variety of life, plant and animal, within a given area). For example, Krummel et al. (1987) showed that the fractal dimension was significantly higher for outlines of untouched versus human-affected forest, probably relating to differences in the scale of human versus natural processes. Olf and Ritchie (2002) point out that “habitat, food and resources for organisms often are found to be statistically self-similar across ecologically relevant ranges of scales” and showed that fractal dimension could distinguish between habitat loss and habitat fragmentation (the former having a lower fractal dimension and more negative effect on biodiversity than the latter). Brown et al. (2002) theorise that this is because “scaling relationships that are self-similar or fractal-like over a wide range of spatial or temporal scales” (p.619) represent a class of emergent ecological phenomena. These relationships demonstrate underlying power laws that appear to be universal with respect to the type of organism or type of environment, offering “clues to underlying mechanisms that powerfully constrain biodiversity” (p.619) and could help us to understand “the diversity of species and complexity of ecosystems in terms of fundamental principles of physical and biological science” (p.619).

The present study thus had three hypotheses:

1. the fractal dimension, D , of the dominant edge in the land-sky silhouette would be higher in images of areas having high biodiversity.
2. the D of the dominant edge in the land-sky silhouette would be higher in images of

areas having a shallow-depth ($\leq 25\text{m}$) of water table.

3. the magnitude of skin conductance response to viewing an image would be inversely correlated to the D of that image.

Hypothesis 1 aimed to establish a primary relationship that one visual aspect of healthy vegetation (as indicated by D of the land-sky silhouette) would relate to biodiversity in the area. Hypothesis 2 would explore whether this biodiversity–D relationship might be due to the amount of accessible water available to vegetation in the area. Hypothesis 3 explored whether any change in human physiological arousal, as measured by skin conductance, in response to viewing an image was also related to the fractal dimension of that image's land-sky silhouette. Hypotheses 1 & 2 were explored in experiment 1, and hypothesis 3 in experiment 2 (which used a selection of images from experiment 1).

2. Experiment 1: Fractal dimension of water-depth and biodiversity images

2.1 Methods

All images used in the experiment were photographed to be representative of the wider location, ensuring there were clear tree-lines away from roads or other sources of potential pollutants. Any obviously human-intended plantings (e.g., around entrances, visitor centres or plantation areas) were avoided. None of the images in either category contained man-made structures (to avoid the possible negative connotations of an urban environment), visible bodies of water or dominant hills to avoid known preference biases (cf. Hagerhall et al., 2004). Purely urban images were also excluded as there would be no direct link between biodiversity measures and the landscape: the fractal dimension of artificial structures would be due to the aesthetic architectural choices made rather than there being any causal link. For example, an urban scene with gothic architecture would have higher fractal dimension than one with modernist buildings (e.g., Joye, 2007), yet this would be unrelated to biodiversity. All photos were digitally photographed, from a similar distance at the same resolution (10MP), and within a few weeks of each other under similar weather conditions. Such precautions were taken to ensure a robust comparison, even though

previous research has indicated that D values are neither sensitive to changes in scale nor dependent on the precise technique by which the land-sky silhouette is extracted (Keller, Crownover & Chen 1987).

2.1.1 Images of shallow/deep water-table sites

Based on the *British Geological Survey* hydrogeological map of Dorset (where the author was based at the time), 10 sites were identified in areas showing a range of depth of underground aquifers. Based on Jackson et al.'s (1999) empirical determination of a maximum tree root depth of 25m, five specific locations were determined to have “shallow” depth water table ($\leq 25\text{m}$ i.e., water table is accessible to tree roots) and five to have “deep” depth water table ($\geq 50\text{m}$ i.e., water table is inaccessible to tree roots). Two images were then taken to be representative of each site, giving twenty images in all: 10 shallow, 10 deep (see Table 1).

2.1.2 Images of high/low biodiversity sites

Species richness (the number of different species identified in a specified area) was chosen as the specific indicator of biodiversity, as there is general consensus that this is a reasonable, simple measure of an area's biodiversity (Gotelli & Colwell, 2001; Natural History Museum, 2009). These were obtained from the website of the UK's *National Biodiversity Network (NBN)* (<http://data.nbn.org.uk/>), which provides publicly available species lists recorded within or overlapping a specific site at 10km resolution. Five sites were chosen based on their inclusion in the *NBN* site and physically accessible (i.e., the sites could be visited to take photographs), and to cover a range of biodiversity (from 3646 to 21683 species). These sites were different from the ones used for the water-table images. A total of 40 images were obtained across this range – see table 2 for a list. For the purpose of the preplanned analysis, sites with species richness at or below the median value were categorised as low biodiversity; sites above the median value were categorised as high biodiversity.

Table 1. List of water-table category sites with depth (m) to water table

Site	Depth to water table (m)	Water-table category
Nine Stones, Winterbourne Abbas	0	Shallow
Balmers Coombe Bottom, Bulbarrow	0	
Cerne Park, Cerne Abbas	20	
Delcombe wood, Bulbarrow	20	
Horse Clump, Winterbourne Abbas	23	
Big Wood, Winterbourne Steepleton	50	Deep
Jubilee Trail, Dry Wood, Winterbourne Steepleton	55	
Twitchings Copse, Bulbarrow	65	
Woolland hill, Bulbarrow	70	
Park Dale, Cerne Abbas	110	

Table 2. List of biodiversity category sites showing species richness for each area

Site (number of photos)	Species Richness	Biodiversity Category
Canford Heath (5)	3,646	Low
Sopley Common (8)	4,159	
Avon Heath (7)	4,733	
Studland Nature Reserve (8)	8,340	High
New Forest National Park (12)	21,683	

2.1.3 Fractal dimension analysis

For each image, the technique of Hagerhall et al (2004) described earlier was used to isolate the silhouette of the dominant edge (landscape-sky silhouette). This was done using the open source software *ImageJ* (available from <http://rsbweb.nih.gov/ij/>), which was also used to calculate the fractal dimension of that edge. For each image, the JPG was converted to binary and the 'find edges' tool used on the sky region. The 'fractal box count' tool with then used with box sizes set to (2, 3, 4, 6, 8, 12, 16, 32, 64) to calculate the fractal dimension.

2.2. Results

An Initial review of the water table category images indicated that there were two possible

outliers. This was confirmed by calculating the two standard deviation range (mean fractal dimension, $D=1.293$. $SD=0.043$, thus the $2SD$ range is from 1.207 to 1.379) and removing two datapoints outwith this range (Cerne Park site: $D=1.207$ and 1.204). For the reduced data set, a Wilcoxon test comparing the fractal dimension for shallow versus deep sites gave a close to significance result: $W=23$, $N=18$, $P=0.07$ (see fig. 2). For the biodiversity category images, a Wilcoxon test comparing fractal dimension for high versus low biodiversity sites gave a significant result: $W=25$, $N=40$, $P<0.001$ (see fig. 2). A post-hoc analysis of image D-value versus ranking by species diversity gave a Spearman rho=0.64.




Figure 2. Comparison of fractal dimension of image silhouette for deep versus shallow water-table and low versus high biodiversity. Error bars indicate one standard deviation.

3. Experiment 2: Physiological reactions to and preference for the images

3.1. Methods

3.1.1. Participants

A total of 50 (34 females, 16 males) unselected participants – primarily university students – $M=19$ years, ranging from 18–31 years) participated in this study. The study was advertised on campus and via email to a potential participant pool maintained by the psychology department.

Students volunteered in return for course credit and all participants were paid a small expenses fee. All gave informed consent before participating and the study had ethical approval from the psychology research ethics committee.

3.1.2. Stimuli

Five photographic images were selected from each the two categories *water-table* and *biodiversity* from experiment 1 to cover the a representative range of fractal dimension found in the whole image set ($D=1.22$ to 1.40). The biodiversity set is shown in figure 3 as an example.



Figure 3: Images from the biodiversity set, showing increasing fractal dimension (D) from top to bottom.

3.1.3. Skin conductance recording

Skin conductance (SC) was recorded for each participant via two electrodes attached to the second phalanx of the index and second fingers of their non-dominant hand. Electrodes were sintered Ag-AgCl round cup electrodes with an 8mm diameter, affixed with adhesive collars and using pH balanced aqueous gel. These were connected via preamplifier to a model SC5-SA recorder with 24-bit A/D conversion (PsyLab/Contact Precision Instruments, London, UK) and interfaced via serial port to a PC running the *Windows* operating system with custom-written software in *Visual Basic 6.0*.

Data analysis was conducted offline on a PC running Linux (Ubuntu 11.04). The raw SC data consisted of two components: the longer-term tonic response and the fast, stimulus-related phasic responses. The tonic component from each participant's data was first removed by filtering

the data with a high pass filter of 0.016 Hz (equivalent to a time constant of 10 seconds), using the GPL program *QtiPlot* (<http://soft.proindependent.com/qtiplot.html>). Subsequent analysis was performed using the GPL program *R* (<http://www.r-project.org/>). A z-transform (i.e., mean-centred and expressed in units of each participant's standard deviation) was used on the SC phasic data to allow between participant comparison and to give a more robust measure for subsequent statistical analysis (Sersen et al., 1978). The 200 sample (five second) period after each stimulus was then isolated to give that participant's series of phasic SC responses (SCRs), one per stimulus image.

3.1.4 Procedure

On arrival, participants were told they were taking part in a study into scene preference that combined conscious (stated preference) and unconscious (skin conductance measures). They were seated in front of a 17" touch-sensitive TFT computer monitor and the procedure explained to them. The SC electrodes were attached and, after listening to a five minute piece of relaxing music (a standard procedure to ensure all participants start from close to their baseline arousal state), they viewed each set of five images in turn. Set and image order were randomized and counterbalanced for each participant, with each image being shown on the monitor for 10 s with a 10 s rest period between each one. SC data were sequentially sampled and saved to hard disk at 40 Hz throughout the session. After they had seen all five images in a set, they were asked, via touch-screen response, to rank those images in order of personal preference, "from MOST to LEAST favourite". The second set was then shown, and again, preferences recorded. Finally, they were debriefed and paid £3 expenses for their participation.

3.2. Results

SCRs were screened to remove null-responses, movement and noise artefacts, resulting in 280 usable response profiles across the participants, for all 10 images. As the z-transformed SCR magnitude data approximated a normal distribution, a Pearson correlation was used ($r_P = -0.11$,

$N=280$, $P=0.03$), showing a small but significant negative relationship between normalised SCR magnitude and fractal dimension of the associated image (see fig. 4). Image preference ratings were also correlated with D using a Spearman rank correlation ($r_s=0.37$, $N=480$, $P<0.001$), showing that preference was positively related to fractal dimension (see fig. 5).




Figure 4. Plot of mean normalized SCR magnitude versus fractal dimension, D




Figure 5. Plot of mean expressed preference versus fractal dimension, D

4. Discussion

This study explored the idea that fractal dimension might represent a stimulus which maps onto the basic human need to live in a healthy environment, and to better understand relationships between the fractal dimension of visual scenes, human physiological responses to such scenes,

and self-reported preference for specific visual environments.

In Experiment 1, fractal dimension varied significantly between high and low biodiversity sites ($P < 0.001$), but not in relation to shallow versus deep water table sites ($P = 0.07$). This suggested that the types and growth patterns of vegetation primarily reflect the ecological health of the wider area. For this experiment, the measure of biodiversity used was “species richness”: the number of species – plant, animal and fungus – observed in a 10km^2 region. The results thus suggested that a single measure of the fractal dimension of the skyline of an image taken within this region could be used as an indication of the ecological health of the region as a whole. This in itself could, if more widely replicated, provide a useful estimate of an area's biodiversity that did not rely on costly surveys.

Experiment 2 showed significant negative relationship between the magnitude of a person's physiological response and the fractal dimension of the dominant edge in a visual image, suggesting that the level of physiological arousal we experience in response to seeing that image (or scene, if we were physically present) relates to the visual complexity of that image or scene. However, as can be seen in figure 4, there are no obvious visual cues which would lead the perceiver to determine that complexity. This relationship was in the direction predicted by existing research in the field of restorative environments (e.g., Ulrich et al., 1991) which typically show reduced arousal for more “natural” images (which would be expected to have higher fractal dimension) when compared to more “urban” images (which would be expected to show more linear geometry and so have lower fractal dimension). As well as directly affecting wellbeing via a more relaxed somatic state, lowered arousal, in the absence of any specific external affective cues, tends to be interpreted as a positive emotional state (e.g., Stevens, 2007) which in itself is associated with enhanced wellbeing (e.g. Lyubomirsky et al., 2005). That the lowered arousal did result in positive affective associations was supported in this experiment by the finding that preference for an image was also positively related to fractal dimension (and thus showed a negative relationship with arousal), replicating earlier research (Hagerhall et al., 2004; Cheung & Wells, 2004; Rogowitz & Voss, 1990).

Combining the results suggests the presence of an evolved response in humans to stimuli associated with a healthy ecosystem. The ecological health of an area is expressed via patterns of vegetative growth that is exhibited as increased fractal dimension. When humans – and, given the similarity of cortical structure, other mammals – observe that scene, reduced physiological arousal occurs in response to visual complexity and this is experienced as a positive emotional state and expressed as a preference. Simply put, it shows that the environments which we *innately* prefer are those which can help restore our wellbeing if they themselves are part of a healthy, functioning ecosystems.

More generally, these findings suggest a way to encourage people to realise that they are not separate from the pressing environmental issues of the day, allowing us to see “the needs of the planet and the person as a continuum” (Roszak, 1992, p. 14) by showing that the properties which allow humans to be physically and mentally healthy are the same ones which are associated with a healthy ecosystem on local and global scales. Human emotional responses to specific environments do not necessarily need to be validated purely by ideological arguments but instead can be seen as involving innate responses to places and situations that represent a meaningful, evolved means of communication between us and our environment. This could allow a reintegration of direct (emotional, reflective) experience in pro-environmental strategies rather than putting all the emphasis on “objective” arguments and perception-of-risk calculations – an idea also seen in some other psychology studies which have shown that early childhood experience of natural settings and outdoor recreation that relies on specific natural features (e.g., white water rafting) are the strongest predictors of subsequent pro-environmental behaviour (e.g., Kals et al., 1999). Rather than changing people's motivation by bombarding them with information and so-called rational arguments, evolutionary arguments also have a place by showing there are inherent motivations, intentions and behaviours that can play a role.

However, the results of this and similar studies could be read as encouraging reductionism i.e., we do not actually need natural environments to be healthy, we just need to ensure that our built environment has a high fractal dimension in terms of visual cues. Such an approach

doubtlessly has a place in “customising visual landscapes and wall art to aid human functioning and stress reduction in mentally demanding environments” and “incorporating favourable visual properties in the design of our everyday environments to foster general well-being” (Hagerhall et al., 2008, p. 1492), yet to do this as our only or primary response to problematic environments would be to impoverish human experience. The intention of this study was to shed light on the pre-existing literature spanning many disciplines which shows that humans are embedded in, and so profoundly affected by, their environment (Stevens, 2010), the “natural” environments being associated with many more benefits/less stressors than “urban” ones. This is unsurprising, given that “natural” features are the ones we, along with all other animals, are evolutionarily adapted to, but it is hoped that the results of this study demonstrate that fractal dimension offers one way of better understanding what we mean by “natural” that goes beyond the usual socially constructed and human-centric approaches.

Acknowledgments

This research was supported by Fundação Bial, Portugal (grant no. 71/08). Thanks are also due to Helen Kirkup for her able assistance in data collection.

References

- Abarbanel, H. D. I (1996). *Analysis of observed chaotic data*. New York: Springer Verlag.
- Balling, J. D., & Falk, J. H. (1982). Development of Visual Preference for Natural Environments. *Environment and Behavior*, 14, 5-28.
- Brown, J. H., Gupta, V. K., Li, B.-L., Milne, B. T., Restrepo, C., & West, G. B. (2002). The fractal nature of nature: power laws, ecological complexity and biodiversity. *Philosophical Transactions of the Royal Society of London B*, 357, 619-626.
- Cheung, K. C., & Wells, N. M. (2004). The natural environment & human well-being: Insights from fractal composition analysis? HarFA - Harmonic and Fractal Image Analysis 2004, pp. 76-82.
- http://www.fch.vutbr.cz/lectures/imagesci/includes/harfa_e_journal.inc.php. Accessed 2 Feb

2011

Glass, L., & Mackey, M. C. (1988). *From clocks to chaos: the rhythms of life*. Princeton, NJ: Princeton University Press.

Gotelli, N. J., & Colwell, R. K. (2001). Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters*, 4, 379-391.

Hagerhall, C. M., Purcell, T., & Taylor, R. (2004). Fractal dimension of landscape silhouette outlines as a predictor of landscape preference. *Journal of Environmental Psychology*, 24, 247-255.

Hagerhall, C. M., Laike, T., Taylor, R. P., Küller, M., Küller, R., & Martin, T.P. (2008). Investigations of human EEG response to viewing fractal patterns. *Perception*, 37, 1488-1494.

Jackson, R. B., Moore, L. A., Hoffmann, W. A., Pockman, W. T., & Linder, C. R. (1999). Ecosystem rooting depth determined with caves and DNA. *PNAS*, 96, 11387-11392.

Joye, Y. (2007). Architectural lessons from environmental psychology: The case of biophilic architecture. *Review of General Psychology*, 11, 305–328.

Kals, E., Schumacher, D., & Montada, L (1999). Emotional affinity toward Nature as a motivational basis to protect Nature. *Environment and Behavior*, 31, 178-202.

Karten, H. J. (1997). Evolutionary developmental biology meets the brain: the origins of mammalian cortex. *PNAS*, 94, 2800-2804.

Keller, J. M., Crownover, R. M., & Chen, R. Y. (1987). Characteristics of natural scenes related to the fractal dimension. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 9, 621–627.

Krummel, J. R., Gardner, R. H., Sugihara, G., O'Neill, R. V., & Coleman, P. R. (1987). Landscape patterns in a disturbed environment. *Oikos*, 48, 321-324.

Lyubomirsky, S., King, L. & Diener, E. (2005). The benefits of frequent positive affect: does happiness lead to success? *Psychological Bulletin*, 131, 803-855.

Mandelbrot, B. B. (1983). *The fractal geometry of nature*. New York: W. H. Freeman and Co.

Natural History Museum (2009). Popular estimates of biodiversity value: using species richness.

<http://www.nhm.ac.uk/research-curation/research/projects/worldmap/diversity/species.htm>.

Accessed 2nd December 2009.

- Olf, H., & Ritchie, M. E. (2002). Fragmented nature: consequences for biodiversity. *Landscape and Urban Planning*, *58*, 83–92.x
- Párraga, C. A. Troscianko, T., & Tolhurst, D. J. (2000). The human visual system is optimised for processing the spatial information in natural visual images. *Current Biology*, *10*, 35-38.
- Parsons, R., & Hartig, T. (2000). Environmental psychophysiology. In J. T. Cacioppo, L. G. Tassinary, & G. G. Berntson (Eds.), *Handbook of psychophysiology*. Cambridge: Cambridge University Press.
- Purcell, A. T., Lamb, R. J., Mainardi Peron, E., & Falchero, S. (1994). Preference or preferences for landscape. *Journal of Environmental Psychology*, *14*, 195–205
- Rayner, K., & Pollatsek, A. (1992). Eye movements and scene perception. *Canadian Journal of Psychology*, *46*, 342–376.
- Rogowitz, B. E., & Voss, R. F. (1990). Shape perception and low dimension fractal boundary contours. *Proceedings of the Conference on Human Vision: Methods, Models and Applications, SPIE/SPSE Symposium on Electron Imaging*, *1249*, 387–394.
- Roszak, T. (1992). *The voice of the Earth*. New York: Simon and Schuster.
- Sersen, E. A., Clausen, J., & Lidsky, A. (1978). Autonomic specificity and stereotypy revisited. *Psychophysiology*, *5*, 60-67.
- Spehar, B., Clifford, C.W.G., Newell, B.R., & Taylor, R.P. (2003). Universal aesthetic of fractals. *Computers & Graphics*, *27*, 813–820.
- Stevens, P. (2007). Affective response to 5 μ T ELF magnetic field-induced physiological changes. *Bioelectromagnetics*, *28*, 109-114.
- Stevens, P. (2010). Embedment in the environment – a new paradigm for wellbeing? *Perspectives in Public Health*, *130*, 265-269.
- Ulrich, R. S., Simons, R. F., Losito B. D., Fiorito, E., Mile, M. A., & Zelson, M. (1991). Stress recovery during exposure to natural and urban environments. *Journal of Environmental Psychology*, *11*, 201-230.
- Ulrich, R. S. (1983). Aesthetic and affective response to natural environment. In I. Altman, & J. F. Wohlwill (Eds.), *Behavior and the natural environment: human behavior and environment –*

advances in theory and research volume 6 (pp. 85-126). New York: Plenum Press.