

Assessing the Roles of Sender and Experimenter in Dream ESP Research

Chris A. Roe*, Simon J. Sherwood*, Louise Farrell*,
Louie Savva*, and Ian S. Baker**

*Centre for the Study of Anomalous Psychological Processes,
University of Northampton, United Kingdom

**Koestler Parapsychology Unit, University of Edinburgh,
United Kingdom

Abstract

This study explored the role of the sender in a dream ESP task by considering the effects of presence of a sender (sender, no sender) and the receiver's expectancy that a sender was present. Forty participants each completed a sender and a no sender trial on consecutive nights by keeping a dream diary of all mentation they could recall when they awoke. The order of trials was randomised across participants. On no-sender nights a randomly selected video clip was played repeatedly from 2:00 until 6:30am; on sender nights a sender would also watch the clip between 6:00 and 6:30am and attempt to communicate its content. Both sender and no sender conditions produced above chance hit rates (30% and 35% respectively), but z scores for similarity ratings did not deviate significantly from chance (sender night: $t_{(39)} = 0.92, p = .18$; no sender night: $t_{(39)} = 1.11, p = .14$, one-tailed). There was no difference in performance in terms of sender conditions ($z = -0.22, p = .41$, one-tailed) or sender expectancy ($z = -0.18, p = .46$, one-tailed), failing to support the proposal that senders play an active role in dream ESP success. Possible improvements in the manipulation of participant expectancy are discussed.

Introduction

J. B. Rhine's advice to those who hoped to study *psi* in the laboratory involved an analogy with making rabbit stew; "If you want to have rabbit stew, first catch the rabbit" (Stanford, 1993, p. 129).¹ Thus if we are to study the action of *psi* in the laboratory we need to ensure that all aspects of the laboratory situation are arranged so as to facilitate (or at least not inhibit) its occurrence. *Psi* phenomena are not renowned for their experimental reliability (e.g., Beloff, 1983, Milton and Wiseman, 1999, see Shapin and Coly, 1985 for an extended discussion), and all too often we have been left to feed on scraps. It might be naïve to expect replication on demand given the effect sizes typically involved (cf. Utts, 1991), but nevertheless there must be a suspicion that *psi* is sensitive to some factors that have not been adequately explored or typically are not controlled for effectively. If different laboratories differ in these subtle respects it could lead superficially similar experiments to generate different outcomes, as some enjoy rabbit stew while others settle for vegetable broth. Efforts to identify potential confounding factors promise to inform us of the necessary conditions to capture *psi* more consistently as well as perhaps offering some insight into its *modus operandi*.

In looking to map these necessary conditions we have recently been especially concerned to consider the sender-receiver-experimenter dynamic as a factor (or collection of factors) that moderates *psi* performance in Ganzfeld ESP trials (Roe, Sherwood and Holt, 2004; Sherwood, Roe, Holt and Wilson, 2005). In the first of these studies (Roe et al., 2004) we attempted to distinguish between the active contribution a sender might make and the positive effects of simply believing that a friend was viewing the target. The direct hit rate was exactly at chance (25%) and, although this was slightly better for sender trials than no sender trials (26.1% versus 23.5%) and better for trials on which the receiver believed there was a sender than when they believed there was not, irrespective of whether there actually was one (33.3% versus 18.2%), there were no significant differences between conditions based on *z* scores of target ratings (for sender status $p = .632$; for sender expectancy $p = .765$). This was disappointing given that of seven previ-

¹After presenting a version of this paper at the International Conference of the Society for Psychical Research in September 2006, Sean O'Donnell questioned Rhine's choice of analogy here, which seems to suggest that *psi* is something to be slain and dismembered by its investigators. We have some sympathy with his suggestion that a better analogy would be of *psi* as a butterfly, delicately to be caught and closely observed in its natural, intact and healthy state.

ous Ganzfeld studies that directly compared sender and no-sender conditions within the same study, all reported better performance with a sender, significantly so for two studies (see Roe et al., 2004 for a more thorough review). We therefore intended to reconsider this aspect in the current study. In a subsequent study we explored whether participants' perceptions of the experimenter and of the experimenter's attempts to generate a warm social ambiance were predictive of performance (Sherwood et al., 2005). Here both participants and the experimenter completed an interaction questionnaire that asked about their mood, expectations of success and sense of rapport with the other participants. Responses on the interaction questionnaire suggested that participants were typically in a good mood, fairly relaxed, optimistic about the trial — though not confident of success — and had a positive perception of the experimenter, all of which were expected to be psi conducive features. Nevertheless, the direct hit rate for this study was nonsignificantly worse than chance (21.1%, $z = -.015$). However, although relationships between these variables and trial outcomes (in terms of z scores for similarity ratings) did not exhibit a clear pattern and tended to vary somewhat from experimenter to experimenter, they did offer some significant overall relationships, such as with receiver mood (negative-positive; $r_s = -.335$), sender optimism ($r_s = .432$) and confidence of success ($r_s = .398$) which we felt warranted further work.

We planned to follow up these findings by conducting a further study that would investigate the effects of both sender and experimenter upon the receiver's ability to identify a target video clip based on correspondences with their own mentation. However, using the Ganzfeld procedure as a method of eliciting psi has proved to be very labour intensive, and may have deterred some participants from volunteering because of the time commitment required and the coordination necessary when involving a sender and receiver. In looking for an alternative method we were encouraged to reconsider dream ESP as a paradigm, since the original Maimonides research programme was notably successful (Child, 1985; Ullman and Krippner, with Vaughan, 1973), and there is evidence of above-chance scoring among subsequent replication attempts with experimental designs using a simplified method that did not require REM monitoring or access to sleep laboratories (e.g., Dalton, Steinkamp and Sherwood, 1999; see Sherwood and Roe, 2003, for a review).

In considering previous dream ESP research we can derive some

75 encouragement for suggesting that sender and experimenter effects
might be evident here too. For example, Ullman et al. (1973), in reflect-
ing on the performance of participants in the Maimonides dream ESP
series, commented that “[T]he active involvement of the agent [sender]
is an important ingredient for success.” (p. 212). The majority of Mai-
80 monides studies investigated telepathy rather than clairvoyance, which
might be interpreted as a tacit assumption that a sender can facilitate
psi hitting in a dream ESP task. However, this presumption was not
supported in Sherwood and Roe’s (2003) summary analysis, which sug-
gested that overall the clairvoyance studies in this series had been more
85 successful than those intended to study telepathy. Of course, making
comparisons across studies may be misleading, since they could have
differed in other ways besides whether or not they involved a sender,
and we expected that a direct comparison within a single study should
clarify this relationship.

90 There is also some suggestion in the database of post-Maimonides
dream ESP studies that some groups of researchers have been markedly
more successful than others (see Sherwood and Roe, 2003, pp. 102–104).
Given the diversity of approaches used in these studies it is difficult to
attribute these differences to one particular cause, but it may be worth-
95 while to consider whether differences in experimenter-participant inter-
action have some effect.

Planned analyses

The main planned analyses are to consider the following hypothe-
ses:

- 100 1. Participant performance, in terms of z score of target rating, will be
higher than chance expectation for each experimental night
2. Participant performance in the actual sender condition will be
higher than that in the no sender condition
3. Participant performance in trials for which the participant believed
105 there was a sender will be higher than for trials for which the par-
ticipant believed there was not a sender

We are grateful to an anonymous referee for suggesting that we
consider differences in success between the two senders and we include
this *post-hoc* analysis here. Following Sherwood et al. (2005), we also

110 planned to conduct exploratory analyses considering covariation of performance (using z -scores of target ratings) with experimenter interaction measures.

Design

115 This study employed a repeated measures design to assess the role of the sender in dream ESP research, with each participant completing a sender and a no sender trial night. Participants remained blind as to which night was which but were asked to nominate on which night they felt there was a sender so as to allow us to consider expectancy effects. Potential psychological experimenter effects were assessed by correlat-
120 ing participant and experimenter ratings of their interaction against task performance. In all cases the dependent variable was pre-specified to be the z score of target clip ratings.

Method

Participants

125 Forty participants were recruited from a variety of sources including the undergraduate population, appeals to the media and an established research database. The sample consisted of 30 females and 10 males² (Mean age = 32.08; range = 19–62; mean score on belief measure = 80.32, range = 19–119, absolute range of scale = 19–133, mid-
130 point = 76), of whom 21 had practised a mental discipline (e.g., meditation/relaxation techniques) at some point and 15 had practised a physical discipline (e.g., Yoga/martial arts). Participants were not selected on the basis of their gender or age; neither were they screened for prior experience or for ability to recall dreams. For trials 1–26, Louie Savva
135 (LS) acted as experimenter; for trials 27–40 Louise Farrell (LF) acted as experimenter.

Apparatus

This study used an automated program for selecting and playing video clips that was developed by Dr Paul Stevens and written in Mi-
140 crosoft Visual Basic.³ Video clips are stored digitally as MPEG files, la-

²One anonymous reviewer speculated that the gender bias toward female participants could have been problematic given that there may be gender differences at this task. *Post hoc* analysis revealed that males performed slightly better than females overall (mean $z = .40$ and $.06$ respectively) but this difference was not significant overall (Mann-Whitney $z = -.93$, $p = .36$) or for either night separately (Mann-Whitney $z = -1.66$, $p = .10$; $z = -.41$, $p = .70$, two-tailed).

³For a more complete description of the program, see Roe, Sherwood, Luke and Farrell (2002).

belled 1a, 1b, 1c etc. The target set consists of 116 minute-long digital video clips arranged in 29 sets of 4. These were the same clips used in our previous Ganzfeld studies (Roe, et al., 2004; Sherwood et al., 2005) and have mainly been produced at the University of Northampton and
145 are drawn from popular television programmes and commercial films, although some have been taken from the pool previously used at Edinburgh. Copies of the target pool are available on DVD from the first author upon request. Randomisation is achieved using the Visual Basic pseudo-random algorithm (rnd), having seeded it using the timer at the start of the program (RANDOMIZE TIMER). Once the "Start" button
150 has been pressed, the computer first selects a target set, then selects one of the 4 clips within that set. The order of presentation of the four clips at judging is similarly randomised. The target sets were selected with replacement. All trials were run at the sender's home (CR or SS) using
155 standard desktop PCs and could be set to play through the night.

Materials

The *Participant Information Form* (PIF) is a 56-item measure that was constructed for general use with parapsychological research at the University of Northampton and was based partially on a version used previously at the University of Edinburgh. It includes questions concerning biographical and contact details (11-items); religious and parapsychological background (5 items); computer experience (2 items); practice of mental/physical disciplines (2 items); belief in luck (2 items); clumsiness and punctuality (2 items); competitiveness (1 item); absorption (2 items); sleep and dreams (4 items); imagination and fantasy-proneness (3 items); creativity (2 items); and physical and mental health (1 item). The remaining items relate specifically to knowledge, belief and experience of anomalous phenomena including telepathy, clairvoyance, precognition, psychokinesis, 'communication with the dead' and
170 out of body experiences (19 items). The form concludes with questions about hypnagogic/hypnopompic experiences in a range of modalities (10 items) and an open question inviting descriptions of personal anomalous sleep-related experiences. Copies of all in-house measures are available from the first author on request.

175 Participants also completed the short extraversion and neuroticism subscales of the EPQ-R⁴ (Eysenck, Eysenck, and Barrett, 1985). Each subscale has 24 items with a dichotomous yes/no response format. A

⁴An analysis of these personality variables will not form part of this paper

belief in paranormal measure, adapted from Thalbourne and Delin's (1993) 18-item *Australian Sheep-Goat Scale* (ASGS) with an additional item asking about their performance in the current study and incorporating a 7-point Likert response scale ranging from strongly agree to strongly disagree, was also completed.

Given the apparent importance of experimenter-participant interactions and expectations of success, at the end of the pre-judging briefing, the experimenters and participants completed a short *Interaction Questionnaire*.⁵ This contained six questions, concerning their personal feelings and expectations and perceptions of the quality of the interactions between experimenter and participants, which they were required to answer by giving ratings on 7-point scales (see Appendix).

Procedure

Once recruited for the experiment, participants were sent — either through the post or by email — the PIF, and they returned this ahead of their trial or brought it in with them on the day of judging. Participants were also sent a paper-copy of a dream diary, which contained instructions to clarify the procedure, information about confidentiality and the experimenter's contact information, as well as space to record their dreams and their associations or comments on dreams that they could recall during the trial period. They kept the dream diary for three consecutive nights; night 1 was a practice night and nights 2 and 3 were experimental nights.

On each night participants slept at their own homes, as normal. Upon waking, either during the night or in the morning, they completed the appropriate section of their dream diary, writing down as much detail concerning their dreams as possible. There was space in the diary for participants to note any associations they may have identified between their dreams and events in their waking lives about which they may have been preoccupied or worried. Participants were also asked to indicate on which of the trial nights they believed there had been a sender and on which there had been no sender. It was intended that the participants make this judgement prior to their meeting with the experimenter but in practice this was not always the case.

A sender (either SS or CR) was allocated to each participant on the basis of availability. Both senders remained blind as to participant identities until after the study was completed. On the evening of Night 2 (the

⁵Experimenters did not complete question 3.

215 first experimental night) the sender flipped a coin to determine whether
 Night 2 would be designated a telepathy or a clairvoyance trial. If Night
 2 was a telepathy trial then by default Night 3 was a clairvoyance trial,
 and vice versa. The experimenter remained blind as to which condition
 had been selected for each trial night until after the participants' judg-
 220 ments had been made and recorded. On both experimental nights the
 sender initiated the experimental software at his own home, where a
 randomly-selected target clip was played from 2:00am until 6:30am. On
 the telepathy trial only, the target clip was played for the same length of
 time but the sender watched the target clip between 6:00am and 6:30am⁶
 225 and attempted to communicate its content to the participant by ESP. The
 sender had no contact with the experimenter concerning the experiment
 until the information was needed about the target sets used for the trial
 nights.

After keeping the dream diary for three nights the participant trav-
 230 elled to the university campus to take part in the judging.⁷ Participants
 were requested to attend judging as soon as possible after completing
 the dream diary — ideally the morning after the last dream night — so
 that dream content might still be retrievable from memory; however,
 appointments were arranged at their convenience. On arrival they were
 235 met by the experimenter, who took them to a reception room where
 refreshments were available. They were then engaged in an informal
 conversation incorporating a brief discussion of their experiences. Af-
 ter this interaction the participant was asked to complete a measure
 that asked them to rate the experimenter along a number of dimen-
 240 sions, such as warmth, spontaneity, and optimism (see Sherwood et
 al., 2004). The experimenter also completed brief ratings of confidence
 about the trial and his/her assessment of the interaction with the partic-
 ipant. These forms were placed in a sealed envelope and sent to an in-
 dependent researcher (IB). Participants were assured that we were only
 245 interested in their honest impressions and that at no time would the
 experimenter be aware of the ratings they had given.

Next, the participant read out their dream diary content; if they

⁶The sending period was set at 30 minutes after initial pilot work suggested that it was difficult for senders to maintain their interest and focus on the clip over a longer time span. In this respect it is comparable with a typical sending period in a Ganzfeld session. Although there is no guarantee that a sending period of 30 minutes will necessarily overlap with a receiver REM period, Braud (1977) has previously reported successful dream ESP performance when sending between 6:00 and 6:30am.

⁷On some trials, where access to the reception rooms was not possible, judging was conducted in the experimenter's own office or the participant's own home.

had not remembered any dream material they still participated in the judging phase and were advised to use ‘intuition’ or to see if the clips prompted any dream recall. The judging phase commenced with the experimenter accessing an SMS text message from the sender that identified the target sets for nights 2 and 3 (but not the identities of the target clips). The sender was shown, via a laptop computer, the four clips that made up the target set (consisting of the target clip and three decoys in random order) for night 2, and these were rated and rank ordered for their degree of correspondence to the dream mentation for that night. This process was repeated for the target set and dream mentation for night 3. Once all judgements were recorded the experimenter contacted the sender via mobile phone to discover the identities of the two target clips and which night had been the sender night.

Once the series was completed, IB was provided with trial outcome data (but no personal information from participants) and analysed the relationships between these and the interaction data sent to him. The other members of the team only saw the results of analyses conducted by IB and were not at any stage provided with the raw data from the interaction measures.

Results

To assess our prediction that participants would award a similarity rating to the target that was higher than the average rating for the three dummy clips for each experimental night, z scores were calculated (see Table 1). The overall mean z score for both nights was slightly positive (actual sender night mean $z = 0.13$, $SD = 0.86$; actual no sender night mean $z = 0.16$, $SD = 0.91$), indicating that the target clips tended to be awarded higher dream correspondence ratings than the other clips, but this did not differ significantly from zero (for no sender night: Wilcoxon $z = -1.06$, $p = .15$; for sender night: Wilcoxon $z = .91$, $p = .18$, one tailed), and the hypothesis was therefore not supported. We did not plan to evaluate performance in terms of direct hits, but for information we note that the proportion of direct hits is nonsignificantly higher than mean chance expectation of 25% for both sender and no sender trials ($z = 0.55$ and 1.28 respectively).

Contrary to expectations, the mean z -score for ratings was greater for no sender trials compared with sender trials (mean $z = 0.16$ versus 0.13), but a Wilcoxon Signed Ranks test found that the difference was

Table 1: Target rank frequencies and z -score based upon ratings for actual sender and no sender conditions ($N = 40$ in each case), with similarity rating (SR) mean z -scores and standard deviations (SD)

	Rank				SR	
	1	2	3	4	mean z	$SD z$
Sender trials	12 (30.0%)	11 (27.5%)	10 (25.0%)	7 (17.5%)	0.13	0.86
No sender trials	14 (35.0%)	7 (17.5%)	13 (32.5%)	6 (15.0%)	0.16	0.91

285 not significant ($z = -0.22$, $p = .41$, one-tailed), suggesting that having a sender did not help participants' performance. Neither condition deviated significantly from the null value of $z = 0$.

As well as looking for gross differences between sender and no sender conditions we were able here to consider sender differences, since this study involved two senders who were quasi-randomly allocated to participants and who contributed 20 trials each. Exploratory analysis of the results by the different senders are given in Table 2, and reveal an interesting but unexpected interaction effect: SS's participants performed as predicted, with greater success on sender nights compared with no sender nights, although this difference is not significant (Wilcoxon $z = 1.23$, $p = .22$, two-tailed); however, the reverse was true for CR's participants, with better performance on no sender nights compared with sender nights, to a degree that was suggestive (Wilcoxon $z = -1.66$, $p = .10$, two-tailed). Thus, whether or not having a sender try to send the target is advantageous may depend on who the sender is. This full interaction trend, which could explain the lack of an overall sender effect, is further illustrated in Figure 1.

 Table 2: Target rank frequencies and z -score based upon ratings for actual sender and no sender conditions ($N = 20$ in all cases), with similarity rating (SR) mean z -scores and standard deviations (SD)

	Rank				SR	
	1	2	3	4	mean z	$SD z$
CR						
Sender	5 (25.0%)	5 (25.0%)	5 (25.0%)	5 (25.0%)	-0.04	0.91
No sender	9 (45.0%)	4 (20.0%)	5 (25.0%)	2 (10.0%)	0.32	0.91
SS						
Sender	7 (35.0%)	6 (30.0%)	5 (25.0%)	2 (10.0%)	0.30	0.79
No sender	5 (25.0%)	3 (15.0%)	8 (40.0%)	4 (20.0%)	0.002	0.83

We also planned to consider the effects upon performance of partic-




Figure 1. Mean z -scores based upon ratings for actual sender and no sender conditions according to sender

ipants' belief that a sender had been operating, and so participants were
 305 asked to nominate on which night they believed there was a sender
 and on which night there was not (of course, they were blind to ac-
 tual sender status). Participants correctly identified the order of sender
 and no-sender conditions for their trial on just 18 of the 40 trials, which
 is slightly less than chance expectation of 20 and suggests that partici-
 310 pants could not accurately identify the sender condition. This allows us
 to consider the effects of 'expectancy' separately from actual condition,
 since the two variables are unrelated. A summary of participant perfor-
 mance on nights when they believed there was a sender operating com-
 pared with nights when they believed there was no sender are given
 315 in Table 3. Contrary to expectations, the mean z -score for ratings was
 slightly greater for the trials when the participants did not believe there
 had been a sender compared to the trials when they believed there had
 been a sender, although a Wilcoxon Signed Ranks test found that this
 difference was not statistically significant ($z = -0.18, p = .46$, one-tailed).
 320 Thus, judgements regarding sender status did not affect performance in
 the dream ESP task.

We also conducted a combined analysis that looked for possible
 interactions between actual sender status and participants' judgements
 that there had been a sender. These data are summarised in Table 4, and

Table 3: Target rank frequencies and z -score based upon ratings for conditions deemed to be sender and no sender ($N = 20$ in all cases), with similarity rating (SR) mean z -scores and standard deviations (SD)

	Rank				SR	
	1	2	3	4	mean z	$SD z$
Deemed Sender	12 (30.0%)	9 (22.5%)	13 (32.5%)	6 (15.0%)	0.13	0.92
Deemed No sender	14 (35.0%)	9 (22.5%)	10 (25.0%)	7 (17.5%)	0.16	0.86

325 show that, in terms of our planned outcome measure of z -scores for sim-
 ilarity ratings⁸, the worst performance was for the condition that was
 expected to be optimal, where participants believed there had been a
 sender and there in fact was, although the differences across conditions
 are not significant (Wilcoxon $z = -.24$, $p = .81$, two-tailed).⁹ Where par-
 330 ticipants wrongly identified sender and no sender conditions there is no
 difference whatever in performance between the two trials (Wilcoxon z
 $= 0.00$, $p = 1.00$, two-tailed). Taken together, these results give little indi-
 cation of any overall sender or sender expectancy effect in this study but
 there is some indication that this may be influenced by who the sender
 335 was; indeed, based on the planned outcome measure of z scores, these
 results do not give evidence of ESP at all (despite the percentage hit
 rates being reasonably consistent with others' Ganzfeld findings).

Finally we conducted exploratory analyses to see whether perfor-
 mance covaried with measures of the quality of interaction between
 340 experimenter and participant. Spearman rank order correlations be-
 tween the different questionnaire ratings and the experimental results
 are summarised in Table 5, and include separate analyses for each ex-
 perimenter. Generally, there is little indication from these that perfor-
 mance at the dream ESP task can be predicted by scores on interaction
 345 measures, with only three of 90 correlations achieving significance. Al-
 though all of these were associated with experimenter or participant
 mood, the effects seem to be reversed for sender and no sender trials,
 making interpretation difficult; since the analyses have not been cor-

⁸In terms of direct hits, the best performance occurred with no sender trials in which the partici-
 pant believed there had been no sender and worst performance is for sender trials where participants
 believed there had been a sender. In terms of mean ranks, performance is again worst with the sender-
 sender condition that was expected to be optimal; performance in the other three conditions is very
 similar.

⁹An omnibus test of these data would not be appropriate given their non-independence (all partici-
 pants contribute to two of the 2×2 cells)

Table 4: Target rank frequencies and z -score based upon ratings for actual sender and no sender conditions separated by participants' judgments of sender status, and similarity rating (SR) mean z -scores and standard deviations (SD)

	Rank				SR	
	1	2	3	4	mean z	$SD z$
Deemed Sender						
Sender present ^a	5 (28%)	4 (22%)	6 (33%)	3 (17%)	.053	.927
No sender present ^b	7 (32%)	5 (23%)	7 (32%)	3 (14%)	.187	.922
Deemed No Sender						
Sender present ^b	7 (32%)	7 (32%)	4 (18%)	4 (18%)	.184	.821
No sender present ^a	7 (39%)	2 (11%)	6 (33%)	3 (17%)	.128	.924

^a $N = 18$

^b $N = 22$

rected for multiple analyses they may simply reflect random noise and
 350 would need to be replicated in future work before they should be in-
 terpreted as a real effect. Some other associations generated relatively
 large coefficients, particularly with confidence, although this is in the
 opposite direction to prediction. There do seem to be some differences
 between experimenters, which perhaps reflect their different personali-
 355 ties and interaction styles.

Discussion

Although hit rates of 30% and 35% over two sets of 40 dream ESP
 trial nights is somewhat above the mean chance expectation of 25%, and
 represents an improvement on the overall hit rate of 21.1% from our
 360 previous study that used Ganzfeld stimulation, deviations from chance
 expectation based on z scores of similarity ratings (our pre-specified
 dependent variable) were not sufficient to give statistical significance
 (respectively, mean $z = 0.13$, $t_{(39)} = 0.92$, $p = .18$, one-tailed; mean z
 = 0.16, $t_{(39)} = 1.11$, $p = .14$, one-tailed), and so we are unable to reject
 365 the null hypothesis. Of course, we may not expect to see evidence of
 psi in any summary measure since this study included conditions that
 were hypothesised to show differential psi performance, although we
 did not confirm our prediction that participants would perform better
 on sender trials than on no-sender trials. This failure to capture any
 370 sender effect is consistent with Sherwood et al.'s (2005) failure to find
 a sender effect in earlier Ganzfeld work and with Sherwood and Roe's
 (2003) finding that post-Maimonides dream ESP studies that had inves-
 tigated clairvoyance had in fact been more successful than those ostensi-

Table 5: Spearman correlation co-efficients for z -score experiment results for the sender and no-sender conditions correlated against the sender and experimenter questionnaire ratings. The co-efficients are shown for the overall results, and also broken down by experimenter

Questions	Overall ($N = 40$)		<i>Experimenter</i>				
	Sender	No Sender	LS ($N = 26$)		LF ($N = 14$)		
			Sender	No Sender	Sender	No Sender	
<i>Participant</i>	Mood	-.03	.08	-.13	-.14	.08	.64*
	Feeling	-.14	-.14	-.27	-.11	-.46	-.14
	Optimism	.23	.18	.20	.10	.22	.28
	Confidence	-.09	-.09	-.04	-.07	-.52	-.15
	Rapport	-.07	-.03	-.03	.07	.00	-.13
	Warmth	-.05	-.04	.00	.08	-.18	-.16
	Spontaneity	.08 ^a	-.13 ^a	.02 ^b	-.14 ^b	.05 ^c	-.01 ^c
	Positiveness	.11 ^a	.02 ^a	.15 ^b	.11 ^b	-.04 ^c	.02 ^c
<i>Experimenter</i>	Mood	-.34*	.22	-.28	.04	-.12	.62*
	Feeling	-.07	-.05	-.08	.10	.14	-.11
	Rapport	-.11	.20	-.21	.27	-.08	.09
	Confidence	-.11	.05	.13	-.10	-.50	.26
	Warmth	.04	.16	-.09	.36	.16	-.23
	Spontaneity	.04	.15	-.06	.31	.31	-.09
	Positiveness	-.10	.23	-.10	.34	-.04	.11

* $p < .03$ (2-tailed).

^a $N = 38$ due to missing values.

^b $N = 25$ due to missing values.

^c $N = 13$ due to missing values.

bly investigating telepathy — despite earlier researchers advocating the
 375 involvement of an agent (e.g., Ullman et al., 1973). One might speculate,
 then, that it may be desirable to concentrate on clairvoyance designs in
 future research, given the advantages this would offer in recruitment
 (since participants could be scheduled one at a time instead of having
 to co-ordinate across two people) and in security (with sensory leakage
 380 and cheating made less likely when no-one needs to be aware of the
 target until after judging is completed), and apparently minimal disad-
 vantages in terms of impact upon effect sizes.

One form in which senders might play a role without being essen-
 tial to any psi process is if they make the task seem inherently more
 385 plausible or help diffuse responsibility for any psi that occurs (see Roe
 et al., 2004, for a fuller discussion), and indeed Roe et al. did report a
 sender expectancy effect, whereby participants performed better when
 they believed that a sender was involved compared with when they
 believed there was no sender (with hit rates of 33.3% and 18.2% respec-

tively), irrespective of actual sender status. In this study there was no indication of any sender expectancy effect, with better performance actually occurring where participants believed there was no sender. This may not be a legitimate comparison, however, since in Roe et al.'s (2004) study participant expectancy was manipulated by giving either true or false information at the beginning of the session, whereas in this study participants decided retrospectively for themselves, typically by reflecting on their dream mentation, whether a trial involved a sender. In this latter case then, participants might have no particular expectancy as to whether or not the first trial will involve a sender, since at that point they have no material on which to base their judgement, and it may be that any expectancy effect would only be expressed on the second trial night (although this too may be problematic, since some participants in the current study only made their judgements after completing both nights).¹⁰

In any case, any potential advantages of involving a sender may have been undermined here by our decision on security grounds to keep the senders and participants isolated from one another to the extent that senders and participants were given no information whatsoever about one another. Several of the participants recruited and run by LF expressed doubt that telepathy would be possible under circumstances in which the people involved had never established any rapport, indeed had never met each other or even knew each other's names. Likewise the senders found it a handicap to have no shared experiences or knowledge about the participant on which to base their sending strategy. We should note, however, that although Honorton et al. (1990) reported a suggestive advantage for friends as senders, Broughton and Alexander (1997) found that participants with a lab-assigned sender achieved a much higher hit rate than those who had a spouse or friend serve in that role. Nevertheless, on reflection, this security measure seems Draconian, and in future studies greater effort would be needed to ensure that some degree of rapport is possible, for example by adopting aspects of the methodology used in remote healing studies (e.g. providing a photograph or the first name of the other party, as in Sicher, Targ, Moore and Smith, 1998).

Analysis of participant-experimenter interaction data did not re-

¹⁰We are grateful to our referees for suggesting that it would be useful in future studies to have participants rate the confidence they have in their estimates and also to ask participants how they decided whether trials had been sender or no sender.

veal any strong trends, and given the scope for committing Type I errors, no confident claims are made from these results. If taken at face value, the best predictors seem to be participant and experimenter mood, although these only emerge in some conditions and not others, and do
430 not confirm our earlier finding of a negative relationship between outcome and receiver mood (Sherwood et al., 2005). Previous promising findings with optimism and confidence of success were not confirmed here. It may be unrealistic to expect experimenter-participant interaction findings to replicate across studies given that the participants (and
435 experimenters in this case) are different, and efforts need to be made to ensure continuity across studies. In any case, in the present study there may have been limited opportunity for experimenter interaction effects to occur; with the switch from a Ganzfeld to a dream ESP design most of the recruitment and running of the study was conducted by telephone,
440 email and by post, so that there was a large reduction in the amount of face-to-face interaction until the judging session, by which time any psi performance is likely to be over (unless it occurs during judging). There might, therefore, have been little opportunity for the interaction to affect performance unless it could act retrospectively. Future tests of
445 interaction effects would need to ensure that there is sufficient interaction between the parties so as to provide an opportunity for this to be influential.

Acknowledgements

We would like to thank the Perrott-Warrick Fund, Trinity College,
450 Cambridge for their support of this project.

References

- Beloff, J. (1983). The repeatability problem in parapsychology. *Parapsychology Review*, 14, 1-4.
- 455 Braud, W. (1977). Long-distance dream and presleep telepathy, In J. D. Morris, W. G. Roll, & R. L. Morris (Eds.), *Research in Parapsychology 1976*. (pp.154-55). Metuchen, NJ: Scarecrow Press.
- Broughton, R. S., & Alexander, C. H. (1997). Autoganzfeld II: An attempted replication of the PRL Ganzfeld research. *Journal of Parapsychology*, 61, 209-226.
- 460 Child, I. L. (1985). Psychology and anomalous observations: The question of ESP in dreams. *American Psychologist*, 40, 1219-1230.
- Dalton, K., Steinkamp, F., & Sherwood, S. J. (1999), A dream GESP experiment using dynamic targets and consensus vote. *Journal of the American Society for Psychical Research*, 93, 145-66.

- 465 Eysenck, S. B. G., Eysenck, H. J., & Barrett, P. (1985). A revised version of the Psychoticism scale. *Personality and Individual Differences*, 6, 21-29.
- Honorton, C., Berger, R. Varvoglis, M., Quant, M., Derr, P., Schechter, E., & Ferrari, D. (1990). Psi communication in the ganzfeld: Experiments with an automated testing system and a comparison with a meta-analysis of earlier studies. *Journal of Parapsychology*, 54, 99-139.
- 470 Milton, J., & Wiseman, R. (1999). Does psi exist? Lack of replication of an anomalous process of information transfer. *Psychological Bulletin*, 125, 387-391.
- Roe, C. A., Sherwood, S. J., & Holt, N. (2004). Interpersonal psi: Exploring the role of the sender in ganzfeld ESP tasks. *Journal of Parapsychology*, 68, 361-380.
- 475 Roe, C. A., Sherwood, S. J., Luke, D. P., & Farrell, L. M. (2002). An exploratory investigation of dream GESP using consensus judging and dynamic targets. *Journal of the Society for Psychical Research*, 66, 225-238.
- Shapin, B. & Coly, L. (1985). *The repeatability problem in parapsychology*. New York, NY: Parapsychology Foundation.
- 480 Sherwood, S. J., & Roe, C. A. (2003). A review of dream ESP studies conducted since the Maimonides dream ESP programme. *Journal of Consciousness Studies*, 10, 85-109.
- Sherwood, S. J., Roe, C. A., Holt, N., & Wilson, S. (2005). Interpersonal psi: Exploring the role of the experimenter and the experimental climate in a ganzfeld telepathy task. *European Journal of Parapsychology*, 20, 150-172.
- 485 Sicher, F., Targ, E., Moore, D. & Smith, H. (1998). A randomised double-blind study of the effects of distant healing in a population with AIDS. *Western Journal of Medicine*, 169, 356-63.
- Stanford, R. G. (1993). Learning to lure the rabbit: Charles Honorton's process-relevant research. In R. K. Rao (Ed.), *Charles Honorton and the impoverished state of scepticism*. (pp. 129-176). Jefferson, NC: McFarland & Co.
- Thalbourne, M. A., & Delin, P. S. (1993). A new instrument for measuring the sheep-goat variable: Its psychometric properties and factor structure. *Journal of the Society for Psychical Research*, 59, 172-186.
- 495 Ullman, M., & Krippner, S., & Vaughan, A. (1973). *Dream telepathy: Experiments in nocturnal ESP*. Jefferson, NC: McFarland & Co.
- Utts, J. (1991). Replication and meta-analysis in parapsychology. *Statistical Science*, 6, 363-403.

Appendix

500 *Interaction questionnaire*

1. How would you rate your current mood?

(Negative — Positive)

2. How do you feel at this moment?

(Tense — Relaxed)

505 3. How do you feel about the prospect of participating in this experiment?

(Pessimistic — Optimistic)

4. How confident are you that today's experiment will be a success?

(Not at all confident — Extremely confident)

510 5. How would you describe the quality of rapport that you have with the Experimenter?

(Extremely poor — Extremely good)

6. How would you rate the quality of the interaction between experimenter and participants?

(Very cold — Very warm)

515 ('Rehearsed' — Spontaneous)

(Very negative — Very positive)