


Journal Pre-proof

Defining the importance of stress reduction in managing cardiovascular disease - the role of exercise

Dejana Popovic, Marija Bjelobrk, Milorad Tesic, Stefan Seman, Sisitha Jayasinghe, Andrew P. Hills, Abraham Samuel Babu, Djordje G. Jakovljevic, Lee Stoner, Cemal Ozemek, Samantha Bond, Mark A. Faghy, Nicolaas P. Pronk, Carl J. Lavie, Ross Arena, PIVOT Network



PII: S0033-0620(22)00015-9

DOI: <https://doi.org/10.1016/j.pcad.2022.01.008>

Reference: YPCAD 1259

To appear in: *Progress in Cardiovascular Diseases*

Please cite this article as: D. Popovic, M. Bjelobrk, M. Tesic, et al., Defining the importance of stress reduction in managing cardiovascular disease - the role of exercise, *Progress in Cardiovascular Diseases* (2021), <https://doi.org/10.1016/j.pcad.2022.01.008>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Elsevier Inc. All rights reserved.

Defining the Importance of Stress Reduction in Managing Cardiovascular Disease - The Role of Exercise

Dejana Popovic^{1, 12}; Marija Bjelobrk²; Milorad Tesic¹; Stefan Seman³; Sisitha Jayasinghe⁴; Andrew P Hills^{4, 12}; Abraham Samuel Babu^{5, 12}; Djordje G Jakovljevic⁶; Lee Stoner^{7, 12}; Cemal Ozemek^{8, 12}; Samantha Bond^{8, 12}; Mark A Faghy^{8, 9, 12}; Nicolaas P Pronk^{10, 12}; Carl J Lavie^{11,12}, Ross Arena^{8, 12} – On behalf of the HL - PIVOT Network

- 1 Division of Cardiology, University Clinical Center of Serbia, Belgrade, Serbia;
University of Belgrade, Faculty of Pharmacy, Belgrade, Serbia
- 2 Division of Cardiology, Institute of Cardiovascular Diseases Sremska Kamenica;
University of Novi Sad, School of Medicine Novi Sad, Serbia
- 3 University of Belgrade, Faculty of Sports and Physical Education, Belgrade, Serbia
- 4 School of Health Sciences, College of Health and Medicine, University of Tasmania,
Launceston, Tasmania
- 5 Department of Physiotherapy, Manipal College of Health Professions, Manipal
Academy of Higher Education, Manipal, India
- 6 Institute for Health and Wellbeing (CSELS), Faculty of Health and Life Sciences,
Coventry University, UK; Translational and Clinical Research Institute, Faculty of
Medical Sciences, Newcastle University, UK
- 7 Department of Exercise and Sport Science, University of North Carolina at Chapel
Hill, Chapel Hill, NC 27599, USA
- 8 Department of Physical Therapy, College of Applied Health Sciences, University of
Illinois at Chicago, Chicago, IL, USA
- 9 Human Sciences Research Centre, University of Derby, Derby, UK

- 10 Health Partners Institute, Minneapolis, MN, USA; University of Minnesota School of Public Health, Minneapolis, MN, USA
- 11 John Ochsner Heart Vascular Institute, Ochsner Clinical School-the University of Queensland School of Medicine, New Orleans , LA
- 12 Healthy Living for Pandemic Event Protection (HL – PIVOT) Network, Chicago, IL

Conflict of interests: The authors have no conflicts of interest to disclose.

Address for correspondence:

Dejana Popovic

Department of Cardiovascular Diseases, Mayo Clinic, Rochester, Minnesota

Address: 200 First Street SW, Rochester, MN 55905

Tel.: +15073199637

E-mail: Popovic.Dejana@mayo.edu

Abstract:

Traditional risk factors for cardiovascular disease (CVD) have long been the focus of preventive strategies. The impact of family stress, depression, anxiety, hostility, pessimism, job strain, social isolation, lack of purpose in life and social support, are well recognized risks for CVD development, however they are under-appreciated in clinical practice guidelines. The purpose of this article is to review the impact of acute and chronic stress on CVD risk, elaborate repositioning in guidelines, with emphasis to approaches for stress reduction. Regular exercise, both aerobic and resistance, leads to better adaptiveness to other types of stress, however, it remains unknown whether the total amount of stress one can receive before negative health effects is unlimited. Evidently, marked reductions in stress related disorders are shown following

formal cardiac rehabilitation programs. Attendance of cardiac rehabilitation is highly recommended for the stress-related mortality risk reduction. Innovative approaches to offset the broad challenges that CVD pose, augmented by sustained exposure to stress, are desperately needed, but hindered by a lack of successful population-level interventions that promote lasting change.

Keywords: Stress, Cardiovascular Diseases, Exercise

Abbreviations:

ANS - autonomic nervous system

BDNF - brain-derived neurotrophic factor

CAD - coronary artery disease

CR - cardiac rehabilitation

CRF - cardiorespiratory fitness

CV - cardiovascular

CVD - cardiovascular disease

HPA - hypothalamic-pituitary-adrenocortical

HRV - heart rate variability

MI - myocardial infarction

PA - physical activity

PS - psychological stress

PSS - Perceived Stress Scale

PTSD - post-traumatic stress disorder

SAM - sympatho-adrenomedullar

SIC - stress-induced cardiomyopathy

SNS - sympathetic nervous system

RAAS - renin-angiotensin aldosterone system

VO₂ - oxygen consumption

Stress is defined as a state of homeostasis being challenged (1,2). While acute stress is known as short-term stress, chronic stress is defined as “long-term” stress, characterized by an ongoing specific environmental condition or stressor with enduring impact (3). Stress effects are biological consequences resulting from the struggle with stressors (4–6). It might include reestablishing homeostasis to promote health (good stress - eustress) (1,2). Conversely, sustress (inadequate stress) or distress (bad stress) can cause damage to the body or even disease (1,2).

Stress conditions, both acute and chronic, appear to be strong risks for cardiovascular (CV) disease (CVD) development, however they are hugely under-considered and under-appreciated in clinical practice (1-3). The impact of family stress, depression, anxiety, outbursts of anger, hostility, pessimism, job strain and insecurity, purpose in life, social support, early life stress and social isolation are well recognized in the pathophysiology of stress disorders (1-3). On the other hand, physical activity (PA) has long been used to reduce stress, but its direct link to stress and CVD is still insufficiently emphasized.

In this review, we will analyze the impact of stress on CV health, stress related mechanisms that promote atherosclerosis and risk factors for unwanted stress reaction. Also, we will address the current place of stress as a risk factor in clinical practice guidelines and its reclassification potential, as well as approaches for stress reduction with emphasis on the role of

PA, exercise and improvements in cardiorespiratory fitness (CRF) in the context of CVD development.

Mechanisms of Stress Reaction

Responses to stressful situations are generally regarded as reactions of the organism to accommodate or compensate for stress (1,2). This reaction is previously described as an activation of the sympathoadrenal system (SAM) and the hypothalamic-pituitary-adrenocortical (HPA) axis (2,7–10). This response occurs during both exercise as well as in psychological stress (PS) (1,2,7–10). Interestingly, there is a delayed response of the HPA axis observed during exercise in comparison to PS and points to involvement of different neurobiological and cognitive emotional mechanisms (8,10,11). However, the activation of the stress system also exhibits marked variability, influenced by gender, age, race, genetic factors, body composition, type of stress and PA (6,8,12–14).

Previous reports have shown the relationship between physical stress and HPA axis activation, which deals mostly with the influence of PA on stress hormones levels (6,8,10). Furthermore, it was shown that higher levels of PA are associated with lower HPA axis reactivity to PS (15,16). As a result, PA has beneficial effects on both physical and mental health, affording protection against the detrimental consequences of chronic stress and stress-related diseases, such as CVD and depression (10,17). For example, it was previously shown that elite athletes demonstrated markedly reduced physiological and PS responsiveness compared to untrained subjects (10). Highly trained individuals have reduced salivary free cortisol and heart rate responses to a PS compared to untrained individuals (10). Moreover, trained individuals were generally calmer and exhibited better mood and lower anxiety throughout stress exposure compared to untrained individuals who showed an increase in anxiety and a decrease in mood

and calmness during stress (10). Since chronic exposure to physical stressors affords redundant activation of the HPA axis and the sympathetic nervous system (SNS), the authors posited that stress reactivity of elite athletes might be explained by an exercise-induced modulation of the hormonal stress-responsive and autonomic nervous systems (ANS)(10). These findings are in line with other studies which report that aerobic training leads to lower CV reactivity to PS (17,18). Furthermore, exercise-induced adaptations may also mitigate the responsiveness to other stressors, such as psychosocial stressors (10). It has also been shown that children and adolescents who are more physically active not only had more optimal physical health but also enjoyed better mental health (19). In this context, a better understanding of stress-responsive physiological systems can lead to the development of multidimensional prevention programs against stress-related disorders, which combine physical and cognitive coping strategies.

Impact of Stress on CV Health and Stress Mechanisms Promoting Atherosclerosis

Stress conditions increase the risk for atherosclerosis and related diseases, including stroke and myocardial infarction (MI) (20). The effect of stress on atherosclerosis involves multiple complex mechanisms that remain to be fully elucidated (21).

The response to stress involves the activation of the autonomic nervous system (ANS) and HPA which adversely affect the CV system both by accelerating the atherosclerotic process and by precipitating the occurrence of a CVD or cerebrovascular event (22,23). Whereas the pathophysiological effects on the progression of atherosclerosis are likely to involve repeated or chronic exposure to stress, stress-related triggering of events among people with an already high atherosclerotic plaque burden might also be a consequence of acute stress responses (22,23). In acute CVD events, excessive levels of circulating catecholamines are known to be a contributing factor to disruption of the fibrous cap of atherosclerotic plaque, coupled with active

inflammation and hypercoagulability and followed by electric instability (24). Stress-related clinical events include ventricular tachycardia and ventricular fibrillation, atrial fibrillation, stroke, MI and other forms of acute coronary syndromes (i.e., unstable angina and stress cardiomyopathy). Relevant pathophysiological changes, in addition to stress-induced ischemia, include: 1) an increase in the levels of pro-inflammatory cytokines and adhesion molecules that contribute to atherosclerotic plaque destabilization and monocyte chemotaxis; 2) release of tissue factor; 3) increased blood viscosity; 4) platelet and coagulation activation; 5) systemic vasoconstriction and an increase in arterial blood-pressure levels; 6) increase in sinus node firing rate and atrioventricular conduction velocity; 7) change in the balance between sympathetic and parasympathetic cardiac control; and 8) increase in myocardial oxygen consumption (22,23).

Previous studies implicate chronic stress as an important driver of metabolic syndrome, diabetes, and CVD development (25). The INTERHEART 128 study (29,972 participants from 52 countries) and the INTERSTROKE study (26,919 participants from 32 countries) both identified PS as one of ten modifiable risk factors for CVD onset. Namely, the activation of the HPA and the SAM axis, provokes the release of hormones such as adrenocorticotropic hormone, cortisol, growth hormone, prolactin, epinephrine and norepinephrine (25). Increased cortisol has many somatic effects, including the precipitation of insulin resistance, increased blood pressure, central redistribution of adiposity and an impaired immune response (26). Increased circulating levels of catecholamines additionally promote insulin resistance, increased blood pressure and heart rate, and systemic inflammation by elevating production of inflammatory cytokines (intercellular adhesion molecule-1, acute phase reactant C-reactive protein and proinflammatory cytokine interleukin-6), which are significantly heightened in chronic stress. Chronic stress can suppress or dysregulate innate and adaptive immune responses by altering the type 1/type 2

cytokine balance, thereby inducing low-grade inflammation and suppressing the function of immuno-protective cells (27). The increase of inflammatory cytokines and the expression of adhesion molecules via certain pathways can induce mononuclear cell aggregation and lymphocyte adhesion (27). Furthermore, chronic stress changes the homeostasis of the sympathetic and vagal nervous systems. Autonomic disorders caused by chronic stress may be a common mechanism that increases risk (27,28). The SNS activation triggers increased vascular tone, myocardial oxygen consumption, platelets activation, as well as activation of renin-angiotensin aldosterone system (RAAS), which results in endothelial dysfunction, and hence, the development of CVD (27,28).

Recent studies suggest that psychological stress triggers endothelial dysfunction (29). As previously stated, psychological stress activates the SNS, RAAS and the HPA axis, triggering increased levels of catecholamines, glucocorticoids and angiotensin II, as well as pro-inflammatory cytokines, respectively. All these mediators can promote endothelial dysfunction due to the increase in oxidative stress and circulating inflammatory mediators, as well as reduced nitrogen-monoxide production and bioavailability. These alterations lead to vascular inflammation and, consequently, the development of atherosclerosis. Furthermore, tumor necrosis factor-alpha and mitogen activated protein kinases signaling pathway play a potential role in the inflammatory process, as well as the adrenergic receptors and CXCL12 chemokine that are involved with hematopoietic activation and the development of atherosclerotic lesions (29).

Several neural centers in the brain's salience network play a critical role in an organism's response to its environment (20,21). The amygdala is one important component of this network that is involved in the emotional perception and physiological response to stressors. As such, the

amygdala could be a critical structure in the central nervous system mechanisms linking stress to CVD (20,21).

As an impressive example, stress has been linked to stress-induced cardiomyopathy (SIC), a transient systolic and diastolic left ventricular dysfunction with wall-motion abnormalities considered an acute heart failure syndrome (30). SIC is a self-resolving syndrome that may present as a mimic to ischemic heart disease or as a complication during critical illness, particularly those diseases involving the neurohumoral axis (31,32). This syndrome has gone by many names in the literature, including “cardiac syndrome X”, “broken heart syndrome”, “ampulla cardiomyopathy”, “tako-tsubo cardiomyopathy”, “transient left ventricular apical ballooning syndrome”, “neurogenic stunned myocardium”, and “neurogenic stress cardiomyopathy” (32,33). It was initially described in the Japanese population in the early 1990s, as a transient cardiomyopathy frequently following stressful events, with the absence of coronary artery disease (CAD) (31,33). Perhaps the best definition presently available is the absence of any established cause of reversible myocardial dysfunction. The majority of patients (80–90%) are women, predominantly postmenopausal women. Many cases (14–70%) are preceded by a stressor, either emotional or physical. In patients without another identifiable cause, a recent acute emotional stress frequently precedes the syndrome. SIC is relatively common in patients with severe neurologic injury, as well as in other situations where catecholamines are clearly elevated, such as in the setting of a pheochromocytoma, status epilepticus, tetanus, sepsis, systemic lupus erythematosus, anorexia nervosa, thyrotoxicosis and diabetic ketoacidosis (34–43). The presumed etiology of SIC is excessive SNS stimulation of the myocardium, likely mediated by the brain. Catecholamines may lead to cardiac dysfunction via microvascular dysfunction or direct toxicity (44). Hormones also may play a role in the pathogenesis of SIC.

Estrogen may attenuate the CV response to catecholamines, explaining the higher frequency of SIC in postmenopausal women (35,45). Patients present with chest pain and dyspnea. Pathologic electrocardiogram and cardiac enzyme changes are present (31,35). Echocardiography and contrast ventriculography detect a dyskinesia with a ballooning of the left ventricular apex (46). The classic pattern of wall motion abnormalities seen on echocardiogram in SIC is one of apical ballooning or hypokinesia of the apex of the heart with hypercontractility of the basal segments. In one-third of all patients, a subtle dyskinesia of the right ventricular apex is also observed (46). However, coronary angiography shows no evidence of a relevant epicardial obstructive disease (31). The natural course of SIC syndrome is near-complete resolution over several days to weeks. A large study of patients presenting with acute coronary syndrome and a pattern of apical ballooning on echocardiogram with normal coronary angiography found no increase in 4-year mortality compared with matched controls and a 1–2% annual recurrence rate (47,48).

Types of Stress and CV Health: Impact of Family Stress, Depression, Posttraumatic Stress Disorder, Anxiety, Outbursts of Anger, Hostility, Pessimism, Job Strain and Insecurity, Long Working Hours, Purpose in Life, Social Support, Early Life Stress and Social Isolation

Numerous factors, such as lifestyle behaviors, psychosocial variables, personality types, and mental illnesses, can have a direct impact on the etiology of CVD. Importantly, these factors can also precipitate or worsen pre-existing CVD conditions (49–52).

A growing body of evidence indicates an inverse relationship between marital quality and CAD morbidity and mortality (53). There is a link between marital quality and carotid artery intima-media thickness and coronary artery calcification burden – common indicators of

atherosclerosis (54,55). In addition, recent reports indicate marital status can influence individual CV reactivity and subsequent manifestations of unwanted blood pressure patterns (56). Similarly, associations between depression/depressive symptoms, post-traumatic stress disorder (PTSD), anxiety, and anger/hostility are well documented (57–59). An individual with depression without prior CVD is 2.7 times more likely to die from CAD over a median follow-up of 8.5 years (60), and 20–30% of CVD patients generally have depression (61). Mechanistically, excessive SNS and/or reduced parasympathetic modulation (62), blunted inhibitory feedback in the HPA axis (63), elevated levels of platelet factor 4 and β -thromboglobulin (64), endothelial dysfunction (65) and increased CRP, tumor necrosis factor- α and interleukins (66) in depression, are considered the main causes of cardiovascular anomalies. The latter includes high blood pressure, reduced heart rate variability (HRV), and worsening of atherosclerosis.

Similarly, the association between characteristics of PTSD, including reliving the traumatic event, avoidance of reminders, hyperarousal, anhedonia, detachment, restricted affect, sleep difficulty, intense negative emotions, and potential CVD is incontrovertible (58,67). Physiologically, it is likely that a combination of changes in brain activity, somatic nervous system, endothelial dysfunction and inflammatory reaction are involved in a plausible link between PTSD and CVD (68). For instance, autonomic imbalance manifested as low HRV (69) and high resting heart rate (70), HPA axis dysfunction reflected as low diurnal cortisol levels (71), increased platelet activation (72), endothelial dysfunction manifested as significant reduction in flow mediated dilatation (73) and inflammatory perturbations with significantly high levels of IL-1 β , IL-6, and interferon γ (74), can be directly associated and/or lead to exacerbation of existing CVD. In addition, behavioral changes that may accompany traumatic events

including, but not limited to smoking, inactivity, obesity, and nonadherence to medications, can also increase the potential for development of CVD (75,76).

Anger and hostility, typically associated with Type A personality traits, and comprising numerous constructs such as physiological, cognitive, phenomenological, and behavioral variables, have been linked with ill-health since antiquity (77). A substantial body of evidence indicates a positive relationship between anger and hostility with increased CVD events in healthy populations, as well as in patients with existing CVD (78,79). Excessive (and in some instances, paradoxical) vasoconstriction mediated by endothelial dysfunction (80), pronounced adrenocorticotrophic responses to socially salient stress (81), and increased intima media thickness (82), are some of the mechanisms linking anger and hostility with potential CVD.

Although pessimism and optimism do not anchor the two ends of one spectrum (83), empirical evidence suggests that the former is associated with increases in CAD, stroke, all-cause mortality (84–86), and the latter with increased protection against CVD (87,88). Many of the protective effects of optimism can be attributed to indirect, positive behavioral changes including increased PA levels, better diet quality, and reductions in smoking (87). A substantial evidence base support associations between pessimism and pathophysiologic dysfunction, mediated via neuro-endocrine (i.e., HPA axis), nervous system (i.e., blood pressure), immune system, genetic, endothelial, and metabolic function modulations (89–92).

Demand for increased productivity and efficiency has increased, leading to increased pressure upon employees and workforces globally (93). Whilst high levels of mental and physical wellbeing are closely associated with increased performance (94), sustained pressures, contractual requirements and incentivized employment opportunities relating to performance have an adverse effect upon work-life balance, absenteeism, and physical and mental wellbeing

(93). Prolonged sedentary profiles are a well-defined risk factor and the product of increased working hours, a reduction in active job roles, increased technological innovation and more recently remote working (95). Stress associated with work has also been identified as a major risk factor for CVD, in a review of 600,000 workers from 27 cohort studies, Kivimäki & Kawachi suggest that stressors from employment such as pressure relating to success, job strain and long working hours (>50 hours per week) increase the incidence of developing CAD and stroke by up to ~40% (96) compared with those free of such stressors. Poor work-life balance caused by job strain and long working hours is detrimental to mental and physical wellbeing, it also associated with broad social impacts that are linked to a 1.5 increased risk of experiencing social isolation (97) and CVD events.

A rapidly growing literature base has also identified and increased the understanding that positive psychological wellbeing can protect against adverse physical health outcomes, including CVD (98). A meta-analysis of 10 prospective studies by Cohen et al demonstrated that having an increased sense of purpose (directed and motivated by valued life goals) is associated with reduced CVD risk (99). Whilst more work is needed to determine the associated mechanisms, the continued presence of stress and the associated biological functions are associated with negative implications for healthy ageing and pre-disposing individuals and increased CVD risk factors (100). The concept of healthy ageing is affected by exposure to ‘early life stress’ which influences the biological responsiveness to future stress and is associated with increased incidence of developing a broad range of chronic conditions such as CVD later in life (101). Whilst the exact mechanistic details are not understood in their entirety, broad associations have been determined in both central and peripheral systems. Exposure to stress in early life alters the biological programming of the cardiovascular system inducing lasting effects on the

responsiveness to stress stimuli, pre-disposing stress during early life as an independent risk factor (102).

Due to the broad risk posed to global health systems and the increasing economic costs, currently estimated at \$351.2 billion per annum (103), innovative approaches to offset and combat the broad challenges that are posed by CVD, which are augmented by sustained exposure to stress, are desperately needed. Approaches to reducing sedentary behaviors and the promotion of sustained activity profiles and healthy lifestyles are well documented but hindered by a lack of successful population-level interventions that promote lasting change. Laughter and enjoyment have been shown to reduce stress and by consequence CVD risk (104) and must be considered in the development of innovative lifestyle interventions that seek to reduce stress, instill work-life balance, and promote increased mental and physical health and wellbeing.

Stress as a Risk Factor in Current Guidelines and Risk Estimation Systems - Reclassification Potential?

Traditional risk factors for CVD have long been the focus of preventive strategies. However, they interplay with psychological factors which additionally strongly impact outcomes. A recent population study identified the impact of various psychological disorders on the incidence of CVD (105). They identified that stress related disorders were strongly associated with early onset CVD in those below and above the ages of 50 years [hazard ratio 1.40 (1.32 to 1.49) and 1.24 (1.18 to 1.30), respectively]. The INTERHEART study demonstrated exposure to long-term stress added to the risk of acute MI even after considering combined exposure to conventional CVD risk factors, with odds ratios increasing from 69-183, when participants were exposed to psychosocial adversity (106). The INTERHEART study also found that psychosocial factors were one among the top three factors contributing to presentation of acute coronary syndrome.

More recently, in the Young Finns study, the authors found favorable childhood experiences to be associated with lower coronary artery calcium scores when assessed 28 years later (107). In adulthood, workplace stress has also been seen to contribute to the risk of incident CVD, by an increase of 40% (108).

Copenhagen City Heart Study demonstrated the higher impact of vital exhaustion than other risk factors for coronary artery disease development (109). Moreover, the relationships between high job strain and incident CVD were only minimally explained by the traditional CVD risk factors in 22086 subjects participating in the Women's Health Study (110), whereas depressive/anxious symptoms explained the highest proportion of the observed high strain-CVD association (21%). All these population studies strengthen the physiological basis that the heart and mind are greatly interlinked with one another and can have a tremendous impact on CV health (111). However, the quantitative data on relative impact of stress versus other traditional risk factors in development of every particular CVD are generally lacking and further studies in this direction are highly warranted.

Based on evidence, it is apparent that stress is a major contributor to CVD, however, given attention in the current guidelines hugely underestimates its role. Many years ago, the World Health Organization recommended the need for clinical history related to psychosocial factors, however, did not provide any evidence-based recommendations on its management (112). The European guidelines provide a class IIa, Level B recommendation for the assessment of psychosocial risk factors along with class I, level A recommendation for multimodal behavioral interventions in those with CVD (113). However, treatment of these conditions with the aim of preventing CVD was a class IIa, level B recommendation. The more recent ACC/AHA guidelines did consider the social determinants of health and did recommend the need for regular

assessment of psychosocial stressors and their treatment through counselling (114). However, no further details were provided.

Contrary to the CVD prevention guidelines, cardiac rehabilitation (CR) statements, however, have emphasized the need for addressing psychosocial components and have made it a core component (115,116). Even guidelines focusing on low-resource settings have emphasized this need (117). Despite this being a core component, the number of programs around the world delivering this component remains low (118,119). This could be due to the limited number of mental health professionals around the world available to provide such care (118).

There are three aspects of the stress experience that complicate its inclusion as a routine consideration in clinical practice. First, as discussed before, stress is a ubiquitous aspect of the human experience, ranging from benign daily stress (e.g., being stuck in traffic) to stress which manifests from lifelong discrimination. Second, the mechanisms by which excessive stress exposure increases CVD risk are multi-factorial and have not been fully elucidated. Using the examples above, regular but acute stress in the workplace may partially increase CVD risk through autonomic nervous system (ANS) dysregulation. Whereas stress resulting from lifelong discrimination may lead to chronic low-grade inflammation and advanced cellular aging (120). Third, both the magnitude of stress and the physiological effects of stressors are likely dependent on resiliency and coping strategies. Using the workplace example, one individual may internalize negative criticism from a line manager and ‘cope’ by engaging in other harmful behaviors (e.g., smoking or binge eating), whereas as second worker may, for example, have learned to better communicate with the line manager, go for a walk, or practice mindfulness. In this way, one can consider that it is not just the stressor that is the CVD risk factor, but rather the collective perception of, reaction to, and subsequent physiological reactivity from the stressor.

While stress is a complex CVD risk factor, it does not mean we are unable to help a given individual. But the operative word is ‘individual’. There is no one scale that can represent the stress by all individuals. Similarly, there is unlikely to be one physiological biomarker to encapsulate the effects of stress on CV health. However, stress as a general entity can be subjectively measured, with an example scale being the Perceived Stress Scale (PSS) (121). The 10-item PSS is one of the most widely administered scales for measuring perceptions of stress over the past month, and has been associated with poor health outcomes, including depression and CVD (122,123). As such, this scale can be used to monitor perceptions of stress and to track reductions that may occur through changing in lifestyle or the incorporation of other stress management strategies.

Strategies that have been used to enhance stress management have traditionally focused on mind-body interventions in which there is often both a PA as well as a cognitive-behavioral/mental component (124, 125). Yoga and tai chi are examples of mind-body therapies which incorporate both physical and mental components (126, 127). Both PA and mental (e.g., cognitive-behavioral) approaches such as mindfulness meditation have been independently linked to reduced stress, or more precisely – reduced perceptions and negative reactivity from stress (128–130). Further work is needed to determine if there is a synergistic effect from combining physical and mental/cognitive stress-reduction strategies, and whether certain strategies should be focused on specific subsets of the population.

Hormonal Background of Psychological and Physical Stress – Is it the Same?

Both physical and PS are primarily mediated via co-activity of the autonomic nervous system (ANS) and a constellation of HPA axis (131). While responsiveness (i.e., magnitude and time course) of both systems is highly stressor specific, the outcomes of the former are realized

through a multitude of autonomic pathways (e.g., SAM system, dopamine system, etc.) whereas outputs of the latter are modified by input from intricate brain regions such as locus coeruleus, amygdala, anterior cingulate cortex, paraventricular nucleus and the hippocampus (131,132). Amongst the effector hormonal milieu that determines the overall stress response, epinephrine/norepinephrine (SAM) and cortisol (HPA) can be considered the ‘major players’, at least in the peripheral tissues (133).

The stress response is both natural and adaptive with optimally reacting systems, and by activating various feedback loops, can return the body to a steady state once the initial threat has subsided (134). The magnitude of the stress response is a key component in one’s progression to ill-health, and existing evidence indicates that a “Goldilocks” response (i.e., intermediate, not too high or low) is the most adaptive and resilient (135). In contrast, both exaggerated and blunted reactivity of the autonomic and hypothalamic pathways are associated with some of the most pervasive CVD conditions such as hypertension, coronary artery calcification and increases in carotid intima media thickness (136–141). These responses are suggestive of an inverted-U shaped association between stress reactivity and disease.

Additionally, duration of exposure is also a key factor that determines adaptiveness or deleterious stress (133). Bursts of short-term stress (i.e., stress that lasts from minutes to hours) interspersed with adequate time for system recovery is thought to be associated with salubrious adaptive effects (142). In contrast, chronic stress (e.g., stress that persists for several hours a day for weeks/months) can lead to maladaptation such as dysregulation of circadian cortisol rhythm (143,144). Depression – which is strongly linked with CVD – is a classic example of ‘stress pathway overdrive’ leading to deleterious health outcomes (145). For instance, existing literature is replete with examples of laboratory stressors such as the Trier Social Stress Test – which

activates the same corticolimbic circuitry implicated in depression (146) – simulating the hyperactivity of corticotropin releasing hormone-adrenocorticotropin-cortisol (i.e., HPA axis) system, that is commonplace in major depression etiology (145,147).

Exercise, as a form of physical stress, shares several characteristics of an acute stressor, and challenges both nervous and endocrine arms of the stress machinery (148). For instance, exercise of sufficient dose (i.e., volume and intensity) can increase circulating levels of many stress-related factors including cortisol, epinephrine/norepinephrine and inflammatory cytokines (149,150).

The Role of Exercise in Managing and Mitigating Stress

Particularly noteworthy, as postulated in the cross-stressor adaptation hypothesis, regular exercise can result in adaptations that lead to reduced sensitivity to subsequent heterotypic (other than exercise) stressors (151). A substantial body of evidence indicate stress-buffering effects (both autonomic and hypothalamic pathways) of regular exercise (152,153).

The 2018 Federal Physical Activity Guidelines Advisory Committee's Scientific Report determined that there was a strong body of evidence supporting the anxiolytic effects of acute and regular physical activity in adults and older adults (154). This was an update to the 2008 report which had concluded that there was insufficient evidence available to make such an observation. Although there is still a relatively small pool of studies in this area compared to more prominent topics (i.e., cardiovascular and skeletal muscle health/function), a growing body of work consistently supports the beneficial effects of both aerobic and resistance exercise in lowering symptoms of stress (155,156). Most randomized controlled studies investigating the effects of aerobic exercise on stress have employed moderate continuous aerobic exercise at volumes comparable to the Federal Physical Activity guidelines (i.e., 70% of peak VO_2 , ~30

minutes per session, 3-5 days per week) (157–159) and therefore, little is known about the potential graded effects of higher intensities and/or volumes of aerobic exercise on stress. However, a previous meta-analysis found a dose-response relation between CRF and the risk of common mental health disorders (160). With higher intensities and volumes of exercise leading to greater improvements in cardiorespiratory fitness (161), it is feasible that chronic high-volume exercise may promote more favorable decreases in stress symptoms over time. Future studies are needed to determine whether this holds true.

Regular resistance exercise training (i.e., at least two days of muscle strengthening activity per week) is a vital component of the PA guidelines as it slows the loss of muscle mass with aging, reduces risk of serious falls and helps individuals maintain their physical independence (154). Moreover, a meta-analysis by Gordon and colleagues concluded resistance exercise may lead to similar improvement in markers of stress as aerobic exercise in chronically ill patients and those with anxiety related disorders (155). Furthermore, initial evidence suggests that resistance exercise intensity may modulate the degree of improvement in stress related disorders in older adults (162). A 12-week resistance exercise training intervention conducted by Tsutsumi and colleagues (162) found greater strength gains in those completing vigorous resistance exercise training (75 to 85% of one repetition maximum, 8-10 repetitions) than those randomized to moderate intensity (55 to 65% of one repetition maximum, 14 to 16 repetitions), but found that the moderate intensity group had a significantly greater decrease in anxiety. Suggesting that moderate intensity resistance exercise training may be more beneficial in improving psychological health. Additional studies will be needed to confirm this.

To date, much uncertainty remains about the mechanism that explain the anxiolytic effects of aerobic and resistance exercise, though certain neurobiological and psychological

mechanisms have been proposed. For example, it is evident that stress-induced depressive pathology contributes to altered brain-derived neurotrophic factor (BDNF) level and, thereby, disruptions in neuroplasticity (163). Conversely, physical activity optimizes BDNF in key brain regions, and as such promotes neuronal health. Perhaps the most recognized value of physical activity is the increase in serum BDNF after acute exercise (i.e., 70% of peak VO_2 for 30 min) (163,164). Improvements in self-efficacy and negative affect may play a key role in decreasing symptoms associated with stress disorders with exercise (165). It is important to note that many exercise interventions have been conducted in group settings which may provide an inadvertent support group atmosphere and facilitate coping stress (166). Nevertheless, with previous studies (167) showing that a single session of aerobic exercise can lower stress levels may suggest that this occurs independent of the potential effect of exercising in a group. Certainly, an individual's exercise preferences and response to group settings may modify stress levels. Accordingly, additional work in this area is needed to establish individualized recommendations on exercise settings relative to individual preferences and responses. There is however sufficient evidence to support the promotion of aerobic and resistance exercise as a method of lowering stress.

Of note, despite the 'stress-busting' effects, excessive doses of exercise can be counterproductive and cause unintended harm by chronically exposing the body to stress hormones (150,168,169). However, given the widespread availability of labor-saving technologies and unprecedented levels of sedentarism spanning most physical activity domains, it is unlikely that significant numbers of individuals will engage in excessive amounts of exercise (170,171).

CR, PS, and Survival

As mentioned above, assessment and management of PS has become a major component of CR programs (172,173). Lavie and colleagues (174) demonstrated that CAD patients have high levels of PS, including depression, anxiety and hostility, with marked reductions in PS following formal CR programs. Milani and colleagues (175, 176) demonstrated that CR patients with depression have 4-5 fold increased mortality compared to those without depression, but depressed CHD patients who attended formal CR had 70% reductions in mortality compared with depressed patients who did not attend CR. Those patients who did not improve CRF with CR continued with high prevalence of depression post-CR and high mortality risk, whereas those who had either small or moderate improvements in CRF with CR had marked reductions in both depression and all-cause mortality. These benefits were even more noted in those with systolic heart failure. Although these benefits were noted with standard CR programs (177), Blumenthal and colleagues (178) noted even greater reductions in CVD events in those with enhanced CR (regular CR with stress management training). Greater efforts are needed to improve PS management, especially depression, and attendance in CR programs for the management of patients with CVD (173, 176, 179).

COVID-19 Pandemic, PS, and Potential for Exercise

Currently, we are still almost 2 years into the COVID-19 pandemic, and we have discussed the very marked effects of the COVID -19, physical inactivity and obesity pandemics/epidemics colliding (180-182). Additionally, COVID-19 has resulted in profound PS at so many areas in our society, which has probably worsened the obesity and physical activity habits (183). Clearly, PA and higher CRF has been associated with significantly less severe complications from COVID-19 (184-186). As we discussed previously with natural disasters (187), there is certainly

great probability that regular PA and exercise would lead to reductions in PS and improvements in prognosis in COVID-19 and future pandemics, as it does with CVD (172, 186).


Conclusion

In summary, stress is a complex and highly subjective construct. Acute and chronic stress are hugely underestimated as risk factors for CVD in clinical practice, and as such there is a lack of recommended coping strategies that may reduce stress-related diseases burden. There is not a one-size fits all approach to managing stress, and any management strategies should be mindful of individual traits and lived experiences. More attention is needed to better understand mechanisms of stress-related negative health outcomes through research, as evidence is needed to confirm the obvious role of stress in overall health and disease. Nonetheless, there are means by which to monitor stress and strategies which are reported to be effective in managing stress. Regular exercise, both aerobic and anaerobic, leads to better adaptiveness to stress (Central Illustration). However, questions remain regarding the total amount of stress one can receive before negative health effects are experienced.

STRESS & EXERCISE



Stress, whether short-term or long-term, leads to biological consequences, which may precipitate a cardiovascular event.



STRESS:

A state of **homeostasis** being challenged; a reaction to accommodate an ongoing stressor.

The response to stress accelerates atherosclerosis, or the buildup of plaque in the arteries.^{1,2} Stress can also trigger cardiac events for those with preexisting plaque buildup.³



ACUTE

Short-term

CHRONIC

Long-term

The stress response of both types lead to an activation of the **autonomic nervous system and HPA axis**.³

CHRONIC STRESS

AFFECTIVE DISORDERS

EVOKED CENTRAL RESPONSES

- Increased HPA
- Increased SNS
- Increased altered behaviors
- ANS dysfunction
- Insulin resistance
- Obesity
- Hypertension
- Inflammation
- Platelet activation
- Endothelial dysfunction
- Ovarian dysfunction
- Lower bone density
- Somatic effects

Many different factors from the environment and a person's background can influence the stress reaction.

STRESSORS

- Marital quality
- Social deprivation
- Vital exhaustion
- Trauma
- Demanding jobs
- Purpose in life
- Socioeconomic status
- Public speaking
- Immunological Stress

&

RISK FACTORS

for unwanted stress reaction:

- Depressive mood
- Anxiety
- Outbursts of anger
- Pessimism
- Hostility
- D-type personality
- Nutrition
- Body composition
- Physical fitness

HORMONAL BACKGROUND

Chronic stress can suppress the immune response and subsequently alter the cytokine balance.⁴ Relatedly, psychological stress activates the sympathetic nervous system, renin-angiotensin system, and the HPA axis. These triggering events have been suggested to promote endothelial dysfunction.⁵



COPING STRATEGIES

To improve health in patients with CVD or risk factors for CVD, multimodal behavioral interventions are recommended as coping strategies, including physical exercise and psychological therapy.



THE ROLE OF EXERCISE

Exercise interventions have shown that even a single session of aerobic exercise can lower stress levels.^{1,4}



Citations

1. Enson I, Greenlee TA, Mohr RW, Petruzzello SA. META-ANALYSIS OF ACUTE EXERCISE EFFECTS ON STATE ANXIETY: AN UPDATE OF RANDOMIZED CONTROLLED TRIALS OVER THE PAST 25 YEARS. *Depress Anxiety*. Aug 2015;32(8):624-634.
2. Yao B, Meng L, Hao M, Zhang Z, Gong T, Guo Z. Chronic stress: a critical risk factor for atherosclerosis. *J Int Med Res*. 2019; 47(4): 1439-1440.
3. Mika Kivimäki and Andrew Steptoe, Effects of stress on the development and progression of cardiovascular disease. *Nature Reviews Cardiology*. 2015; 11: 215-223.
4. Dhabhar FS. Effects of stress on immune function: the good, the bad, and the beautiful. *Immunol Res*. 2014; 58:199-216.
5. Horvane Cöco and Ana Maria de Oliveira. Endothelial Dysfunction Induced by Chronic Psychological Stress: A Risk Factor for Atherosclerosis. *Cardiovasc Pharmacol open access*. 2015; 4:5.

Central Illustration. Both physical and psychological stress trigger stress response. While psychological stress leads to multiple disorders, physical allows adaptation to stress ameliorating effects of psychological stress.

References:

1. Lu S, Wei F, Li G. The evolution of the concept of stress and the framework of the stress system. *Cell Stress*. 2021;5(6):76-85.
2. Goldstein DS, Kopin IJ. Evolution of concepts of stress. *Stress*. 2007;10(2):109-120.
3. Hammen C, Kim EY, Eberhart NK, Brennan PA. Chronic and acute stress and the prediction of major depression in women. *Depress Anxiety*. 2009;26(8):718-723.
4. Rom O, Reznick AZ. The Stress Reaction: A Historical Perspective. *Adv Exp Med Biol*. 2016;905:1-4.
5. Peters A, McEwen BS, Friston K. Uncertainty and stress: Why it causes diseases and how it is mastered by the brain. *Prog Neurobiol*. 2017;156:164-188.
6. Popovic D, Popovic B, Flecas-Solarovic B, et al. The interface of hypothalamic-pituitary-adrenocortical axis and circulating brain natriuretic peptide in prediction of cardiopulmonary performance during physical stress. *Peptides*. 2013;47:85-93.
7. Yaribeygi H, Panahi Y, Sahraei H, Johnston TP, Sahebkar A. The impact of stress on body function: A review. *EXCLI J*. 2017;16:1057-1072.
8. Ahtiainen JP, Pakarinen A, Kraemer WJ, Häkkinen K. Acute hormonal and neuromuscular responses and recovery to forced vs maximum repetitions multiple resistance exercises. *Int J Sports Med*. 2003;24(6):410-418.

9. Di Luigi L, Guidetti L, Baldari C, Romanelli F. Heredity and pituitary response to exercise-related stress in trained men. *Int J Sports Med.* 2003;24(8):551-558.
10. Rimmel U, Zellweger BC, Marti B, et al. Trained men show lower cortisol, heart rate and psychological responses to psychosocial stress compared with untrained men. *Psychoneuroendocrinology.* 2007;32(6):627-635.
11. Martikainen S, Pesonen AK, Lahti J, et al. Higher levels of physical activity are associated with lower hypothalamic-pituitary-adrenocortical axis reactivity to psychosocial stress in children. *J Clin Endocrinol Metab.* 2013;98(4):E619-E627.
12. Garrido P. Aging and stress: past hypotheses, present approaches and perspectives. *Aging Dis.* 2011;2(1):80-99.
13. Charmandari E, Kino T, Chrousos GP. Primary generalized familial and sporadic glucocorticoid resistance (Chrousos syndrome) and hypersensitivity. *Endocr Dev.* 2013;24:67-85.
14. Pompili M, Serafini G, Innominata M, et al. The hypothalamic-pituitary-adrenal axis and serotonin abnormalities: a selective overview for the implications of suicide prevention. *Eur Arch Psychiatry Clin Neurosci.* 2010;260(8):583-600.
15. Gröpel P, Urner M, Pruessner JC, Quirin M. Endurance- and Resistance-Trained Men Exhibit Lower Cardiovascular Responses to Psychosocial Stress Than Untrained Men. *Front Psychol.* 2018;9:852.
16. Arvidson E, Dahlman AS, Börjesson M, Gullstrand L, Jonsdottir IH. The effects of exercise training on hypothalamic-pituitary-adrenal axis reactivity and autonomic response to acute stress-a randomized controlled study. *Trials.* 2020;21(1):888.

17. Spalding TW, Lyon LA, Steel DH, Hatfield BD. Aerobic exercise training and cardiovascular reactivity to psychological stress in sedentary young normotensive men and women. *Psychophysiology*. 2004;41(4):552-562.
18. Cardoso CG Jr, Gomides RS, Queiroz AC, Pinto LG, da Silveira Lobo F, Tinucci T, Mion D Jr, de Moraes Forjaz CL. Acute and chronic effects of aerobic and resistance exercise on ambulatory blood pressure. *Clinics (Sao Paulo)*. 2010;65(3):317-25.
19. Eime RM, Young JA, Harvey JT, Charity MJ, Payne WK. A systematic review of the psychological and social benefits of participation in sport for children and adolescents: informing development of a conceptual model of health through sport. *Int J Behav Nutr Phys Act*. 2013;10:98.
20. Kershaw KN, Lane-Cordova AD, Carnethon MR, Tindle HA, Liu K. Chronic Stress and Endothelial Dysfunction: The Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Hypertens*. 2017;30(1):75-80.
21. Johnstone SE, Baylin SB. Stress and the epigenetic landscape: a link to the pathobiology of human diseases?. *Nat Rev Genet*. 2010;11(11):806-812.
22. Yao BC, Meng LB, Hao ML, Zhang YM, Gong T, Guo ZG. Chronic stress: a critical risk factor for atherosclerosis. *J Int Med Res*. 2019;47(4):1429-1440.
23. Kivimäki M, Steptoe A. Effects of stress on the development and progression of cardiovascular disease. *Nat Rev Cardiol*. 2018;15(4):215-229.
24. Moreno PR. Pathophysiology of plaque disruption and thrombosis in acute ischemic syndromes. *J Stroke Cerebrovasc Dis*. 2001;10(2 Pt 2):2-9.
25. Glaser R, Kiecolt-Glaser JK. Stress-induced immune dysfunction: implications for health. *Nat Rev Immunol*. 2005;5(3):243-251.

26. Morais JBS, Severo JS, Beserra JB, et al. Association Between Cortisol, Insulin Resistance and Zinc in Obesity: a Mini-Review. *Biol Trace Elem Res.* 2019;191(2):323-330.
27. Dhabhar FS. Effects of stress on immune function: the good, the bad, and the beautiful. *Immunol Res.* 2014;58(2-3):193-210.
28. Chumaeva N, Hintsanen M, Ravaja N, et al. Interactive effect of long-term mental stress and cardiac stress reactivity on carotid intima-media thickness: the Cardiovascular Risk in Young Finns study. *Stress.* 2009;12(4):283-293.
29. Côco H, de Oliveira AM. Endothelial Dysfunction Induced by Chronic Psychological Stress: A Risk Factor for Atherosclerosis. *Cardiovascular Pharmacology: Open Access.* 2015;04(05).
30. Sanghvi SK, Harris DM. Anticoagulation and stress-induced cardiomyopathy. *J Thromb Thrombolysis.* 2019;47(1):1-7.
31. Pelliccia F, Kaski JC, Crea F, Camici PG. Pathophysiology of Takotsubo Syndrome. *Circulation.* 2017;135(24):2426-2441.
32. Farid A, Dufresne W, Farid B, Amsterdam EA. A Stressful Situation: Takotsubo Cardiomyopathy. *Am J Med.* 2018;131(3):253-256.
33. Boyd B, Solh T. Takotsubo cardiomyopathy: Review of broken heart syndrome. *JAAPA.* 2020;33(3):24-29.
34. Buchmann SJ, Lehmann D, Stevens CE. Takotsubo Cardiomyopathy-Acute Cardiac Dysfunction Associated With Neurological and Psychiatric Disorders. *Front Neurol.* 2019;10:917.

35. Gupta S, Goyal P, Idrees S, Aggarwal S, Bajaj D, Mattana J. Association of Endocrine Conditions with Takotsubo Cardiomyopathy: A Comprehensive Review. *J Am Heart Assoc.* 2018;7(19):e009003.
36. Ikenouchi-Sugita A, Yoshimura R, Hori H, et al. Plasma catecholamine metabolite levels and the activities of psychiatric symptoms in systemic lupus erythematosus. *Hum Psychopharmacol.* 2013;28(2):198-202.
37. Pacak K. Pheochromocytoma: a catecholamine and oxidative stress disorder. *Endocr Regul.* 2011;45(2):65-90.
38. Silva JE, Bianco SD. Thyroid-adrenergic interactions: physiological and clinical implications. *Thyroid.* 2008;18(2):157-65.
39. Jenkins ZM, Castle DJ, Eikelis N, Phillipou A, Lambert GW, Lambert EA. Autonomic nervous system function in women with anorexia nervosa. *Clin Auton Res.* 2021. doi: 10.1007/s10286-021-00836-z. Epub ahead of print.
40. Sharma VK, Singh TG. Chronic Stress and Diabetes Mellitus: Interwoven Pathologies. *Curr Diabetes Rev.* 2020;16(6):546-556.
41. van der Poll T. Effects of Catecholamines on the Inflammatory Response. *Sepsis.* 2001;4(2).
42. Thwaites CL, Yen LM, Cordon SM, et al. Urinary catecholamine excretion in tetanus. *Anaesthesia.* 2006;61(4):355-359.
43. Shimizu M, Kagawa A, Takano T, Masai H, Miwa Y. Neurogenic stunned myocardium associated with status epilepticus and postictal catecholamine surge. *Intern Med.* 2008;47(4):269-273.

44. Abraham J, Mudd JO, Kapur NK, Klein K, Champion HC, Wittstein IS. Stress cardiomyopathy after intravenous administration of catecholamines and beta-receptor agonists [published correction appears in *J Am Coll Cardiol*. 2009 May 12;53(19):1828. Kapur, Navin [corrected to Kapur, Navin K]]. *J Am Coll Cardiol*. 2009;53(15):1320-1325.
45. Ueyama T, Hano T, Kasamatsu K, Yamamoto K, Tsuruo Y, Nishio I. Estrogen attenuates the emotional stress-induced cardiac responses in the animal model of Tako-tsubo (Ampulla) cardiomyopathy. *J Cardiovasc Pharmacol*. 2003;42 Suppl 1:S117-S119.
46. Citro R, Okura H, Ghadri JR, et al. Multimodality imaging in takotsubo syndrome: a joint consensus document of the European Association of Cardiovascular Imaging (EACVI) and the Japanese Society of Echocardiography (JSE). *Eur Heart J Cardiovasc Imaging*. 2020;21(11):1184-1207.
47. Boland TA, Lee VH, Bleck TP. Stress-induced cardiomyopathy. *Crit Care Med*. 2015;43(3):686-693.
48. Cocco G, Chu D. Stress induced cardiomyopathy: A review. *Eur J Intern Med*. 2007;18(5):369-379.
49. Sharifi-Rad J, Rodrigues CF, Sharopov F, et al. Diet, Lifestyle and Cardiovascular Diseases: Linking Pathophysiology to Cardioprotective Effects of Natural Bioactive Compounds. *Int J Environ Res Public Health*. 2020;17(7):2326.
50. Versteeg H, Spek V, Pedersen SS, Denollet J. Type D personality and health status in cardiovascular disease populations: a meta-analysis of prospective studies. *Eur J Prev Cardiol*. 2012;19(6):1373-1380.
51. De Hert M, Detraux J, Vancampfort D. The intriguing relationship between coronary heart disease and mental disorders. *Dialogues Clin Neurosci*. 2018;20(1):31-40.

52. Broers ER, Gavidia G, Wetzels M, Ribas V, Ayoola I, Piera-Jimenez J, Widdershoven JWMG, Habibović M; Do CHANGE consortium. Usefulness of a Lifestyle Intervention in Patients With Cardiovascular Disease. *Am J Cardiol.* 2020;125(3):370-375.
53. Wang HX, Leineweber C, Kirkeeide R, et al. Psychosocial stress and atherosclerosis: family and work stress accelerate progression of coronary disease in women. The Stockholm Female Coronary Angiography Study. *J Intern Med.* 2007;261(3):245-254.
54. Gallo LC, Troxel WM, Kuller LH, Sutton-Tyrrell K, Edmondowicz D, Matthews KA. Marital status, marital quality, and atherosclerotic burden in postmenopausal women. *Psychosom Med.* 2003;65(6):952-962.
55. Joseph NT, Kamarck TW, Muldoon MF, Manuck SB. Daily marital interaction quality and carotid artery intima-medial thickness in healthy middle-aged adults. *Psychosom Med.* 2014;76(5):347-354.
56. Smith TW, Baron CE, Deits-Leuchner C, Uchino BN, Berg CA. Is it me or you? Marital conflict behavior and blood pressure reactivity. *J Fam Psychol.* 2020;34(4):503-508.
57. Carney RM, Freedland KE. Depression and coronary heart disease. *Nat Rev Cardiol.* 2017;14(3):145-155.
58. Edmondson D, von Känel R. Post-traumatic stress disorder and cardiovascular disease. *Lancet Psychiatry.* 2017;4(4):320-329.
59. Batelaan NM, Seldenrijk A, Bot M, van Balkom AJ, Penninx BW. Anxiety and new onset of cardiovascular disease: critical review and meta-analysis. *Br J Psychiatry.* 2016;208(3):223-231.

60. Surtees PG, Wainwright NW, Luben RN, Wareham NJ, Bingham SA, Khaw KT. Depression and ischemic heart disease mortality: evidence from the EPIC-Norfolk United Kingdom prospective cohort study. *Am J Psychiatry*. 2008;165(4):515-523.
61. Bradley SM, Rumsfeld JS. Depression and cardiovascular disease. *Trends Cardiovasc Med*. 2015;25(7):614-622.
62. Barton DA, Dawood T, Lambert EA, et al. Sympathetic activity in major depressive disorder: identifying those at increased cardiac risk? *J Hypertens*. 2007;25(10):2117-2124.
63. Stetler C, Miller GE. Depression and hypothalamic-pituitary-adrenal activation: a quantitative summary of four decades of research. *Psychosom Med*. 2011;73(2):114-126.
64. Delle Chiaie R, Capra E, Salviati M, Trabucchi G, Fancheri C, Corrado A, Biondi M. Persistence of subsyndromal residual symptoms after remission of major depression in patients without cardiovascular disease may condition maintenance of elevated platelet factor 4 and β -thromboglobulin plasma levels. *J Affect Disord*. 2013;150(2):664-7.
65. Chen H, Zhang L, Zhang M, et al. Relationship of depression, stress and endothelial function in stable angina patients. *Physiol Behav*. 2013;118:152-158.
66. Pizzi C, Manzoli I, Mancini S, Costa GM. Analysis of potential predictors of depression among coronary heart disease risk factors including heart rate variability, markers of inflammation, and endothelial function. *Eur Heart J*. 2008;29(9):1110-1117.
67. Remch M, Laskaris Z, Flory J, Mora-McLaughlin C, Morabia A. Post-Traumatic Stress Disorder and Cardiovascular Diseases: A Cohort Study of Men and Women Involved in Cleaning the Debris of the World Trade Center Complex. *Circ Cardiovasc Qual Outcomes*. 2018;11(7):e004572.

68. Tawakol A, Ishai A, Takx RA, et al. Relation between resting amygdalar activity and cardiovascular events: a longitudinal and cohort study [published correction appears in *Lancet*. 2017 Feb 25;389(10071):804] *Lancet*. 2017;389(10071):834-845.
69. Lakusic N, Fuckar K, Mahovic D, Cerovec D, Majsec M, Stancin N. Characteristics of Heart Rate Variability in War Veterans with Post-Traumatic Stress Disorder after Myocardial Infarction. *Military Medicine*. 2007;172(11).
70. Tardif JC. Heart rate as a treatable cardiovascular risk factor. *Br Med Bull*. 2009;90:71-84.
71. Morris MC, Compas BE, Garber J. Relations among posttraumatic stress disorder, comorbid major depression, and HPA function: a systematic review and meta-analysis. *Clin Psychol Rev*. 2012;32(4):301-315.
72. Bedi US, Arora R. Cardiovascular manifestations of posttraumatic stress disorder. *J Natl Med Assoc*. 2007;99(6):642-649.
73. Grenon SM, Owens CD, Alley C, et al. Posttraumatic Stress Disorder Is Associated with Worse Endothelial Function Among Veterans. *J Am Heart Assoc*. 2016;5(3):e003010.
74. Passos IC, Vasconcelos Moreno MP, Costa LG, et al. Inflammatory markers in post-traumatic stress disorder: a systematic review, meta-analysis, and meta-regression. *Lancet Psychiatry*. 2015;2(11):1002-1012.
75. Zen AL, Whooley MA, Zhao S, Cohen BE. Post-traumatic stress disorder is associated with poor health behaviors: findings from the heart and soul study. *Health Psychol*. 2012;31(2):194-201.

76. Kronish IM, Edmondson D, Li Y, Cohen BE. Post-traumatic stress disorder and medication adherence: results from the Mind Your Heart study. *J Psychiatr Res.* 2012;46(12):1595-1599.
77. Schulman JK, Stromberg S. On the value of doing nothing: anger and cardiovascular disease in clinical practice. *Cardiol Rev.* 2007;15(3):123-132.
78. Strike PC, Steptoe A. Behavioral and emotional triggers of acute coronary syndromes: a systematic review and critique. *Psychosom Med.* 2005;67(2):179-185.
79. Chida Y, Steptoe A. The association of anger and hostility with future coronary heart disease: a meta-analytic review of prospective evidence. *J Am Coll Cardiol.* 2009;53(11):936-946.
80. van der Meer RE, Maas AH. The Role of Mental Stress in Ischaemia with No Obstructive Coronary Artery Disease and Coronary Vasomotor Disorders. *Eur Cardiol.* 2021;16:e37. doi: 10.15420/ecr.2021.20.
81. al'Absi M, Bongard S, Levalle WR. Adrenocorticotropin responses to interpersonal stress: effects of overt anger expression style and defensiveness. *Int J Psychophysiol.* 2000;37(3):257-265.
82. Räikkönen K, Matthews KA, Sutton-Tyrrell K, Kuller LH. Trait anger and the metabolic syndrome predict progression of carotid atherosclerosis in healthy middle-aged women. *Psychosom Med.* 2004;66(6):903-908.
83. Scheier MF, Carver CS. Dispositional optimism and physical health: A long look back, a quick look forward. *Am Psychol.* 2018;73(9):1082-1094.

84. Kubzansky LD, Sparrow D, Vokonas P, Kawachi I. Is the glass half empty or half full? A prospective study of optimism and coronary heart disease in the normative aging study. *Psychosom Med.* 2001;63(6):910-916.
85. Pänkäläinen M, Kerola T, Kampman O, Kauppi M, Hintikka J. Pessimism and risk of death from coronary heart disease among middle-aged and older Finns: an eleven-year follow-up study. *BMC Public Health.* 2016;16(1):1124.
86. Nabi H, Koskenvuo M, Singh-Manoux A, et al. Low pessimism protects against stroke: the Health and Social Support (HeSSup) prospective cohort study. *Stroke.* 2010;41(1):187-190.
87. Boehm JK, Chen Y, Koga H, Mathur MB, Vittinghoff LL, Kubzansky LD. Is Optimism Associated With Healthier Cardiovascular-Related Behavior? Meta-Analyses of 3 Health Behaviors. *Circ Res.* 2018;122(8):1119-1134.
88. Whitfield JB, Zhu G, Landers JG, Martin NG. Pessimism is associated with greater all-cause and cardiovascular mortality, but optimism is not protective. *Sci Rep.* 2020;10(1):12609.
89. Ikeda A, Schwartz J, Peters JL, et al. Optimism in relation to inflammation and endothelial dysfunction in older men: the VA Normative Aging Study. *Psychosom Med.* 2011;73(8):664-671.
90. Zalli A, Carvalho LA, Lin J, et al. Shorter telomeres with high telomerase activity are associated with raised allostatic load and impoverished psychosocial resources. *Proc Natl Acad Sci U S A.* 2014;111(12):4519-4524.
91. Räikkönen K, Matthews KA. Do dispositional pessimism and optimism predict ambulatory blood pressure during school days and nights in adolescents? *J Pers.* 2008;76(3):605-30.

92. Endrighi R, Hamer M, Steptoe A. Associations of trait optimism with diurnal neuroendocrine activity, cortisol responses to mental stress, and subjective stress measures in healthy men and women. *Psychosom Med.* 2011;73(8):672-678.
93. Litchfield P, Cooper C, Hancock C, Watt P. Work and Wellbeing in the 21st Century †. *Int J Environ Res Public Health.* 2016;13(11):1065.
94. Haddon J. The impact of employees' well-being on performance in the workplace. *Strategic HR Review.* 2018;17(2).
95. Faghy MA, Duncan MJ, Pringle A, Meharry JB, Pascoe CMP. UK university staff experience high levels of sedentary behaviour during work and leisure time [published online ahead of print, 2021 Feb 25]. *Int J Occup Saf Ergon.* 2021;1-8.
96. Kivimäki M, Kawachi I. Work Stress as a Risk Factor for Cardiovascular Disease. *Curr Cardiol Rep.* 2015;17(9):630.
97. Steptoe A, Kivimäki M. Stress and cardiovascular disease: an update on current knowledge. *Annu Rev Public Health.* 2013;34:337-354.
98. Kim ES, Delaney SW, Kubzansky LD. Sense of Purpose in Life and Cardiovascular Disease: Underlying Mechanisms and Future Directions. *Curr Cardiol Rep.* 2019;21(11):135.
99. Cohen R, Bavisani C, Rozanski A. Purpose in Life and Its Relationship to All-Cause Mortality and Cardiovascular Events: A Meta-Analysis. *Psychosom Med.* 2016;78(2):122-133.
100. Kubzansky LD, Huffman JC, Boehm JK, et al. Positive Psychological Well-Being and Cardiovascular Disease: JACC Health Promotion Series. *J Am Coll Cardiol.* 2018;72(12):1382-1396.
101. Brent DA, Silverstein M. Shedding light on the long shadow of childhood adversity. *JAMA.* 2013;309(17):1777-1778.

102. Murphy MO, Cohn DM, Loria AS. Developmental origins of cardiovascular disease: Impact of early life stress in humans and rodents. *Neurosci Biobehav Rev.* 2017;74(Pt B):453-465.
103. Benjamin EJ, Muntner P, Alonso A, et al. Heart Disease and Stroke Statistics-2019 Update: A Report From the American Heart Association [published correction appears in *Circulation.* 2020;141(2):e33]. *Circulation.* 2019;139(10):e56-e528.
104. Vlachopoulos C, Xaplanteris P, Alexopoulos N, et al. Divergent effects of laughter and mental stress on arterial stiffness and central hemodynamics. *Psychosom Med.* 2009;71(4):446-453.
105. Song H, Fang F, Arnberg FK, et al. Stress related disorders and risk of cardiovascular disease: population based, sibling controlled cohort study. *BMJ.* 2019;365:11255.
106. Yusuf S, Hawken S, Ounpuu S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet.* 2004;364(9438):937-952.
107. Juonala M, Pulkki-Råback L, Elovainio M, et al. Childhood Psychosocial Factors and Coronary Artery Calcification in Adulthood: The Cardiovascular Risk in Young Finns Study. *JAMA Pediatr.* 2016;170(5):466-472.
108. Virtanen M, Heikkila K, Jokela M, et al. Long Working Hours and Coronary Heart Disease: A Systematic Review and Meta-Analysis. *American Journal of Epidemiology.* 2012;176(7).
109. Schnohr P, Marott JL, Kristensen TS, Gyntelberg F, Grønbaek M, Lange P, Jensen MT, Jensen GB, Prescott E. Ranking of psychosocial and traditional risk factors by importance for coronary heart disease: the Copenhagen City Heart Study. *Eur Heart J.* 2015;36(22):1385-93.

110. Slopen N, Glynn RJ, Buring JE, Lewis TT, Williams DR, Albert MA. Job strain, job insecurity, and incident cardiovascular disease in the Women's Health Study: results from a 10-year prospective study. *PLoS One*. 2012;7(7):e40512. doi: 10.1371/journal.pone.0040512. Epub 2012 Jul 18.
111. Esler M. Mental stress and human cardiovascular disease. *Neurosci Biobehav Rev*. 2017;74(Pt B):269-276.
112. World Health Organization. Prevention of Cardiovascular Disease: Guidelines for Assessment and Management of Cardiovascular Risk.; 2007. Available at: DocHdl1tmpTarget (who.int)
113. Piepoli MF, Hoes AW, Agewall S, et al. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J*. 2016;37(29):2315-2381.
114. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines [published correction appears in *J Am Coll Cardiol*. 2019;74(10):1429-1430] [published correction appears in *J Am Coll Cardiol*. 2020;75(7):840]. *J Am Coll Cardiol*. 2019;74(10):e177-e232.
115. Cowie A, Buckley J, Doherty P, et al. Standards and core components for cardiovascular disease prevention and rehabilitation. *Heart*. 2019;105(7):510-515.

116. Balady GJ, Williams MA, Ades PA, et al. Core components of cardiac rehabilitation/secondary prevention programs: 2007 update: a scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *J Cardiopulm Rehabil Prev.* 2007;27(3):121-129.
117. Grace SL, Turk-Adawi KI, Contractor A, et al. Cardiac Rehabilitation Delivery Model for Low-Resource Settings: An International Council of Cardiovascular Prevention and Rehabilitation Consensus Statement. *Prog Cardiovasc Dis.* 2016;59(3):303-322.
118. Supervia M, Turk-Adawi K, Lopez-Jimenez F, et al. Nature of Cardiac Rehabilitation Around the Globe. *E Clinical Medicine.* 2019;13:46-56.
119. Turk-Adawi K, Supervia M, Lopez-Jimenez F, et al. Cardiac Rehabilitation Availability and Density around the Globe. *E Clinical Medicine.* 2019;13:31-45.
120. Chen X, Velez JC, Barbosa C, et al. Smoking and perceived stress in relation to short salivary telomere length among caregivers of children with disabilities. *Stress.* 2015;18(1):20-28.
121. Terrill AL, Gjerde JM, Carofalo JP. Background Stress Inventory: Developing a Measure of Understudied Stress. *Stress Health.* 2015;31(4):290-8.
122. Lee EH. Review of the psychometric evidence of the perceived stress scale. *Asian Nurs Res (Korean Soc Nurs Sci).* 2012;6(4):121-127.
123. Redmond N, Richman J, Gamboa CM, et al. Perceived stress is associated with incident coronary heart disease and all-cause mortality in low- but not high-income participants in the Reasons for Geographic And Racial Differences in Stroke study. *J Am Heart Assoc.* 2013;2(6):e000447. Published 2013 Dec 19.

124. O'Reilly GA, Cook L, Spruijt-Metz D, Black DS. Mindfulness-based interventions for obesity-related eating behaviours: a literature review. *Obes Rev.* 2014;15(6):453-461.
125. Gu J, Strauss C, Bond R, Cavanagh K. How do mindfulness-based cognitive therapy and mindfulness-based stress reduction improve mental health and wellbeing? A systematic review and meta-analysis of mediation studies [published correction appears in *Clin Psychol Rev.* 2016;49:119]. *Clin Psychol Rev.* 2015;37:1-12.
126. Tsai JC, Wang WH, Chan P, et al. The beneficial effects of Tai Chi Chuan on blood pressure and lipid profile and anxiety status in a randomized controlled trial. *J Altern Complement Med.* 2003;9(5):747-754.
127. Lim SA, Cheong KJ. Regular Yoga Practice Improves Antioxidant Status, Immune Function, and Stress Hormone Releases in Young Healthy People: A Randomized, Double-Blind, Controlled Pilot Study. *J Altern Complement Med.* 2015;21(9):530-538.
128. Creswell JD, Pacilio LE, Lindsay EK, Brown KW. Brief mindfulness meditation training alters psychological and neuroendocrine responses to social evaluative stress. *Psychoneuroendocrinology.* 2014;44:1-12.
129. Imboden C, Gerber M, Beck J, Holsboer-Trachsler E, Pühse U, Hatzinger M. Aerobic exercise or stretching as add-on to inpatient treatment of depression: Similar antidepressant effects on depressive symptoms and larger effects on working memory for aerobic exercise alone. *J Affect Disord.* 2020;276:866-876.
130. Edwards MK, Loprinzi PD. Affective Responses to Acute Bouts of Aerobic Exercise, Mindfulness Meditation, and Combinations of Exercise and Meditation: A Randomized Controlled Intervention. *Psychol Rep.* 2019;122(2):465-484.

131. Wippert PM, Rector M, Kuhn G, Wuertz-Kozak K. Stress and Alterations in Bones: An Interdisciplinary Perspective. *Front Endocrinol (Lausanne)*. 2017;8:96.
132. McEwen BS. Physiology and neurobiology of stress and adaptation: central role of the brain. *Physiol Rev*. 2007 Jul;87(3):873-904. doi: 10.1152/physrev.00041.2006. PMID: 17615391.
133. Dhabhar FS. The short-term stress response - Mother nature's mechanism for enhancing protection and performance under conditions of threat, challenge, and opportunity. *Front Neuroendocrinol*. 2018;49:175-192.
134. Chrousos GP. Stress and disorders of the stress system. *Nat Rev Endocrinol*. 2009;5(7):374-381.
135. Turner AI, Smyth N, Hall SJ, et al. Psychological stress reactivity and future health and disease outcomes: A systematic review of prospective evidence. *Psychoneuroendocrinology*. 2020;114:104599.
136. Brydon L, Steptoe A. Stress-induced increases in interleukin-6 and fibrinogen predict ambulatory blood pressure at 3-year follow-up. *J Hypertens*. 2005;23(5):1001-1007.
137. Matthews KA, Katholi CR, McCreath H, et al. Blood pressure reactivity to psychological stress predicts hypertension in the CARDIA study. *Circulation*. 2004;110(1):74-78.
138. Jennings JR, Kamarck TW, Everson-Rose SA, Kaplan GA, Manuck SB, Salonen JT. Exaggerated blood pressure responses during mental stress are prospectively related to enhanced carotid atherosclerosis in middle-aged Finnish men. *Circulation*. 2004;110(15):2198-2203.
139. Phillips AC, Hunt K, Der G, Carroll D. Blunted cardiac reactions to acute psychological stress predict symptoms of depression five years later: evidence from a large community study. *Psychophysiology*. 2011;48(1):142-148.

140. Hamer M, Endrighi R, Venuraju SM, Lahiri A, Steptoe A. Cortisol responses to mental stress and the progression of coronary artery calcification in healthy men and women. *PLoS One*. 2012;7(2):e31356.
141. Steudte-Schmiedgen S, Stalder T, Schönfeld S, et al. Hair cortisol concentrations and cortisol stress reactivity predict PTSD symptom increase after trauma exposure during military deployment. *Psychoneuroendocrinology*. 2015;59:123-133.
142. Dhabhar FS, Mcewen BC. Bi-directional effects of stress on immune function: Possible explanations for salubrious as well as harmful effects. *Psychoneuroimmunology*, two-volume set edn. Elsevier Inc. 2007:723-760.
143. Zefferino R, Di Gioia S, Conese M. Molecular links between endocrine, nervous and immune system during chronic stress. *Brain Behav*. 2021;11(2):e01960. Epub 2020 Dec 8.
144. Saul AN, Oberyszyn TM, Daugherty C, et al. Chronic stress and susceptibility to skin cancer. *J Natl Cancer Inst*. 2005;97(23):1760-1767.
145. van Praag HM. Can stress cause depression? *Prog Neuropsychopharmacol Biol Psychiatry*. 2004;28(5):891-907.
146. Cusi AM, Nazarov A, Holshausen K, Macqueen GM, McKinnon MC. Systematic review of the neural basis of social cognition in patients with mood disorders. *J Psychiatry Neurosci*. 2012;37(3):154-169.
147. Mazurka R, Wynne-Edwards KE, Harkness KL. Sex Differences in the Cortisol Response to the Trier Social Stress Test in Depressed and Nondepressed Adolescents. *Clinical Psychological Science*. 2018;6(3).
148. Mastorakos G, Pavlatou M, Diamanti-Kandarakis E, Chrousos GP. Exercise and the stress system. *Hormones (Athens)*. 2005;4(2):73-89.

149. Misra M. Neuroendocrine mechanisms in athletes. *Handb Clin Neurol*. 2014;124:373-386.
150. Hackney AC, Lane AR. Exercise and the Regulation of Endocrine Hormones. *Prog Mol Biol Transl Sci*. 2015;135:293-311.
151. Ensari I, Schwartz JE, Edmondson D, Duran AT, Shimbo D, Diaz KM. Testing the cross-stressor hypothesis under real-world conditions: exercise as a moderator of the association between momentary anxiety and cardiovascular responses. *J Behav Med*. 2020;43(6):989-1001.
152. Hamer M, Taylor A, Steptoe A. The effect of acute aerobic exercise on stress related blood pressure responses: a systematic review and meta-analysis. *Biol Psychol*. 2006;71(2):183-190.
153. Zschucke E, Renneberg B, Dimeo F, Westenberg T, Ströhle A. The stress-buffering effect of acute exercise: Evidence for HPA axis negative feedback. *Psychoneuroendocrinology*. 2015;51:414-425.
154. Physical Activity Guidelines Advisory Committee. 2018 Physical Activity Guidelines Advisory Committee Scientific Report. Washington DC; 2018. Available at: 2018 Physical Activity Guidelines Advisory Committee Scientific Report - Healthy People 2030 | health.gov
155. Gordon BR, McDowell CP, Lyons M, Herring MP. The Effects of Resistance Exercise Training on Anxiety: A Meta-Analysis and Meta-Regression Analysis of Randomized Controlled Trials. *Sports Med*. 2017;47(12):2521-2532.
156. Stubbs B, Vancampfort D, Rosenbaum S, et al. An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: A meta-analysis. *Psychiatry Res*. 2017;249:102-108.

157. Rosenbaum S, Tiedemann A, Sherrington C, Curtis J, Ward PB. Physical activity interventions for people with mental illness: a systematic review and meta-analysis. *J Clin Psychiatry*. 2014;75(9):964-974.
158. Powers MB, Medina JL, Burns S, et al. Exercise Augmentation of Exposure Therapy for PTSD: Rationale and Pilot Efficacy Data. *Cogn Behav Ther*. 2015;44(4):314-327.
159. Gaudlitz K, Plag J, Dimeo F, Ströhle A. Aerobic exercise training facilitates the effectiveness of cognitive behavioral therapy in panic disorder. *Depress Anxiety*. 2015;32(3):221-228.
160. Kandola A, Ashdown-Franks G, Stubbs B, Osborn DPJ, Hayes JF. The association between cardiorespiratory fitness and the incidence of common mental health disorders: A systematic review and meta-analysis. *J Affect Disord*. 2019;257:748-757.
161. Ross R, de Lannoy L, Stotz PJ. Separate Effects of Intensity and Amount of Exercise on Interindividual Cardiorespiratory Fitness Response. *Mayo Clin Proc*. 2015;90(11):1506-1514.
162. LeBouthillier DM, Asmundson GJG. The efficacy of aerobic exercise and resistance training as transdiagnostic interventions for anxiety-related disorders and constructs: A randomized controlled trial. *J Anxiety Disord*. 2017;52:43-52.
163. Phillips C. Brain-Derived Neurotrophic Factor, Depression, and Physical Activity: Making the Neuroplastic Connection. *Neural Plast*. 2017;2017:7260130. doi: 10.1155/2017/7260130. Epub 2017 Aug 8.
164. Ströhle A, Stoy M, Graetz B, et al. Acute exercise ameliorates reduced brain-derived neurotrophic factor in patients with panic disorder. *Psychoneuroendocrinology*. 2010;35(3):364-368.

165. Kandola A, Vancampfort D, Herring M, Rebar A, Hallgren M, Firth J, Stubbs B. Moving to Beat Anxiety: Epidemiology and Therapeutic Issues with Physical Activity for Anxiety. *Curr Psychiatry Rep.* 2018;20(8):63.
166. Lindheimer JB, O'Connor PJ, Dishman RK. Quantifying the placebo effect in psychological outcomes of exercise training: a meta-analysis of randomized trials. *Sports Med.* 2015;45(5):693-711.
167. Ensari I, Greenlee TA, Motl RW, Petruzzello SJ. META-ANALYSIS OF ACUTE EXERCISE EFFECTS ON STATE ANXIETY: AN UPDATE OF RANDOMIZED CONTROLLED TRIALS OVER THE PAST 25 YEARS. *Depress Anxiety.* 2015;32(8):624-634.
168. Gleeson M. Immune function in sport and exercise. *J Appl Physiol (1985).* 2007;103(2):693-699.
169. Walsh NP, Whitham M. Exercising in environmental extremes : a greater threat to immune function?. *Sports Med.* 2006;36(11):941-976.
170. Katz DL. The Mass of Humanity and the Weight of the World: Obesity and the Environment at a Confluence of Causes. *Curr Obes Rep.* 2016;5(4):386-388.
171. Owen N, Healy GN, Matthews CE, Dunstan DW. Too much sitting: the population health science of sedentary behavior. *Exerc Sport Sci Rev.* 2010;38(3):105-113.
172. O'Keefe EL, O'Keefe JH, Lavie CJ. Exercise Counteracts the Cardiotoxicity of Psychosocial Stress. *Mayo Clin Proc.* 2019;94(9):1852-1864.
173. Lavie CJ, Menezes AR, De Schutter A, Milani RV, Blumenthal JA. Impact of Cardiac Rehabilitation and Exercise Training on Psychological Risk Factors and Subsequent Prognosis in Patients With Cardiovascular Disease. *Can J Cardiol.* 2016;32(10 Suppl 2):S365-S373.

174. Lavie CJ, Milani RV. Adverse psychological and coronary risk profiles in young patients with coronary artery disease and benefits of formal cardiac rehabilitation. *Arch Intern Med*. 2006;166(17):1878-1883.
175. Milani RV, Lavie CJ. Impact of cardiac rehabilitation on depression and its associated mortality. *Am J Med*. 2007;120(9):799-806.
176. Milani RV, Lavie CJ. Reducing psychosocial stress: a novel mechanism of improving survival from exercise training. *Am J Med*. 2009;122(10):931-938.
177. Milani RV, Lavie CJ, Mehra MR, Ventura HO. Impact of exercise training and depression on survival in heart failure due to coronary heart disease. *Am J Cardiol*. 2011;107(1):64-68.
178. Blumenthal JA, Sherwood A, Smith PJ, et al. Enhancing Cardiac Rehabilitation with Stress Management Training: A Randomized, Clinical Efficacy Trial. *Circulation*. 2016;133(14):1341-1350.
179. Kachur S, Menezes AR, De Schutter A, Milani RV, Lavie CJ. Significance of Comorbid Psychological Stress and Depression on Outcomes After Cardiac Rehabilitation. *Am J Med*. 2016;129(12):1316-1321.
180. Arena R, Lavie CJ; HL-PIVOT Network. The global path forward - Healthy Living for Pandemic Event Protection (HL - PIVOT). *Prog Cardiovasc Dis*. 2021;64:96-101.
181. Hall G, Laddu DR, Phillips SA, Lavie CJ, Arena R. A tale of two pandemics: How will COVID-19 and global trends in physical inactivity and sedentary behavior affect one another? *Prog Cardiovasc Dis*. 2021;64:108-110.
182. Sanchis-Gomar F, Lavie CJ, Mehra MR, Henry BM, Lippi G. Obesity and Outcomes in COVID-19: When an Epidemic and Pandemic Collide. *Mayo Clin Proc*. 2020;95(7):1445-1453.

183. Dubey S, Biswas P, Ghosh R, Chatterjee S, Dubey MJ, Chatterjee S, Lahiri D, Lavie CJ. Psychosocial impact of COVID-19. *Diabetes Metab Syndr*. 2020;14(5):779-788.
184. Sallis R, Young DR, Tartof SY, Sallis JF, Sall J, Li Q, Smith GN, Cohen DA. Physical inactivity is associated with a higher risk for severe COVID-19 outcomes: a study in 48 440 adult patients. *Br J Sports Med*. 2021;55(19):1099-1105.
185. Brawner CA, Ehrman JK, Bole S, Kerrigan DJ, Parikh SS, Lewis BK, Gindi RM, Keteyian C, Abdul-Nour K, Keteyian SJ. Inverse Relationship of Maximal Exercise Capacity to Hospitalization Secondary to Coronavirus Disease 2019. *Mayo Clin Proc*. 2021;96(1):32-39.
186. Lavie CJ, Sanchis-Gomar F, Arena R. Fit Is It in COVID-19, Future Pandemics, and Overall Healthy Living. *Mayo Clin Proc*. 2021;96(1):7-9.
187. Lavie CJ, Gerber TC, Lanier WL. Hurricane Katrina: the infarcts beyond the storm. *Disaster Med Public Health Prep*. 2009;3(3):131-5.

Conflict of interest: None.