

What workers can tell us about post-COVID workability

Jennifer Lunt,^a Sally Hemming,^b Kim Burton,^c James Elander,^a and Amy Baraniak^a

^a University of Derby; ^b University of Loughborough; ^c University of Huddersfield

Correspondence: Jennifer Lunt j.lunt@derby.ac.uk

Cite as:

Lunt, J., Hemming, S., Burton, K., Elander, J., & Baraniak, A. (2022). Post-COVID work ability: what can workers tell us? *Occupational Medicine*, kqac086.

<https://doi.org/10.1093/occmed/kqac086>

Abstract

Background: The apparent functional impact of post-COVID-19 syndrome has work ability implications for large segments of the working age population.

Aims: To understand obstacles and enablers around self-reported work ability of workers following COVID-19, to better guide sustainable workplace accommodations.

Methods: An exploratory online survey comprising quantitative and qualitative questions was disseminated via social media and industry networks between December 2020 and February 2021, yielding usable responses from 145 workers. Qualitative data was subjected to content analysis.

Results: Over half of the sample (64%) were from health, social care, and education sectors. Just under 15% had returned to work, and 53% and 50% reported their physical and psychological work ability respectively as moderate at best. Leading work ability obstacles were multi-level, comprising fatigue, the interaction between symptoms and job, lack of control over job pressures, inappropriate sickness absence management policies, and lack of COVID-aware organisational cultures. Self-management support, modified work, flexible co-developed graded return-to-work planning, and improved line management competency were advocated as key enablers.

Conclusions: Assuming appropriate medical management of any pathophysiological complications of COVID-19, maintaining or regaining post-COVID work ability might reasonably follow a typical biopsychosocial framework enhanced to cater for the fluctuating nature of the symptoms. This should entail flexible, regularly reviewed and longer-term return-to-work planning addressing multi-level work ability obstacles, co-developed between workers and line managers, with support from human resources, occupational health professionals, and a COVID-aware organisational culture.

Key words: post-COVID-19 symptoms, work ability, long COVID, biopsychosocial, return-to-work, vocational rehabilitation, workplace accommodations.

Key learning points

What is already known about this subject?

- The prevalence of post-COVID-19 symptoms is greatest amongst the working age population relative to other age cohorts.
- Their impact upon physical, cognitive and psychological functioning implies detrimental effects on sustained work ability.
- Vocational rehabilitation approaches based on the biopsychosocial framework can address multi-level obstacles to working with long-term health conditions.

What does this study add?

- The self-reported insights of workers recovering from COVID-19 have identified commonly experienced post-COVID work ability obstacles. Their fit within the biopsychosocial framework implies cross-organisation and sector applicability.
- Workers' perspectives on workplace accommodations for overcoming post-COVID work ability obstacles are unpacked, for use by practitioners, employers and employees, including variation from typical vocational rehabilitation practices.
- Workers' perspectives include perceived benefits that employers could gain from accommodating workers' post-COVID recovery.

What impact this may have on practice and policy?

- Variations from typical vocational rehabilitation appears to be a matter of emphasis. Longer-term, flexible, co-developed and regularly reviewed return to work plans appear to be particularly necessary for accommodating the unpredictable nature of post-COVID recovery and other conditions sharing similar unpredictable symptom characteristics, including chronic fatigue syndrome.
- A biopsychosocial approach provides an appropriate framework for identifying and overcoming work-relevant post-COVID work ability obstacles for use by employers, employees and Occupational Health Practitioners.
- Case studies of successful vocational rehabilitation for workers experiencing post-COVID -19 syndrome could help shape a narrative that early sustained RTW is possible where suitable workplace accommodations are agreed.

Introduction

Just over 1 million people self-reported as having 'long COVID' symptoms within the United Kingdom at the start of September 2021 [1]. Prevalence was highest amongst 35 to 69-year-olds, corresponding to a broad spectrum of the working age population. Long COVID refers to symptoms that continue or develop after acute COVID-19, this includes both ongoing symptomatic COVID-19 (from 4 to 12 weeks) and post-COVID-19 syndrome (symptoms beyond 12 weeks) [2,3]. A survey of 3,300 workers self-reporting as having the syndrome found 90% experienced fatigue [4]. Other leading symptoms included diminished cognitive capacity ('brain-fog'), shortness of breath, pain and muscle-ache. Symptom clusters can be disproportionate to those experienced in the acute infection phase [2]. The enduring, fluctuating multi-system nature of symptoms has obvious implications for work ability (WA) [5] and for vocational rehabilitation (VR). For example, of 138 health care workers reporting symptoms, 32% described themselves as struggling to cope up to 4 months post infection [6].

To address the problem, the NHS introduced a number of long COVID clinics for multidisciplinary assessment and rehabilitation [7]. However, service access does not appear universal [8]: there are still relatively few clinics and waiting lists are reported as long [9]. It seems pertinent to ask whether the limited resources could be supported by existing VR approaches. Current good practice in VR generally recognises that 'good' work is good for health [10, 12, 13]; requires a person-centred approach, line manager input and early intervention [11]. Recently developed return-to-work (RTW) guidance on workers recovering from COVID-19 for use by health care professionals [14] and workers [15] seems to follow this approach, advocating: regular contact with affected workers; assessment and regular review of work-relevant health needs; joint identification by manager and employer of obstacles hindering work resumption; reasonable temporary workplace accommodations by which obstacles can be overcome [10, 16,17] and documented within risk assessments, fit notes or RTW plans. Reflecting the biopsychosocial health perspective, obstacles can be health/symptom related, psychological, occupational, or social/contextual [17], while workplace accommodations (or adjustments) can encompass phased return and working pattern, workload, or job responsibility/task adjustments [17, 18].

The relative recency of long COVID means that such guidance drew on established VR principles rather than direct evidence derived from work-relevant experiences of workers recovering from COVID-19. To address this gap, this paper reports findings from an online survey to quantitatively establish the WA status of workers recovering from COVID-19, together with qualitative exploration of their work-relevant recovery experiences, views on workplace accommodations necessary for sustained RTW/WA and benefits for employers in making accommodations. The findings could clarify whether current RTW guidance for occupational health professionals for long-COVID is sufficiently fit-for-purpose.

Methods

An exploratory online cross-sectional survey comprising a mixture of quantitative and qualitative open-ended items was developed using Qualtrics^{XM} [19], piloted for usability and disseminated online between mid-December 2020 and February 2021. Ethical approval for the survey was gained from the ethics committee of the University of Derby's College of Health, Psychology and Social Care. Participants were recruited via weekly social media posts to COVID-19 and long COVID support groups and opportunistically via research team contacts with U.K based online construction industry, occupational health, academic, professional, carer and organisational networks. To allow for the limited COVID-19 testing towards pandemic outset, UK workers who had either tested positive for COVID-19 or suspected they had were considered eligible. A total of 145 eligible responses were received.

The survey was created using Qualtrics^{XM} (Qualtrics, 2021). Data was exported to IBM SPSS Statistics for Windows, version 26 for quantitative analysis (IBM, 2021). Survey items encompassed demographics, health status, RTW/WA status and views of: RTW obstacles; enablers and benefits to employers for enabling RTW (see Table 1 for structural summary).

INSERT TABLE 1

For health status (see Table 2 for summary statistics), the presence of pre-existing mental or physical medically diagnosed health conditions was assessed by asking participants to select from a list of generic condition labels [20]. Prevalence of post-viral symptoms was determined by asking participants to rate what proportion of a list of post-viral symptoms [21] they had experienced. The list was refined through team consensus. Work ability was assessed using the two single-item Work ability Assessment Inventory 2 (WAI2) scale selected due to its standardisation upon generic working populations, construct validity and brevity [5]. Views of anticipated or actual RTW obstacles and enablers were sought via open-ended items. Views about the benefits to employers for accommodating COVID-19 were sought to help create a RTW business case.

A content analysis [22] was conducted to identify the frequency of meaningful 'categories' of RTW qualitative data from the open-ended items, as indication of their relative priority or importance to workers. Categories refers to groups of words with similar meaning of connotations [23]. The procedure modelled Bowling's [24]. Two researchers independently re-read open-ended responses captured within Excel spreadsheets, discussed emerging categories, and agreed labels. For RTW obstacles and enablers, categories were separated according to individual (physical and psychological), job/work support, organisational and external groupings as per the biopsychosocial VR model [11]. For each open-ended question, online coding templates were created for documenting categories, labels, example quotes, and frequencies by which they arose. The first researcher then coded all responses for each item, counting category frequency. The second researcher then conducted inter-rater checks on all coding. Disagreements were resolved by the first

researcher checking the second researcher's coding decisions, accepting or discussing and resolving areas of disagreement. Categories receiving more than 10 counts were included in this analysis.

INSERT TABLE 2

Results

Of the 145 usable responses, 88% were female participants, 70% self-reported as key workers and 70% reported occupying predominantly non-managerial roles. Ages ranged from 25 to 65 years. The most frequently represented sectors comprised health and social care (50%), educational (15%) and professional, scientific and technical (10%). Of those that indicated their role, 25 (17%) were nurses, 22 (15%) were medics, 14 (10%) were from the allied health professions, 17 (12%) were teachers, and 9 (6%) were social workers or support workers.

In terms of health status, over half (52%) of participants reported pre-existing mental or physical health conditions. Most (see Table 2) reported having contracted COVID-19 more than 6 months previously with their symptoms continuing for longer than 6 months. Nearly 35% said their initial symptoms were mild to moderate, the remainder reporting severe symptoms (at home, 44%, or hospitalised, 12%). Most (91%) self-reported as having one or more of the listed post-viral symptoms.

In terms of work status, just under 15% had fully returned to work, and 16% had partially returned. From the open-ended responses, 9 (6%) of participants depicted the RTW process as "*straightforward*". Twenty-nine (20%) portrayed it as "*difficult*," and a further 17 (12%) reported "*multiple attempts*", with 10 (7%) claiming returning to work to specifically trigger relapse: "*I have made 3 attempts to come back to work and relapsed every time*".

Just 8% of the full sample rated their physical WA to work as good or very good, and 10 % described their mental WA to work as good or very good. As reported elsewhere [25] significant relationships were found between WA and COVID-19 duration.

For RTW obstacles, enablers, and employer benefits derived from qualitative data are listed within Tables 3 to 5, with category frequency and supporting quotes indicated. Category labels from these tables are italicised within the following analysis.

INSERT TABLE 3

Fatigue and *poor concentration* (see Table 3) represented symptoms most frequently portrayed as obstacles at the individual level. The *relapsing nature of symptoms* was also widely attributed as hampering "*return-to-work planning*". Psychological-related obstacles comprised concerns over maintaining: *social distancing* (avoiding reinfection); *safe practice*

(with implications for personal as well as patient safety in caring roles), and *professional identity*.

At the job level, an *interaction between symptoms and physical demands* in terms of physical load, including “*heavy lifting*,” or duration of physical activity including “*being on your feet all day*,” posed key obstacles. Similarly, the *interaction between symptoms and cognitive demands* was ascribed as a leading obstacle. This applied to having to “*concentrate*,” “*word find*” and to meta-cognitive tasks including having to “*multitask*,” engage in “*strategic thinking*,” “*chair meetings*,” “*teach*” or hold sustained “*conversations*.” Inadequate *control over job pressures* due to, “*meeting deadlines*” featured strongly amongst the obstacles. Difficulties upholding usual *working patterns*, especially where “*early starts*” and “*long days*” were involved, also represented leading obstacles. These job-related obstacles reflect situations where people are struggling with their usual job requirements in the face of work-relevant symptoms.

For work support, leading reported obstacles related to *line management*, *peers/colleague behaviour*, *occupational health (OH)* and *human resources (HR)*. Line manager behaviour-related obstacles included: inadequate reactions including “*not conducting risk assessments*,” failing to “*believe what you are saying*,” inadequate understanding over “*what long-COVID meant*,” failing to implement “*OH recommendations*,” and placing pressure on worker recovery on the premise that an employee should “*hit it the ground running on return*.” Other obstacles included being “*inaccessible*” or threatening job loss: “*you risk losing your job if you phone in off-sick*”. Covering for off-sick colleagues for protracted periods was described as contributing to “*uncertainty*,” “*burnout*” and “*resentment*” amongst peers due to being in “*harms-way*” more often. Negative attributes of OH and HR support comprised: being “*physically unavailable*” in the case of OH; being “*slow*” and constraining sick pay in the case of HR, potentially compounded by unawareness of national absence guidance for COVID-19: “*My line manager and HR weren’t aware of the national guidance regarding COVID absence*.”

At the organisational level, implementation of *sickness absence policy* was portrayed as “*quite rigid*,” providing only “*limited sick-pay within the first few years*,” catering for short-term as opposed to “*long-term illness*” and leading to COVID-19 omission from absence reporting systems: “*if Covid is not named on fit note it does not trigger absence management*”.

Fears over *job security* also emerged as an organisational wide obstacle, potentially compounding “*the stress of recovery*.” Apparent widespread attitudes “*that you must be healthy to be successful*” and that being back at work means “*you are back fully or not at all*” were implied to make it “*harder to recover properly*.” Inadequate knowledge of the nature of recovery was also frequently cited (see Table 3). Inadequate knowledge and shared attitudes can be regarded as reflective of *organisational culture*.

A commonly encountered external obstacle concerned *access to suitable health care*, attributed to either: inadequate understanding “*from professionals such as GPs of long-COVID;*” difficulties in obtaining “*GP appointments;*” and “*no access to long-COVID clinics.*” Transport obstacles encompassed the effects of a “*commute upon fatigue*”, using “*public transport*”, or “*walking from car-parks to a place of work*”.

INSERT TABLE 4

Leading RTW enablers are listed in Table 4. *Self-management* was widely supported, yet aspects, including *pacing* were considered impractical for some jobs: “*you can't take a break when needed, and you can't even sit down most of the time.*” *Flexible working* was also widely considered as enabling “*work from home*”, “*rest facilities at work*” and “*flexible attitudes to working time.*” While *graded return-to-work* featured strongly, participants having undergone phased return cautioned that it could take longer than “*4 weeks,*” and should omit any large steps:

“*I was expected to go from a few weeks of reduced hours ... to full time and full duties. This was not graded return, and still being ill, I found this impossible to manage.*”

Changes to jobs and/or tasks were also extensively supported. In terms of their duration, 46 (32%) participants indicated that adjustments might need to be *long-term*:

“*It takes as long as it takes, which may be permanent if we are permanently disabled.*”

In terms of support, various suggestions for improving *line manager competencies* were made. “*Regular catch-ups,*” “*face-to-face meetings*” with a “*single point of contact*” were suggested as enabling. Close communication between HR and line managers “*according to a shared return-to-work plan*” was also proposed for enabling joined-up support. *Managing peer expectations*, improving *occupational health and health care access and utility*, creating more COVID-centric *sickness absence policies*, targeting organisational wide *awareness* levels and attitudes with respect to COVID-19 were widely supported, so that “*illness is not viewed as an inconvenience or stigma*”. Working from home was attributed by 31 (21%) participants as supporting their WA by providing “*control*” over “*when to work and what to focus on.*”

INSERT TABLE 5

Table 5 details the themes participants identified as benefits to employers accruing from making workplace accommodations, along with supporting quotes. In order of frequency, ability to *retain specialist skills*, *fostering commitment*, enabling *sustained return-to-work* and *productivity* were cited.

Discussion

This exploratory investigation of the implications of COVID-19 on WA has identified key obstacles to resuming former WA levels, and relevant workplace accommodations. Findings are based on the actual or anticipated experiences of workers who believed they were recovering from SARS-CoV-2 infection, the majority of whom appeared to have post-COVID-19 syndrome to varying degrees. A small minority had fully returned to work. The majority self-reported their physical and psychological WA as moderate at best.

The obstacles to RTW most frequently highlighted (>25 participants) spanned multiple domains, comprising: fatigue; the interaction between symptoms and physical job demands; inadequate control over job pressures; inappropriate sickness absence management policies; and lack of COVID-aware organisational cultures. Highlighting the most commonly described obstacles should not obscure the significance of others, due to their potential interaction including between the physical and cognitive demands of a role.

Those most commonly described RTW enablers (<25 participants) comprised: self-management of symptoms alongside workplace demands; graded RTW planning where viable; modified job tasks or responsibilities, and improved line-management competency. Since these are participant-generated, they are not necessarily exhaustive of all potential accommodations.

INSERT TABLE 6

A summary of data-derived potential workplace accommodations that employers can make for workers with post-COVID-19 syndrome, are provided in Table 6. In return for making these accommodations, the findings suggest that employers will benefit according to: the retention of specialist skills; worker commitment; productivity, and sustained WA. While these benefits help advance a business case, it is recognised that the sample is skewed to essential health, social care and education professionals. Consequently, further research is necessary to generalise the findings to managing the RTW of more diverse occupational groups, especially those who have been classified as key workers, including delivery drivers, supermarket staff and care workers, as a way of managing skill shortages within the wider workforce. The industry skew could partly explain the sample's high proportion of female workers, mirroring that within health and social care within the UK and more widely [27]. A higher proportion of female participants also reflects the predominance of women reporting long COVID [28].

Although the sample size is small due to the study's exploratory nature, the fit of findings to the biopsychosocial rehabilitation framework implies transferability across organisations, industries and public/private sectors [17]. Furthermore, while this study has unpacked some of the challenges that workers experiencing long COVID have encountered on RTW, there was evidence that some workers found the process straightforward. Accumulating authentic case studies supporting this narrative could strengthen the case for post-COVID-19 WA.

Moreover, the findings indicate that regularly updated guidance with a suite of workplace accommodations is necessary to support individual WA/RTW trajectories, covering the varying job contexts in which these operate, and the emerging evidence on potential syndromes underpinning long COVID. Some demarcation may be necessary according to: any lasting pathophysiological damage, degree of cognitive dysfunction, psychological trauma, and use of physical activity given the continued debate surrounding its use for chronic-fatigue-syndrome rehabilitation and post-viral fatigue [29]. To optimise VR utility, further exploratory research may be warranted to determine whether RTW obstacles varied according whether SARS-CoV-2 was contracted at or outside work, or according to type of health care accessed, such as long-COVID clinics or occupational health service.

In judging if contemporary VR guidance derived from evidence for chronic health conditions that precedes the COVID-19 pandemic [14,15] is fit for purpose for accommodating post-COVID-19 syndrome, these findings highlight some nuances that deserve consideration. Firstly, workplace accommodations are usually advocated as temporary [14,17]. The present findings underscore a need for flexible, longer-term and regularly reviewed accommodations to allow for the potentially protracted, unpredictable multi-system nature of post-viral symptoms.

Secondly, an early RTW is recognised as necessary for mitigating potential long-term sickness absence and disability [17], and calls are made for facilitating working *while* recovering on the premise that should permit more rapid resumption of usual WA [18]. The present study highlights that initial RTW planning might need to select tasks that have reduced personal or public safety risks or lower cognitive complexity: this could allow cognitive functioning levels to reemerge unhampered by the pressures to perform in safety critical roles at the point of RTW. Tasks requiring meta-cognitive skills or patient/client interaction may need to be deferred until cognitive functioning is sufficient.

Third, realistic personal and workplace expectations about ability to work have been highlighted as necessary [18, 26]. Expectations that a worker recovering from COVID-19 should be fully productive on RTW might need to be countered to prevent unhelpful pressures on the rehabilitation process. While full fitness before RTW should not be expected, the returning worker, their line manager and peers, leadership and OH practitioners may need to modify beliefs around the need for full fitness and productivity. Persuasive COVID-awareness raising programmes targeting such attitudes could help create more rehabilitation conducive organisational cultures.

Given the reported variation in recovery experiences, these findings imply that supporting workers' autonomy to self-manage job demands alongside symptoms could provide them with the flexibility to meaningfully fulfil at least some job requirements while recovering [26]. Findings also reinforce the view that line managers should play an active role co-developing RTW plans with workers, with OH and HR providing specialist input required [18]. Given the individualistic nature of the work-relevance of post-COVID-19 symptoms and

their accommodation needs, line managers and workers are best placed to work out the optimal requirements to regain WA, assuming clinical screening where appropriate.

Finally, these findings indicate a long-term and flexible approach to workplace health management as potentially important for allowing the large number of workers apparently struggling after COVID-19 to sustainably regain WA, including those with pre-existing conditions. Other conditions presenting similarly unpredictable symptom patterns and RTW obstacles such as chronic fatigue syndrome [13, 30] should also benefit from such flexibility. A skew to female participants and essential workers could make this approach pertinent to health and social care. Use of the biopsychosocial framework to overcome multi-level obstacles to WA, coupled with support for working-while-recovering wherever reasonable, should afford a more person-centred approach that can contend with the unpredictable characteristics of post-COVID-19 symptoms.

References

1. Office of National Statistics. Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK:7 October 2021. <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/prevalenceofongoingsymptomsfollowingcoronaviruscovid19infectionintheuk/7october2021> (2 Nov 2021, date last accessed)
2. NICE, SIGN, RCGP. *NICE guideline [NG188], COVID-19 rapid guideline: managing the long-term effects of COVID-19 2020*. <https://www.nice.org.uk/guidance/ng188> (29 Mar 2021, date last accessed)
3. Carfi, A., Bernabei, R., Landi, F., Group ftGAC-P-ACS. Persistent Symptoms in Patients After Acute COVID-19 *Journal of the American Medical Association*, 2013; 324: 603-605.
4. Trade Union Congress (TUC). Worker's Experience of Long Covid <https://www.tuc.org.uk/research-analysis/reports/workers-experiences-long-covid> [2 Nov 2021, date last accessed).
5. Ebener M, Hasselhorn HM. Validation of Short Measures of Work Ability for Research and Employee Surveys. *Int J Environ Res Public Health*. 2019;1618:3386.
6. Gaber TAZK, Ashish A, Unsworth A. Persistent post-covid symptoms in healthcare workers, *Occupational Medicine*. 2021;71:144–146.
7. National Health Service (NHS). *Long-Covid Clinics* <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2021/06/C1312-long-covid-plan-june-2021.pdf> (15 Dec 2021, date last accessed).
8. Ladds E, Rushforth, A, Wieringa S, Taylor S, Rayner C, Husain L, & Greenhalgh T. Persistent symptoms after Covid-19: qualitative study of 114 “long Covid” patients and draft quality principles for services. *BMC health services research*. 2021: 20: 1-13.
9. Bawden, T. People with long covid facing months-long wait for treatment specialist NHS clinic. <https://inews.co.uk/news/nhs-figures-reveal-a-huge-shortfall-in-long-covid-treatment-1310377> (16 Dec 2021, date last accessed).
10. Waddell, G, Burton, AK. *Is work good for your health and well-being?* 2006. TSO, London, UK. ISBN 9780117036949
11. Waddell G, Burton AK, Kendall N. *Vocational rehabilitation – what works, for whom, and when?* 2008. TSO, London UK. ISBN 9780117038615
12. Fadyl JK, Anstiss D, Reed K, Levack WMM. Living with a long-term health condition and seeking paid work: qualitative systematic review and thematic synthesis. *Disability and rehabilitation*. 2020:1-11.
13. Joosen M, Frings-Dresen M, Sluiter J. Process and Outcome Evaluation of Vocational Rehabilitation Interventions in Patients with Prolonged Fatigue Complaints. *International Journal of Behavioral Medicine*. 2011;18:160-171.
14. Faculty of Occupational Medicine. *Guidance for health care professionals on return to work for long-covid*. 2021. https://www.fom.ac.uk/wp-content/uploads/longCOVID_guidance_04_small.pdf [(14 June 2021, date last accessed)

15. Society of Occupational Medicine. *Covid-19 Return to Work Guide. For recovering workers* [https://www.som.org.uk/COVID-19 return to work guide for recovering workers.pdf](https://www.som.org.uk/COVID-19%20return%20to%20work%20guide%20for%20recovering%20workers.pdf) [14 June 2021, date last accessed]
16. Grant, M., Rees, S., Underwood, M. *et al.* Obstacles to returning to work with chronic pain: in-depth interviews with people who are off work due to chronic pain and employers. *BMC Musculoskeletal Disord.* 2019; 20: 486
<https://doi.org/10.1186/s12891-019-2877-5>
17. Waddell, G., Burton, AK. *Concepts of rehabilitation for the management of common health problem.* 2004. TSO, London UK. ISBN 0 11 703394 4
18. Kendall, N., Burton, K., Lunt., J., Mellor, N. Daniels, K. *Development of an Intervention Toolbox for Common Health Problems in the Workplace*, HSE Research Report RR 1053. 2016. <https://www.hse.gov.uk/research/rrhtm/rr1053.htm> (30 March 2021, date last accessed)
19. Qualtrics. *Products*,
<https://www.qualtrics.com/uk/?rid=ip&prevsite=en&newsite=uk&geo=GB&geomatch=uk> (30 March 2021, date last accessed)
20. Public Health England (PHE) *Long Term Conditions, 2020*
<https://fingertips.phe.org.uk/indicator-list/view/NnltYiPSX0> (12 December 2020, date last accessed)
21. National Health Service (NHS). Myalgic encephalomyelitis or chronic fatigue syndrome (ME/CFS) <https://www.nhs.uk/conditions/chronic-fatigue-syndrome-cfs/> (16 December 2021, date last accessed)
22. Hsieh HF, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res.* 2005; 9:1277-88.
23. Weber, R. P. *Basic content analysis* (2nd ed) 1990. Sage Publications, Inc.
24. Bowling, A. *Research methods in health: investgating health and health services.* 2014. 4th edition , Maidenhead, GB. McGraw Hill; Open University Press, p. 536.
25. Lunt, J., Hemming, S., Elander, J., Baraniak, A., Burton, K., & Ellington, D. Experiences of workers with post-COVID-19 symptoms can signpost suitable workplace accommodations. *International Journal of Workplace Health Management*, 2022; 15: 359-374.
26. Sampere M, Gimeno D, Serra C, Plana M, López JC, Martínez JM, Delclos GL, Benavides FG. Return to work expectations of workers on long-term non-work-related sick leave. *J Occup Rehabil.* 2012; 22 :15-26.
27. Organisation for Economic Co-operation and Development (OECD). Women are well-represented in health and long-term care professions, but often in jobs with poor working conditions <https://www.oecd.org/gender/data/women-are-well-represented-in-health-and-long-term-care-professions-but-often-in-jobs-with-poor-working-conditions.htm> (30 May 2022, date last accessed)
28. Statista. Number of people with long COVID symptoms in the United Kingdom (UK) as of April 2022, by gender and extent of activity limitations

<https://www.statista.com/statistics/1257434/people-with-long-covid-in-the-uk-by-gender-and-activity-limits/> (30 May 2022, date last accessed)

29. British Medical Journal (BMJ). NICE backtracks on graded exercise therapy and CBT in draft revision of CFS guidance BMJ 2020;371:m4356
30. Huibers MJ, Leone SS, Kant IJ, Knottnerus JA. Chronic fatigue syndrome-like caseness as a predictor of work status in fatigued employees on sick leave: four year follow up study. *Occup Environ Med.* 2006, 63:570-2.

Table 1: Survey Summary

<i>Health Status (5 items)</i>	
Quantitative questions	<ol style="list-style-type: none"> 1. Which of the following best describes your COVID- 19 experience at its most serious? (<i>Mild/ at home, severe at home, hospitalised, hospitalised and ICU</i>) 2. How long ago? (<i>current, within the last month, between a month/ 6 months ago, more than 6 months ago</i>) 3. COVID-19 Duration? (<i>less than a week, one -two weeks, - two weeks – one month, between a month and 6 months, more than 6 months</i>)
Open Ended	<ol style="list-style-type: none"> 4. How would you say your health has been affected by COVID-19? 5. How would you say your existing conditions have been affected by COVID-19?
<i>Work ability/ Return-to-work Status (7 items)</i>	
Quantitative (RTW)	<ol style="list-style-type: none"> 6. Have you resumed work? (<i>fully, partially, not yet, not anticipating to resume work, did not stop working</i>) 7. Psychological Work ability - How do you rate your current work ability with respect to the psychological and demands of your work? (<i>1 very good, 2 rather good, 3 moderate, 4 rather poor, 5 poor</i>) 8. Physical Work ability: How do you rate your current work ability with respect to physical demands of your work? (<i>1, very good, 2 rather good, 3 moderate, 4 rather poor, 5 poor</i>)
Open ended (RTW)	<ol style="list-style-type: none"> 9. If you have resumed work, how easy have you found it? (For example, how easy was it continue working without any further sickness absence).
<i>Return-to-work Obstacles (1 item, 4 parts)</i>	
Quantitative (RTW)	<ol style="list-style-type: none"> 10. What do you view as the obstacles that make or have made return-to-work harder (<i>individual (e.g. health, psychological), job, support (managerial), organisational, external</i>)? <ol style="list-style-type: none"> a. Individual (<i>e.g. health, psychological</i>), b. Job (<i>e.g. working patterns, physical and psychological demands</i>) c. Support (<i>e.g. line manager, peer, human resources, occupational health</i>), d. <i>Organisational</i> (<i>e.g. shared attitudes about health at work, reporting systems, absence management procedures, flexible working polices, communication practices, job security</i>) e. <i>External</i> (<i>e.g. access to health care, ability to get to work, family support</i>)
<i>Return-to-work Enablers (1 item, 4 parts)</i>	
Open Ended	<ol style="list-style-type: none"> 11. What enablers would make or have made your return-to-work easier? <ol style="list-style-type: none"> a. Individual (<i>e.g. health, psychological</i>), b. Job (<i>working patterns, physical and psychological demands</i>) c. Support (<i>e.g. line manager, peer, human resources, occupational health</i>), d. <i>Organisational</i>: (<i>e.g. shared attitudes about health at work, reporting systems, absence management procedures, flexible working polices, communication practices, job security</i>) e. <i>External</i>. <i>e.g. (access to health care, ability to get to work, family support)</i> 12. <i>How do you think these obstacles enablers differ according to whether you are having to return-to-work or return to working (at home)?</i> 13. How long do you think any adjustments for making resumption of work easier should reasonably last?
<i>Benefits for employees</i>	
	<ol style="list-style-type: none"> 14. What do you think might be/are the benefits for the employer in supporting your recovery?

Table 2: Summary Statistics

Variables*	Frequency (%)**
Pre-existing conditions (n=75)	
<i>Respiratory</i>	14 (10)
<i>Diabetes</i>	2 (1)
<i>Cardiovascular problems</i>	3 (2)
<i>Musculoskeletal problems</i>	5 (3)
<i>Mental Health</i>	9 (6)
<i>Other</i>	42 (29)
COVID-19 experiences at its worst? (n=132)	
<i>Mild/Moderate at home</i>	50 (35)
<i>Severe at home</i>	65 (45)
<i>Hospitalised</i>	17 (12)
COVID-19 How long ago? (n=132)	
<i>Current</i>	4 (3)
<i>Within the last month</i>	6 (4)
<i>Between a month and 6 months ago</i>	28 (19)
<i>More than 6 months ago</i>	94 (65)
COVID-19 Duration (n=132)	
<i>1-2 weeks</i>	10 (7)
<i>2 weeks - 1 month</i>	8 (6)
<i>1 - 6 months</i>	35 (24)
<i>6 months+</i>	79 (55)
Range of post-viral symptoms experienced (n=132) (excessive fatigue, confusion, trouble with concentrating/brain fog, headaches, aches and pains in muscles, stiff joints, sore throat, chest pain, rash, upset stomach)	
<i>All of these</i>	22 (15)
<i>Most of these</i>	81 (56)
<i>Some of these</i>	21 (15)
<i>A few of these</i>	6 (4)
<i>None</i>	2 (1)
Have you resumed work? (n=88)	
<i>Fully</i>	21 (15)
<i>Partially</i>	23 (16)
<i>Not yet</i>	38 (26)
<i>Not anticipated</i>	5 (3)
<i>Did not stop working</i>	1(1)
Physical work ability (n=88)	
<i>Very good</i>	3 (2)
<i>Rather Good</i>	8 (6)
<i>Moderate</i>	22 (15)
<i>Rather poor</i>	30 (21)
<i>Poor</i>	25 (17)
Psychological work ability (n=88)	
<i>Very good</i>	2 (1)
<i>Rather Good</i>	14 (9)
<i>Moderate</i>	25 (17)
<i>Rather poor</i>	27 (19)
<i>Poor</i>	20 (14)

*(No. of responses to specific question)

**Percentage of full sample (N=145) provided

Table 3: Leading Return to Work Obstacles

Obstacle (Anticipated or Experienced)	n	Example Quotes
Individual Level (physical or psychological factors)		
Fatigue	29	<i>"Need for huge amount of rest...never had an illness I could not push through before."</i>
Social distancing	23	<i>"Not able to social distance at work with patients"</i>
Poor concentration	20	<i>"Each two days of work could cause me three days of brain fog and short-term memory loss"</i>
Relapsing nature of symptoms	13	<i>"Rollercoaster nature of symptom severity making it impossible to plan work"</i>
Safety (personal or patient safety)	12	<i>"Cognitive symptoms made nursing unsafe."</i>
Expectations/professional Identity	10	<i>"Feeling worried about not being able to perform as well as I used to"</i>
Job (factors in the immediate work environment)		
Symptom interaction with physical job demands of a given job	29	<i>"Absolutely could not safely care for patients nor handle the physicality of nursing"</i>
Control over job pressures	28	<i>"High pressure in my own caseload, which has increased since the original lockdown"</i>
Working patterns	17	<i>My days are 7.30 to 6.40pm at the earliest-there is no way I would manage these "</i>
Symptom interaction with cognitive job demands of a given job	12	<i>"Working IT systems is difficult with brain fog and concentration is hard too"</i>
Work Support (managerial support factors)		
Human Resources/ Occupational Health Support (-) (the quantity and quality of human resources services and/or occupational health services support undermines work ability)	14	<i>"They paid me for the time, I was positive with Covid but now I have long Covid don't know what pay I will get if any</i>
Line management (-)	12	<i>"Not believing you and saying you are over exaggerating, telling you to push yourself, saying you risk losing your job if phone in sick."</i>
Peer/colleague behavior (-)	11	<i>"Peers and colleagues are already burnt out and exhausted from the pandemic, will face resentment from being out for so long"</i>
Organisational Support (wider organisational factors)		
Sickness absence policies	26	<i>"Organisation is generally supportive... but the HR policies are rigid e.g. limited company sick pay in first few years, refusal to furlough."</i>
Organisational Culture (Awareness of COVID and collective attitudes about health and work)	27	<i>"It would be great if people understood fatigue is not the same tiredness"</i>

Job Security		<i>"Once you are back, you are back fully or not at all"</i>
	15	<i>"Not sure if they will keep my job open until I'm able to return that will be very phased."</i>
External factors (societal factors that affect ability to work)		
Access to suitable health care	33	<i>"I struggled to get GP appointment. Waiting for appointment at long-Covid clinic for 3 months"</i>
Transport issues	26	<i>"I would need to be given special permit to allow me to park at work"</i>

Table 4: Leading Return-to-Work Enablers

Enabler	n	Example Quotes
Individual		
Self-management	27	<p><i>"Resting when I can and ensuring I take medication as prescribed"</i></p> <p><i>"Asking for help and being open about my situation"</i></p> <p><i>Self-coaching"</i></p> <p><i>"Knowing limits"</i></p> <p><i>"Understand which medications will help me"</i></p>
Pacing and taking breaks	15	<p><i>"If I become tired/ near my limit I need to stop immediately"</i></p> <p><i>"Taking regular breaks"</i></p>
Job and Work Support		
Flexible working arrangements	32	<p><i>"Being allowed to do short periods of work when well"</i></p> <p><i>"Reduced hours"</i></p> <p><i>"Leaving early if needed"</i></p> <p><i>"Later starts"</i></p>
Graded return to work	30	<p><i>"My GP has recommended working, 2 hours a day initially, one hour teaching and one hour admin"</i></p>
Changing jobs/tasks/responsibilities	29	<p><i>"Obviously, I need to get back to patient facing work but I think the non-complex stuff would be better to start back in"</i></p> <p><i>"Changes in 'workload,' 'roles,' 'duties' or 'caseload"</i></p> <p><i>"Having tasks within my ability."</i></p> <p><i>Initially undertaking "repetitive work,"</i></p> <p><i>"Job sharing"</i></p> <p><i>"Periods of supernumerary"</i></p> <p><i>"More complex tasks deferred"</i></p>
Improve line management competency	31	<p><i>"Managers should have a conversation with you about what may help and how dealing with fatigue, pain etc can be lessened"</i></p> <p><i>"Provide a prescriptive approach"</i></p> <p><i>"Developing return-to-work plans in partnership"</i></p> <p><i>"Have "conversations with employees about what will help"</i></p> <p><i>"Have interpersonal skills training"</i></p> <p><i>"Keeping in contact"</i></p> <p><i>"Help to prioritise workloads and backlogs"</i></p>
Improve occupational health/health care access and utility	14	<p><i>"Precise documentation by GPs of "symptoms within fit-notes"</i></p> <p><i>"A precise diagnosis"</i></p> <p><i>"Realistic diagnosis"</i></p> <p><i>"OH referrals"</i></p> <p><i>"OH assessments prior to return-to-work" "Cognitive function assessments"</i></p> <p><i>"Mental health support."</i></p>
Managing peer expectations	11	<p><i>"Briefing colleagues that a person back at work may not be fully recovered"</i></p> <p><i>"Providing "appropriate training" to cover for an off-sick colleague"</i></p>
Organisational		
Sickness Absence Policy Modifications	14	<p><i>"Sickness policy needs to be looked at in relation to long covid"</i></p> <p><i>"User friendly" for when unwell</i></p> <p><i>"Include bullying and harassment"</i></p> <p><i>"Discount COVID-sickness absence"</i></p> <p><i>"Ensure visibility and accessibility"</i></p>
Organisation-wide COVID-Awareness	14	<p><i>"Creating knowledge about post-Covid and organ damage"</i></p>

Organisation-wide positive attitudes about COVID and work ability	13	<i>“Culture of accepting that illness is not an inconvenience or a stigma”</i> <i>“Illness is not viewed is not an inconvenience or stigma”</i>
---	----	--

Table 5: Benefits to Employers

Benefit	n	Example Quotes
Retention of specialist skills	20	<i>“[Avoid]expending resource in hiring someone new and training them in a highly specialised area”</i> <i>“Train team members and supporting other departments”</i>
Fostering commitment	20	<i>“Knowing that they will be supported if unlucky enough to struggle with health will contribute to positive workplace, job satisfaction and productivity”</i>
Sustained return-to-work	15	<i>“Ensuring staff don’t go off again by returning too early”</i>
Productivity	12	<i>“More support might result in a quicker recovery and more productive in long run”</i>

Table 6: Potential Workplace Accommodations for Post-COVID 19 Workability

- **Self-management:** Providing workers training and time to self-manage their symptoms alongside job demands through pacing (including taking frequent breaks), developing awareness of their own limits/boundaries, noticing changes in symptoms, stress management etc.
 - **Co-developed graded RTW plans:** Co-developed between managers and workers, provide flexible graded RTW plans that allow for fluctuating symptoms that are largely unbounded by time. Include OH assessments, OH referral prior to returning to work and cognitive assessments where appropriate. Avoid large step-ups in hours or work demands.
 - **Flexible working patterns:** Including frequent breaks, reduced hours or modifications to start and finish times. Provide rest-facilities.
 - **Changes to jobs/tasks and responsibilities:** Including amended duties with more complex tasks deferred, temporary changes to role, simplified/reduced workload gradually building over time. Consider temporary job-sharing and supernumerary.
 - **Practical and emotional support from line managers:** Including practical support in prioritising workload, support with back-log, keeping in touch with workers while they are off-sick and resume working, and having compassion-based conversations about health issues. Provide communication training to support this where needed. Have regular catch-ups with a single point of contact such as a line manager.
 - **Briefing peers:** Including managing expectations on what to expect from colleagues returning to work with post-viral symptoms, providing training to enable peers to cover for off-sick colleagues allowing open discussions with colleagues and supporting reciprocity – the possibility that one day colleagues may need similar support.
 - **Modify sickness management policies:** Including making them more COVID-friendly and user-friendly, discounting COVID-sickness absence, including bullying and harassment considerations, and heightening their visibility and accessibility.
 - **Awareness raising programmes:** Running organisational wide programmes for raising awareness about the nature of COVID-recovery, creating realistic expectations about what is possible on RTW (e.g. that it may not be possible to be fully functioning straight-away, but that it is still possible to make a useful contribution in less than perfect health, e.g. by coaching others) and creating compassionate organisational cultures
-