

Higher on the mountains: dynamics of agro-pastoral practices in a low-latitude mountain system (Karkas Mts, central Iran) during the Mediaeval Warm Period and Little Ice Age

Morteza Djamali^{1,2*}

Emmanuel Gandouin¹

Arash Sharifi^{3,4}

Philippe Ponel¹

Abdolmajid Naderi-Beni²

Alireza Naqinezhad⁵

Jacques-Louis de Beaulieu¹

Elodie Brisset¹

Nafiseh Samadi⁶

Marjan Mashkour⁷

Dahvya Belkacem¹

Michelle Leydet¹

Marta Garcia⁸

Kazuyo Tachikawa⁸

François Demory⁸

Edouard Bard⁸

¹Institut Méditerranéen de Biodiversité et d'Ecologie - IMBE (Aix Marseille Univ, Avignon Univ, CNRS, IRD), Europôle de l'Arbois, 13545 Aix-en-Provence, France

²Gobabeb-Namib Research Institute, Walvis Bay, Namibia

³Department of Marine Geosciences, Rosenstiel School of Marine, Atmospheric, and Earth Science, University of Miami, Miami, FL 33149, USA.

⁴Beta Analytic Inc., Isobar Science, Miami, FL, USA

⁵Department of Plant Biology, Faculty of Basic Sciences, University of Mazandaran, Mazandaran, Iran

⁶University of Tehran, General office for developmental projects and logistics, Tehran, France

⁷Archéozoologie, Archéobotanique (UMR 7209, AASPE), CNRS, Muséum National d'Histoire Naturelle, CP55, 55 rue Buffon 75005 Paris, France.

⁸Aix Marseille Univ, CNRS, IRD, INRAE, Coll France, CEREGE, Aix-en-Provence, France

Abstract

Little is known about the impact of the Mediaeval Warm Period (MWP: 950-1250 CE) and the Little Ice Age (LIA: 1250-1850 CE) on high-altitude agro-pastoral practices and the vertical mobility of human communities in the mountain regions of interior Southwest Asia. Although the area experienced significant socio-political changes during the last millennium, the socio-environmental interactions during these climatic periods remain poorly understood. This study presents a geochronologically well-constrained, multi-proxy geochemical and palaeoecological record from a high-altitude (2500 m) wetland in the Karkas Mountains, located on the central Iranian desert plateau. All proxies, including bioindicators (pollen, non-pollen palynomorphs, fossil insects) and X-ray fluorescence (XRF) intensity variations, reveal two contrasting patterns of hydroclimatic conditions and agro-pastoral practices during the MWP and the LIA. While the MWP is characterised by intensive cereal cultivation and the probable establishment of permanent agricultural communities, the LIA shows intensified grazing pressure on mountain steppe vegetation, suggesting a shift towards a more mobile pastoral lifestyle—one that persisted into the 20th century. Interestingly, this shift coincides with the Mongol invasion of the Iranian plateau, though it may not necessarily be related to it. Based on the Gahak wetland record and regional palaeoclimatic data, we conclude that the MWP in the mountainous areas of central Iran was marked by milder winters and possibly shorter summer droughts, in contrast to the LIA, which had harsher winters and longer-lasting snow cover. Our findings highlight the significant role of high-altitude agro-pastoral communities in supplying food to the lowland urban centres of mediaeval Persia, at least during the MWP.

Keywords: Mediaeval Climate Anomaly, Little Ice Age, Cereal cultivation, Climate impact, Southwest Asia

Introduction:

The last two millennia have witnessed frequent climatic fluctuations recorded in both continental and marine archives (McGregor et al., 2015; PAGE2k Consortium, 2017). These fluctuations have impacted the mean annual and seasonal patterns of temperature and precipitation (Büntgen et al., 2016), the extent of glaciers and snow cover (Koch and Clague, 2011; Hughes, 2014; Kokh, 2015; Rowan, 2017; Kjær et al., 2022), peat accumulation (Magnan et al., 2018), lake and wetland water levels (Del Socorro Lozano-Garcia et al., 2007), and drought dynamics (Flohr et al., 2017; Fleitmann et al., 2022). They have also impacted human societies by affecting agricultural production (e.g. Pfister, 1981; Pfister and Brázdil, 2006; Peregrine, 2020; Waldinger, 2022).

Although there is a consensus on the global nature of these climatic fluctuations, their hydroclimatic consequences (e.g. glacier mass balance) appear to have varied regionally (Matthews and Briffa, 2005). The social vulnerability to the economic impacts of these hydroclimatic variations could thus have differed even over short geographical distances, as evidenced in central Europe where two adjacent regions with differing social organisations, topographic setting, and microclimatic conditions reacted very differently to the Little Ice Age (LIA) fluctuations (Pfister and Brázdil, 2006). Such spatial variations highlight the need for multiple case studies to understand the spatio-temporal variations in socio-environmental

interactions related to these significant late Holocene climatic phases (Matthews and Briffa, 2005; Pfister and Brázdil, 2006).

In Europe and North America, numerous studies have investigated the effects of the Mediaeval Warm Period (MWP) and LIA on human socio-economy (Thomson and MacDonald, 2020; Waldinger, 2022). However, little is known about the extent of their impacts on societies in other parts of the world, including interior Southwest Asia. Both intervals have left visible ecological and geological fingerprints in various sedimentary archives of the region (Kroonenberg et al., 2007; Ramezani et al., 2008; Djamali et al., 2009; Ramezani, 2013; Leroy et al., 2013; Haghani et al., 2015; Flohr et al., 2017; Bayer Altin and Kayan, 2020). Indeed, the interior of Southwest Asia, including the Iranian plateau, is among the most water-stressed regions of the world (Barlow et al., 2015; Emadodin et al., 2019), where the climatic anomalies of the Common Era have repeatedly impacted societies throughout history (Büntgen et al., 2016; Gurjaskaite et al., 2018; Fleitmann et al., 2022). However, the potential of natural and documentary archives in this region has not yet been fully explored to investigate how these frequent climate variations, characterised by numerous extreme events, may have affected human socio-economy during times of large-scale migrations, invasions, and conflicts in mediaeval Persia.

In this study, we present a multi-proxy record of geochemical, palynological, and entomological data with a robust chronology from a high-altitude (>2500 m), low-latitude (33°N) mountain site, with the aim of understanding the socio-environmental interactions inferred from variations in the proxy data. Our specific objectives are as follows:

1. To reconstruct the climatic variations of the last two millennia and their impacts on wetland hydrology and sedimentation,
2. To track vegetation dynamics and floristic changes in the mountain steppes surrounding the wetland in response to these climatic variations and the dynamics of agro-pastoral practices, and
3. To discuss the socio-cultural implications of the reconstructed climatic and ecological changes.

Study area

Gahak wetland is a semi-permanent shallow wetland located in the Karkas Mountains of central Iran (Figs. 1 & 2). The wetland (33°38'8.31"N; 51°29'50.78"E; 2517 m a.s.l.), with a catchment basin of approximately 21 km², is surrounded by high mountains reaching up to 3224 m (Mt Kolah Barfi) and receives water from snowmelt, run-off, and numerous small springs. Traces of a qanat system can be identified at the western end of the wetland (Fig. 2B). The wetland's surface area can vary substantially on a seasonal and inter-annual basis depending on the amount of annual precipitation. The wettest part of the wetland, in the east, is waterlogged almost all year.




Figure 1 Location of Gahak wetland in the Karkas Mountains (grey dotted line “KM”) in western central Iranian plateau is marked with a red star. Black stars denote the regional paleoclimate records referred to in this study. Climate diagrams of the city of Kashan in the desert and the Abyaneh village in the mountain show marked

changes in annual/seasonal precipitation and temperature as well as the duration of dry season along the desert-mountain gradient. Red horizontal bars show the number of dry months (8 for Kashan and 6 for Abyaneh). Temperature values in bold to the top-left of the diagrams correspond to the mean of the maxima of warmest month of the year. Temperatures to the bottom-left of the diagrams represent the absolute minimum temperature of the whole period (bold italic) and mean of the minima of coldest month (narrow italic).

Climate. Gahak is close to meteorological station of the mountain village of Abyaneh (Fig. 1, lower right panel) with 213 mm/yr and 11.9°C of annual precipitation and mean annual temperature, respectively. Although located at high elevations (>2500 m.a.s.l.), the dry season lasts for six months from May to October. Four months of the year (December to March) experience frosts. The peak of precipitation happens in late winter, early spring. There is a steep climatic gradient from the high mountains to the desert environments surrounding Karkas Mts. In Kashan (Fig. 1, lower left panel), the duration of dry season increases to eight months, annual temperature goes up to 19.1°C and annual precipitation is 138 mm/yr far below the defining limits of a true desert environment (Quinn, 2009). In Global Bioclimatic Classification system, Kashan falls within Mediterranean desertic-continental while the high mountain areas of Karkas display a Mediterranean xeric to pluviseasonal-continental bioclimate with shorter dry seasons (Djamali et al., 2011).

Vegetation. The upland vegetation in the 2000-3500 m elevation range is covered by a typical species-rich Irano-Turanian Mountain steppe dominated by *Artemisia* cf. *aucheri*, *Cousinia* spp., *Euphorbia* spp., *Bromus* spp., *Stipa turkestanika*, *Eryngium* sp., *Astragalus* spp., *Acantholimon* sp., *Acanthophyllum* sp., *Achillea* sp., and *Echinops* sp. (Fig. 3). Submitted to increasing grazing pressure at lower elevations, *Astragalus* spp. become the dominant plant and together with *Artemisia aucheri* form a community. This *Artemisia* species is different from *A. sieberi* which is widely distributed in desert areas surrounding Karkas Mountains (Djamali et al., 2020). Some large and old trees are cultivated including *Populus* and *Salix* which timber are found in the ruins of unknown-age stone buildings around the wetland.

Three vegetation zones can be respectively distinguished within the Gahak wetland which is indeed a mountain fen: open permanently-waterlogged zone (easternmost corner of the wetland), semi-permanently wet peatlands, and seasonally flooded peripheral zone. The open waters of the eastern part of the wetland are surrounded by emergent *Phragmites australis* followed by submerged aquatic plants of *Ranunculus trichophyllus*, *Zannichellia palustris*, and *Chara* sp. The peat surface is mainly dominated by *Carex orbicularis* subsp. *kotschyana*, *Trifolium pratense*, *Agrostis stolonifera*, *Blysmus compressus*, *Pedicularis sibthorpii*, *Dactylorhiza umbrosa*, *Festuca rubra*, *Poa pratensis*, *Eleocharis quinqueflora*, *E. uniglumis*, *Cirsium* sp., *Triglochin palustris*, *Juncus gerardii*, *Plantago maritima*, *P. lanceolata*, and *Scorzonera* sp. In the hygromorphic soils close to surrounding upland dry steppes, the following plants are frequently observed on the peripheral zone: *Hordeum* cf. *violaceum*, *Plantago maritima*, *Mentha longifolia*, *Glaux maritima*, *Trichophorum pumilum*, *Medicago* spp., *Bolboschoenus affinis*, *Alopecurus pratensis*, *Juncus articulatus*, *Carex songarica* and *Juncus inflexus*. The latter zone is currently (July 2018) overgrazed and it is expected that livestock grazing will advance to the peatland at the end of the dry season.

Geology. The geology of the Gahak drainage basin is dominated by both carbonate and siliciclastic rocks ranging from Triassic to Tertiary (Fig. 2A). The immediate slopes have outcrops composed of Jurassic to Cretaceous conglomerates, sandstones and limestones. However, the Eocene volcanoclastics (andesites and tuffs) constitute the largest surface outcrops of geological formations in the catchment.

Agro-pastoral practices. Today, the wetland surroundings form the summer highland pasture ('yaylak') for the semi-nomadic people of Kashan area who practice transhumance and high elevation agro-pastoralism during the summer months. The grazing of highland pastures by livestock (sheep and goat) starts from April to October when the mobile agro-pastoral communities of Kashan suburbs are present on the site. No rain-fed wheat farming is currently practiced because of the insufficient spring rainfall. The winter wheat is sown in October-November and the germinated plants are left the whole winter covered under snow for several months. The wheat fields need to be irrigated three or four times in late spring and early summer to compensate for the long dry season of the area lasting six months (Fig. 1). The wheat grains are harvested from August to September. In late spring and summer, many vegetables including eggplants, cucumber, and tomatoes are also cultivated and irrigated using water from the wetland and surrounding springs.




Figure 2- Geological (A) and Geomorphological (B) maps for Gahak Wetland area with the location of the coring site. Panel B is a zoom on the inset rectangle to the right of panel A. Geology map is based on the 1:100,000 geological map of Kashan (Radfar, 1993). Geomorphological map is based on Google Earth images and field observations.




Figure 3 Gahak wetland photographed in Spring 2018 as viewed from the North. Coring site is located in the water-logged area to the left of the photo (arrow).

Materials and Methods

Field works. Three sediment cores were extracted with stratigraphy intact using a Russian peat borer. They were labeled Gahak 1, Gahak 2, and Gahak 3 each in two drives not exceeding 130 cm in total. We selected the most completely retrieved sister core sections (Core 2-1 and 3-2) to create a composite section of 120 cm length that we analyzed in this study. The cores were taken from the peat surface in the eastern part of the wetland which remains wet most of the year (Fig. 2B). A red soil horizon located at 120-130 cm horizon marked the base of all cores and possibly the beginning of wetland sedimentation. The cores were sealed in non-reactive plastic sheets, placed within halved PVC core liners and stored in a cold chamber. In July 2018, a botanical excursion/survey was also conducted around the Gahak Wetland to prepare a checklist of main aquatic and semi-aquatic plants colonizing the wetland and its peripheral area.

X-Ray Fluorescence (XRF) and magnetic core scanning. The Gahak 2-3 composite section was subsampled longitudinally by U-channels with a 20 mm x 20 mm cross-sectional area. A high-resolution, non-destructive XRF core scanning was performed at 5-mm resolution using an ITRAX scanner (Cox Analytical Systems; Croudace et al., 2006) to measure the relative abundance of Si, K, Ca, Ti, Mn, Fe, Rb, Sr, Zr, and incoherent/coherent scattering ratio (hereafter inc/coh, indicator of organic matter and pore water content). The Mo tube was used as an X-ray source at 30 kV, 45 mA with 15 s of counting. The U-channels were then scanned with a recently-developed high-resolution magnetic scanner to measure the variations in the intensity of artificially-induced isothermal remanent magnetization (IRM) (Demory et al., 2019). This new technique is more performant in detecting more subtle changes in magnetic properties of sediments compared to the commonly measured magnetic susceptibility. XRF measurements were performed at Inorganic Geochemistry Unit and magnetic field (IRM) measurements were performed at Rock Magnetic Laboratory at CEREGE (Aix-en-Provence, France).

Chronology. A total of seven samples (3 peat and 4 organic mud) were selected along the core and sent to Poznan Radiocarbon Laboratory for ^{14}C analysis. Ages and their associated errors were calibrated using IntCal20 calibration dataset (Reimer et al., 2020) in OxCal version 4.4 program (Bronk Ramsey, 1995) and age-depth models were established using both CLAM and Bchron R packages (Blaauw, 2010; Parnell et al., 2008). Ages and their associated uncertainties were reported in both 1-sigma and 2-sigma ranges in calendar years after Common Era (cal yr CE).

Palynological sample preparation. A total of 24 subsamples for pollen analysis were taken every 5 cm along the core each presenting one to two cm^3 . Samples were prepared for pollen and (NPPs) Non-Pollen Palynomorphs analysis using standard extraction technique (Moore et al., 1991). Pollen percentages were calculated using Pollen Sum (sum of upland plants excluding aquatics/subaquatics) for terrestrial plants and Total Pollen Sum (sum of upland plants including aquatics and sub-aquatics) for aquatic plants. The distinction between the above ecological categories was based on both botanical literature and our field observations. Non-Pollen Palynomorphs (NPPs) (and *Triticum*-type pollen) were quantified as concentrations (true number of objects per cm^3 of sediment). To calculate pollen and NPP concentrations, *Lycopodium* tablets (Batch no. 1030, University of Lund) were added to samples and the formula presented by Stockmarr (1971) was used. Pollen percentage and NPP concentration diagrams were created in Tilia/TGView (Grimm 2004/2005). Pollen identifications were assisted using pollen atlases of Europe and the Mediterranean (Reille 1992, 1995, 1998; Beug, 2004; van Zeist and Bottema, 1977) and the Pollen Reference Collection of Persia and Arabia hosted at IMBE (Aix-en-Provence, France). NPP identifications were assisted by a number of publications (van Geel and Aproot, 2006; Cugny, 2011; Shumilovskikh and van Geel, 2020, among others).

Fossil insects. The whole core was cut into 10-cm thick slices for extraction of fossil insect remains. The fossil insect extraction was performed following the method recommended by Coope (1986). In summary, samples were wet-sieved at 300 μm mesh, the residues were floated on paraffin and sorted under a binocular microscope at low magnification. The insect remains were then preserved in alcohol or glued to cardboard pieces. Taxonomic identifications were made by direct comparison with modern specimens of a reference collection. Definition of the ecological groups of the fossil coleopteran assemblages is based on the modern literature, mainly Koch (1989a, b, 1992).

Statistical analyses

A constrained sum-of-squares cluster analysis (CONISS; Grimm 1987) was used with the Rioja R package (Juggins 2022) to identify zones in the stratigraphy for XRF elemental abundances and entomological data. The significance of each zone was tested with the broken stick model (Bennett 1996) which revealed seven significant zones. However, we slightly modified the CONISS modeled zonation and reduced the number of zones from seven to five by taking into account the downcore variations of elements and their geochemical response to various environmental conditions. CONISS analysis for pollen and NPP diagrams was performed

using the ‘coniss’ program included in Tilia/TGView (Grimm, 1977). We refer to pollen zones by “P-” and to NPP zones by “NPP-” (e.g. P-A1 & NPP-A1 zones).

In order to compare data with other regional paleoclimate records, a Loess smoothing procedure (Local Polynomial Regression) was applied between age and each variable such as Ti XRF intensity data from Gahak (this study) and Neor (Sharifi et al., 2015) as well as $\delta^{18}\text{O}$ from Gejkar (Flohr et al., 2017). The smoothing parameter is equal to 0.05 which represents the proportion of observations used to local regression. Loess smoothing procedure (Cleveland et al., 1992) was performed in R.

And finally, a one-way non parametric ANOVA of Kruskal-Wallis and related Post Hoc test, the Conover’s all-pairs test (Conover and Iman, 1979), was performed on Ti intensity variations, sum of farming-associated pollen, and pastoral pollen indicators. The objective was to test difference between the main time intervals represented in Gahak record; the Mediaeval Warm Period, the Little Ice Age, and the two episodes pre-dating and post-dating these phases. Results are classically represented with box-plot and letters that summarize significant difference or similarity between groups per periods.

Spectral and Wavelet Analysis

To track the degree of coherence between elemental pairs from the XRF measurements as a function of time, wavelet semblance analyses were performed based on the continuous wavelet transform method (Cooper and Cowan, 2008) using a modified MATLAB script. Time series were detrended by subtracting the least-square best-fit line from the original data using the built-in MATLAB command. Furthermore, time series were linearly interpolated at an equally spaced time step prior to the analysis.

Dominant periodicities within an unevenly spaced time series were identified by utilizing the Lomb-Scargle periodogram (Lomb, 1976; Scargle, 1982) approach using PAST software (Hammer et al., 2001) version 3.22. Prior to the analysis, the data was detrended using the built-in MATLAB command.

Results and interpretation

Sedimentary facies

Based on sediment textural properties, five sedimentological units can be recognized (Fig. 4). The lowest sedimentary unit (S-U1) is a massive gray clay with no visible layering and very few organic debris and pale mottling that occupies the depth of 94 to 120 cm. It seems to show a seasonally-inundated topographic depression marking the ‘pre-wetland environment’. The unit is overlaid by a 22-cm gray organic mud layer, containing a significant amount of siliciclastic minerals (S-U2). The organic mud layer is composed of lacustrine carbonate mud and contains intact aquatic snail shells and ostracods and is rich in plant debris. It indicates the

deposition in a low-energy shallow wetland to lake environment and corresponds to the ‘wetland formation phase’. The organic mud layer gradually changes into a carbonate mud with white color and without significant fossil content. This sedimentary unit (S-U3) is most probably deposited in an oligotrophic freshwater shallow water body and marks a phase of ‘lake formation’ (S-U4). The upper part of the core from 58 cm to the surface comprises a fibrous peat accumulation that contains a sandy peat layer around the depth of 10 cm. The contact with S-U3 is sharp and shows an abrupt change in hydrological conditions.

Radiocarbon chronology and age-depth model

Results of the seven radiocarbon dates and their associated uncertainties are presented in Table 1.

Table 1 Detailed radiocarbon age results.

Sample	Depth (cm)	¹⁴ C lab code	material	¹⁴ C age (yr BP)	¹⁴ C posterior calibrated ages (cal CE)		
					Calibrated ¹⁴ C ages (cal yr CE)	Median (cal yr CE)	HPD (cal yr BP)
Gahak 3-2-6	6-7	Poz-118150	Peat	127.63 ± 0.37	1837 ± 66	1860	1803-1931 (79%)
Gahak 3-2-38	38-39	Poz-118149	Peat	240 ± 30	1704 ± 93	1655	1516-1822 (91%)
Gahak 3-2-49	49-50	Poz-104459	Peat	405 ± 30	1497 ± 56	1464	1430-1628 (95%)
Gahak 3-2-74	74-75	Poz-104458	Organic mud	715 ± 30	1294 ± 32	1283	1257-1387 (95%)
Gahak 3-2-76	76-77	Poz-154221	Organic mud	945 ± 30	1100 ± 41	1128	1026-1168 (95%)
Gahak 3-2-83	83-84	Poz-154222	Organic mud	965 ± 30	1094 ± 41	1115	1022-1161 (95%)
Gahak 3-2-90	90-91	Poz-118148	Organic mud	550 ± 30	1380 ± 37	1407	1316-1435 (95%)




Figure 4 Sedimentary log and age-depth model of Gahak 2/3 sediment core. The excluded age is colored in red.

The age-depth model was generated based on smoothing spline regression (Fig. 4). We also generated an age-depth model based on Bayesian inference (not shown here), as we have previously demonstrated the effectiveness of such age models in identifying age outliers in anthropogenically disturbed small wetlands in semi-arid Iran (Djamali et al., 2021). The Bchron Bayesian model automatically excluded the age of 1380 ± 37 cal yr BP at 91-90 cm depth, which we also excluded from the Clam smooth-spline model (highlighted in red in Fig. 4). In the absence of radiocarbon ages (due to the lack of organic sediment) for the lower 40 cm of the core, we retained the down-core extrapolation based on the linear regression model proposed by Clam. The robustness of the record (e.g., between 85 and 65 cm depth) is presented as the age uncertainties, shown by the grey-filled curve, to the left of the proxy panels (Fig. 5 and Figs. 8-10).

Elemental proxies

The sedimentary profile against down-core variations in IRM field values and XRF-based relative abundance of a selection of elements including Si, Ti, Fe, K, Ca, Sr, and three elemental ratios (Zr+Rb)/Sr, Si/Ca, and Fe/K are illustrated in Fig. 5. The vertical scale is based on depth with corresponding calibrated radiocarbon ages marked for each 5cm depth intervals (both in BP and CE).

Several episodes of enhanced elemental intensities were identified along the core. Overall, the XRF zones obtained through stratigraphically-constrained zonation (SCZ) show a good correlation with major sedimentary units. The abundances of refractory elements (Si and Ti; Ca), redox-sensitive element (Fe), and mobile elements (K, Sr and Rb) show down-core covariations. The correlation is inverse between Ca and Sr intensities with the rest of the elements from the beginning of the record at 480 CE to 1470 CE. The correlation between element pairs is also evident in the cross-wavelet transform (CWT) analyses for the elemental pairs (Fig. 6 A-D) and their corresponding semblance wavelet analysis. These correlations suggest a single mechanism for Ti, Si, Fe, and K input to the Gahak wetland.

XRF-U1 (Depth: 120-94 cm / Age: 480-910 CE): Pre-wetland environment. Throughout this unit, the refractory lithogenic elements (Si and Ti) show a decreasing trend while Ca and Sr abundances show increasing values (Fig. 5). The unit is completely fit with the lowest sedimentary unit (S-U1), comprising massive gray clay possibly deposited in a poorly drained floodplain.

XRF-U2 (Depth: 94-70 cm / Age: 910-1301 CE): Wetland formation and first erosion events. This unit corresponds to the organic mud sedimentary unit. High (Zr+Rb)/Sr and Si/Ca ratios reflect the balance between clastic and carbonate components where high values typically reflect coarse grained siliciclastic terrigenous intervals (Dypvik and Harris 2001; Lo et al. 2017). The ferromagnetic minerals (see IRM curve) show significantly high values with two remarkable peaks centered around 76 and 90 cm depth (970 and 1140 CE). These peaks are interpreted as evidence of intensive and rapid soil erosion events that could be related to agro-pastoral events (see next sections).

XRF-U3 (Depth: 70-55 cm / Age: 1301-1467 CE): Lake formation. The unit corresponds to a lacustrine facies. The sediment is composed of lacustrine carbonate mud rich in plant debris and aquatic snail shells and ostracods. The transition of the XRF-U2 to XRF-U3 unit is very sharp and reflected in almost all element intensity curves well seen in downcore variation in elemental z-scores Fig. S2). Ca and Sr abundances attain their highest values in the entire record. Carbonates and calcium-rich rocks are frequently available for weathering in the catchment basin of the Gahak wetland (Fig. 2A). It seems that carbonate deposition happened through biogenic mediation and/or concentration through evaporation (Gierlowski-Kordesch, 2010). Fe/K curve shows significant peaks during this period, suggesting abrupt terrigenous input to the basin either through riverine or aeolian input (Martins et al. 2007; Mulitza et al.

2008; Lupker et al. 2010; Govin et al. 2012; Burone et al. 2013; Perez et al. 2016). The Mn/Fe curve also present the same abrupt peak during this time frame pointing toward more oxic water column and high detrital input due to either lower lake levels or increase in riverine input of oxygenated water (Makri et al., 2021). Aeolian input and inflow of water-borne clastics are other mechanisms for deposition of carbonates in such environments (Gierlowski-Kordesch, 2010). The (Zr+Rb)/Sr and Si/Ca ratios are very low and Fe/K as well as Mn/Fe ratios are high during this time interval. Since the Ca and Sr reached their highest values and the Sr-Ca plot (Fig. S1) shows a dichotomy during this time period it can be speculated that Ca and Sr were most likely transported in dissolved form through riverine input while the fine-grained terrigenous particles may have been transported via aeolian input.




Figure S1 Variation of Sr versus Ca showing different trend for 650-480 B.P. time frame compared to the rest of the record.




Figure 5 Down-core XRF records of elemental abundances (XRF intensity) for refractory (S and Ti), redox-sensitive (Fe) and mobile (K, Ca, Sr) elements as well as some elemental ratios. The dotted lines in the first six panels represent the 6-degree polynomial trend line. The inc/coh ratio is plotted to the left as a proxy for relative

organic content (peat) in sediment. Sedimentary log and magnetic field (IRM) are also plotted to compare to XRF elemental variations.

XRF-U4 (Depth: 55-15 cm/Age: 1467-1809 CE): Peat accumulation. The lake environment is replaced with a peat bog during the transition of XRF-U3 to XRF-U4. The peat would have lasted for the rest of the record until today. The peat formation and accumulation is recorded by coherent/incoherent XRF scattering ratio (inc/coh) curve (Fig. 5) even if inc/coh value is an indicator of relative abundance of lighter elements relative to heavier elements, not a direct indicator of organic matter (Chawchai et al., 2016). The upper part of the Unit XRF-U4 is characterized by a significant peak of IRM values centered around 1750 CE and major peaks of Si/Ca and (Zr+Rb)/Sr values around 1640 CE and after 1750 CE. These peaks are related to significant inputs of coarser grained siliciclastic terrigenous materials to the wetland.




Figure S2 Downcore variation in elemental z-scores.

XRF-U5 (Depth: 15-0 cm/Age: 1809-1934 CE): Anthropogenic disturbance of the peat. The unit is characterized by extremely high values of IRM field related to high content of eroded siliciclastic sands which can visually be observed intercalated in the fibrous peat. Accordingly, intensity of all refractory lithogenic elements as well as redox-sensitive, and mobile elements show peaks in this unit. Except for a small peak at the top of the record, no visible change is observed in the Fe/K ratio which corresponds with an abrupt decline in Si/Ca and (Zr+Rb)/Sr ratios potentially suggesting terrigenous input were dominated by fine grained particles.




Figure 6 Cross-wavelet transform (CWT) for elemental pairs (A-D) and their corresponding semblance wavelet analysis capture covariations in elemental abundances over time (note that in Fig. 5, the XRF records are plotted by depth). Red areas in the CWT panels indicate positive amplitude, while dark blue indicates negative amplitude. In the panels referring to semblance analysis, the degree of correlation between the relative abundances of two elements is shown in red, corresponding to a semblance of +1 (positive correlation), cyan for a semblance of zero, and dark blue for a semblance of -1 (negative or inverse correlation). For most of the record, the relative abundances of refractory and mobile elements show significant agreement (+1), except for calcium (Ca), which shows an inverse correlation. These correlations suggest a single source of elemental input to the Gahak wetland. However, the correlation between Ca and Ti is disrupted at some point after 300 years before present.

Dominant periodicities in the Gahak elemental ratio $(Zr+Rb)/Sr$, using the Lomb-Scargle periodogram (Lomb, 1976; Scargle, 1982), revealed several periodicities centred around 220, 98, 63, 48, and 22 years B.P. The 220-year periodicity aligns well with the Suess/de Vries cycle, which corresponds to 190 to 230-year solar excursions (Steinhilber et al., 2012; Ma and Vaquero, 2020). The 98-year periodicity is close to the 84- to 92-year Gleissberg cycle (Peristykh and Damon, 2003). Peristykh and Damon (2003) also confirmed that the Gleissberg cycle is modulated by the 208-year Suess/de Vries cycle, which is reflected in their combination tones, including the ~61-year cycle. They also identified a well-resolved 44.9-year periodicity as the second harmonic of the 84- to 92-year Gleissberg cycle. These cycles are close to the 63-year and 48-year periodicities observed in the Gahak record. Finally, the 22-year periodicity corresponds to the sunspot activity cycle caused by a 22-year magnetic dynamo cycle known as Hale Cycle (Mursula et al. 2002). The correlation between the results of the spectral analysis on Gahak record and well-known solar cycles suggests that the climate of uplands of the central Iranian plateau is highly sensitive to the variations in solar irradiance.




Figure 7 Lomb-Scargle periodogram for Gahak using downcore XRF $(Zr+Rb)/Sr$ ratios.

Pollen and Non-Pollen Palynomorphs (NPPs)

In total, 80 pollen taxa and 17 NPP types were identified, counted, and transformed into percentages and concentrations (grains per cm^3), respectively. A simplified percentage pollen diagram with major pollen taxa attributed to different ecological groups, is illustrated in Fig. 8. NPP concentrations and their ecological groups are also illustrated in Fig. 9. Stratigraphically-confined vertical zonation schemes of both diagrams are remarkably similar and suggest almost the same subdivision of pollen and NPP records. Although, the broken stick test suggests three major zones for each diagram, we defined five major zones by giving special importance to some key pollen and NPP indicators (e.g., *Triticum*).

All pollen spectra show the predominance of an open landscape (tree percentages remaining constantly under 10%). *Quercus* (oak) pollen has a continuous curve but displays the background values reflecting the Zagros deciduous forest in western Iran (based on our unpublished data from modern pollen traps of Kashan). Traces of cultivated trees in lower elevations (*Juglans* and *Platanus*) and desert shrubs (*Calligonum* group: *Calligonum* and *Pteroporum* spp.) growing in central Iranian plateau are also observed. Overall, pollen assemblage composition suggests an Irano-Turanian mountain steppe dominated by *Artemisia*, *Cousinia*, Fabaceae, and Plumbaginaceae constituted the regional vegetation during the last millennium (Djamali et al., 2012). However, the subtle changes in relative abundance of herbaceous pollen and particularly those pollen and NPPs associated with agro-pastoralism and wetland environment reveal remarkable changes in the landscape and in the anthropogenic activities.

P-A1/NPP-A1 (Depth: 120-97 cm/Age: 480-860 CE):

Pre-wetland environment. The vegetation record starts with high values of Cichorioideae and Asteroideae associated with significant amounts of Cyperaceae all suggesting a seasonally-flooded poorly-drained flat alluvial plain colonized by plants characteristics of hydromorphic soils (zone A1). Water table may have slightly raised at the P-A1 to A2 transition (Fig. 8) as evidenced by slightly increased pollen of Typhaceae and *Potamogeton/Triglochin*. These changes accompanied with the first peak of Cyperaceae at the same transition indicate the beginning of the wetland formation also correlated with a marked change in sediments.




Figure 8 Percentage diagram for a selection of pollen and spores from Gahak sediment core.

P-A2/NPP-A2 (Depth: 97-72 cm/Age: 860-1272 CE):

Wetland formation and establishment of a sedentary farming community. Significant amounts of Cyperaceae pollen and their associated fungi *Entorrhiza* sp., a peak of *Botryococcus*, and increasing values of Rhabdocoela during P-A2/NPP-A2 together suggest a

permanently waterlogged ecosystem with high organic productivity. *Entorrhiza* is a genus of fungi whose species are mostly obligate parasites of Cyperaceae and Juncaceae and cause galls on root tips of these families worldwide (e.g., Bauer et al., 2015; Chater and Smith, 2018; Reiss et al., 2019).

The most spectacular feature of the Gahak pollen record is a continuous peak of *Triticum* (wheat) pollen along the P-A2 to P-A3 (NPP-A2 to B1). The identification of the wheat pollen is based on morphological criteria presented in the seminal work of Beug (2004). Wheat pollen percentages and concentrations reach exceptionally high values exceeding 28% and 21000 grains/cm³, respectively (Figs. 8 and 12). The almost continuous curve (although with low values) of Solanaceae and *Polygonum aviculare*-type during the same phase is noteworthy because the species of *Solanum* (i.e., *S. nigrum* and *S. villosum*) and to some extent *Polygonum aviculare* subspecies are mostly weeds in farms and gardens of most of Iran from 0 to 2400 m a.s.l. (Mosaferi et al., 2015; Eskandari et al., 2019; Eskandari et al., 2022).

Spores of several phytopathogenic fungi also show significant peaks almost correlating with *Triticum* pollen curve (cf P-A2 to A3 with NPP-A2 to B1). Of particular interest is the almost continuous curve of the hyphopodia of *Gaeumannomyces*, a genus of sac fungi (ascomycetes) with several species well-known to cause the take-all disease infecting root system of many Poaceae and particularly the cereals (Ritz, 2005). Different former varieties of the species *Gaeumannomyces graminis* (Walker, 1972), now split into several new species including *G. graminis*, *G. avenae*, and *G. tritici* which are host-specific pathogens of cereal crops including wheat, barley, and oat (Hernández-Restrepo et al., 2016). *Gaeumannomyces* hyphopodia appear simultaneously with wheat cultivation strongly suggesting its association with the cereals cultivated in Gahak.




Figure 9 Concentration diagram for 17 non-pollen palynomorphs. Concentrations are in spore numbers per cm³ of sediment.

Another significant correlation of a non-pollen palynomorph with the *Triticum* curve is observed in *Thecaphora* spore curve (Fig. 9). Although we could not find any evidence of *Thecaphora* species directly infecting the wheat in modern literature, the genus belongs to a group of smut fungi commonly pathogens of cereal crops including the *Triticum* (Ram and Singh, 2004; Arias et al., 2021). *Thecaphora* species are reported as pathogens of several crops

all around the world but mostly the crops originated from South America. Common examples are *T. frezii* infecting peanuts and *T. solani* infecting potato. In the Old World, most of the phytopathogenic *Thecaphora* are found in Asteraceae, Caryophyllaceae and Polygonaceae families. In Iran, *T. schwarzmaniana* has recently been reported as a pathogen of wild rhubarb species *Rheum ribes*, a wild plant used as vegetable and in traditional medicine (Vasighzadeh et al., 2014). However, unlike other pollen records of Iran e.g. that of Lake Urmia (Djamali et al., 2008), we did not find *R. ribes* pollen in the Gahak record. Similarly, *T. dahuangis* has more recently been reported to infect Chinese rhubarb *R. palmatum* (Arias et al., 2021). The co-occurrence of *Thecaphora* curve with agricultural phase of Gahak suggests the possibility that this smut fungi was associated with a crop or crop-associated weed in the past.

P-A3/NPP-B1 (Depth: 72-61 cm/Age: 1272-1405 CE):

Decrease of cereal cultivation and formation of a lake system. The cereal cultivation continues with lower intensity during the P-A3 zone (NPP-B1) zones and *Artemisia* pollen starts increasing, heralding a major change in the vegetation and anthropogenic activities in Gahak which will culminate during the next pollen assemblage zone. The pollen concentration values of *Triticum* show more drastic fall compared to the pollen percentages (Figs. 8 and 12) suggesting that agricultural activities abruptly declined around 1250 CE following the most commonly accepted chronology of MWP to LIA climatic transition (e.g. Matthews and Rebuffa, 2005). The near absence of *Botryococcus* and lowest values of aquatic pollen, both indicate the oligotrophic conditions with lower organic productivity and substantially reduced aquatic vegetation. This is in accordance with the deposition of lacustrine a carbonate mud of S-U3 sedimentary unit (Fig. 4).

P-B1/NPP-B2 (Depth: 61-47 cm/Age: 1405-1559 CE):

Development of pastoralism and seasonal occupation of the site. During the P-B1 (NPP-B2), *Artemisia* became very important in the steppe vegetation dominating the pollen assemblages for about 160 years (Fig. 7). This remarkable *Artemisia* increase in the landscape is contemporaneous with a peak of the green algae *Pediastrum* showing an open water lacustrine environment (Fig. 9). Nowadays, *Artemisia* (*A. aucheri*) grows in well-drained soils which receive a high amount of snow-melt water during spring. Indeed, *Artemisia aucheri* currently forms a well-established community in the subalpine zone of Karkash Mountains with frequent snowfall. High amount of *Artemisia* pollen, peak of *Pediastrum*, and a lithology dominated by a lacustrine mud together suggest a significant amount of snowfall and reduced annual evapo-transpiration in Gahak environment during the period almost corresponding to the 15th-16th centuries CE (1400-1580 CE). Significant peaks of *Sporormiella* and some other dung-associated fungal spores (NPP-B2; Fig. 9) suggest a clear transition from sedentary agriculture to mobile pastoral life-style during the transition of the P-A3 to P-B1 (NPP-B1 to NPP-B2) transition at the very beginning of the 15th century CE.

Anthropogenic pollen indicators of pastoralism which were present with low values from the beginning of the record, start increasing from the P-B1 pollen zone until the end of the pollen

diagram (Fig. 8). In the context of the Karkas Mountains, our floristic observations of high elevation mountain steppes suggest that *Plantago lanceolata*-type, *Rumex*, *Verbascum*-type, *Cousinia*, and *Salvia pratensis*-type all indicate grazing activities. A significant increase in *Trifolium* pollen from the P-B1 onwards is noteworthy. The *Trifolium*-type pollen in the Gahak sequence seems identical to the pollen of the red clover (*Trifolium pratense*) widely cultivated in Eurasia as a fodder crop and has contributed to the feeding the livestock during the seasonal occupation of the site in summer months which is still practiced nowadays. It is one of the major plants found in the wet meadows around the Gahak wetland. Combined together, both coprophilous NPPs and coprophagous beetles (see next section) are continuously present during the upper half of the Gahak sequence suggesting a continuous intensive pastoral pressure on the ecosystems in the area since the beginning of the 15th century.

P-B2/NPP-B3 (Depth: 47- ca 10 cm/Age: 1559-1861 CE): Intensification of pastoral pressure on the mountain steppe.

The pollen indicators of grazing observed in P-B1/NPP-B2, increase and form continuous curves in the P-B2 pollen zone (Fig. 8). Significant presence of coprophagous fungi corroborates the strong pastoral pressure on the steppe vegetation during the NPP-B3 subzone (Fig. 9). A peak of Cyperaceae and presence of *Pedicularis*, suggest a densely vegetated wetland with relatively high-water tables. *Pedicularis* is currently found on the peat surface in the eastern part of the wetland. Towards the end of the subzone, a peak of the fungal cells belonging to NPP type HdV-200 correlate with a decline of Cyperaceae. The HdV-200 is mostly associated with biological decomposition of desiccated plant remains and helophytes (van Geel et al., 1989) suggesting seasonal dry conditions in Gahak at the mid- to late 19th century.

P-B3/NPP-B4 (Depth: ca 10-0 cm/Age: 1861-1952 CE):

The uppermost 5 cm of the Gahak record is particularly characterized by explosion of coprophagous fungal spores and high values of pollen indicator of grazing. *Artemisia* which currently dominates the subalpine zone of the area reaches its highest peak in the entire record. The peak of Caryophyllaceae most probably corresponds to the thorny cushion species of *Acanthophyllum* which are dominant in surrounding mountain steppes. All the evidence suggests a very high pastoral pressure on the environment of the Gahak wetland during the last one and half century.

Entomological data

31 taxa were identified based on a total of 259 fragments and 137 individuals, mostly belonging to Coleoptera (90%) with only two taxa belonging to Heteroptera and one taxon to Hymenoptera (Fig. 10). Identifications were limited to genus and family levels due to the lack of reference materials from the region. The coleopteran assemblages were relatively poor with average 13 taxa per sample. The two lowest samples (120-100 cm depth) were completely

devoid of insect remains. The richest sample in terms of number of fragments (n=58) was between 50-40 cm depth and the richest sample in terms of number of individuals (n=26) was between 80-70 cm depth. The taxa were clustered under five ecological groups based on their common habitats and one group of uncertain ecological affinity; phytophagous, aquatic, riparian, coprophagous, and uncertain (Fig. 10). The grouping is based on literature (see methodology section) and our field observations. Stratigraphically-confined vertical zonation of the entomological diagram suggests two major assemblage zones referred to as Ent-A and Ent-B (Figure 9). The lower assemblage zone Ent-A (100-50 cm: 810-1522 CE) is characterized by high abundance of both aquatic and riparian taxa and with low abundance but almost constant presence of coprophagous taxa. The upper zone Ent-B assemblage zone (50-0 cm: 1522-1952 CE) is characterized by lower abundance of aquatic and riparian taxa but higher abundance and more continuous presence of phytophagous and coprophagous taxa.




Figure 10. Palaeoentomological record of Gahak with the number of fragments and inferred individuals for each 10-cm thick sample. Identified taxa were put into five ecological groups.

The absence of insect remains in the lower 20 cm of the core suggest less water-logged conditions and/or more frequent subaerial exposure of the site causing the decaying of the fossil materials. This part corresponds to the pre-wetland phase of the site revealed in pollen and NPP diagrams. For the rest of the diagram, aquatic and riparian taxa are mostly present indicating that the wetland and water-logged conditions have existed at least since 800 CE. Continuous presence of abundant remains of aquatic beetle genera *Ochthebius* and *Enochrus* (Fig. 11) further suggests that the Gahak wetland did not undergo complete seasonal desiccation until very recently. These permanent water-logged conditions correspond to the MWP and most of the LIA. The above beetle taxa can thus be considered as bioindicators for the seasonality of wetland's moisture or water table because they can attest to the year-round water-logged conditions.




Figure 11. Four key beetle genera frequently encountered in Gahak palaeoentomological record. a. *Ochthebius* sp. (1.7 mm long), b. *Enochrus* sp. (5.9 mm long), c. *Carpelimus* sp. (3.6 mm long), and d. *Aphodius* sp. (6 mm long). a and b are indicators of high wetland water tables with no seasonal desiccation. c is a typical riparian taxon growing on wet mud flats, and d is a dung beetle suggesting the abundance of livestock around the site.

Coprophagous taxa are present almost all along the core suggesting that pastoral activities were practiced around the Gahak wetland since the beginning of the record. The upper half of the Gahak record displays more abundant fragments and individuals of dung beetles (*Aphodius* spp. and *Onthophagus* spp.). This can be linked to two environmental factors, the taphonomic processes and changes in the intensity of grazing activities. Contrary to the lower half of the diagram in which the wetland had higher water tables, the upper half of the diagram is characterized by organic sediment build-up and by lower water-tables and/or gradual terrestrialization of the wetland. This may have made the wetland floor more accessible to the livestock grazing and direct dropping of animal dung onto the sediment surface in shallow water. More abundant dung beetles are also compatible with ever increasing values of pollen indicators of grazing (see *Plantago lanceolata*-type), nitrophilous taxa (see *Trifolium*-type pollen) and coprophilous fungal spores (Ponel *et al.*, 2011) (Figs. 7 and 8). As far as the agricultural activities are concerned, only the presence of *Chaetocnema* sp. is noted, a genus which can occasionally be a pest of Poaceae including cereals. In the record, it is found in three intervals with cereal cultivation (compare Figs. 8 and 10).

The more noticeable occurrence of Heteroptera in the lower half of the diagram is also interesting and may indicate relatively higher temperatures. The diversity and abundance of this thermophilous group of insects would have thus increased during the Mediaeval Warm Period.

Discussion

MWP and LIA thermal and moisture variations reflected in Gahak record

High elevation wetlands and peat accumulations are invaluable sedimentary archives that can record changes in snow cover extension, alpine/subalpine flora and vegetation, agro-pastoral practices, and erosional events (Miehe et al., 2009; Djamali et al., 2009; Bell et al., 2021). On dry slopes of the Iranian mountains in the vicinity of desert areas, they form ‘green islands’ harboring a unique plant biodiversity (Naqinezhad et al., 2009; Naqinezhad et al. 2021) and providing an important source of drinking water for livestock and small-scale farming activities. Gahak wetland is one of these wetlands whose multi-proxy record provides a wealth of information on the combined effects of high frequency climatic variations and anthropogenic agro-pastoral practices on lake’s hydro-sedimentary environment and surrounding montane steppes. Not only the MWP and LIA are clearly recorded in Gahak sequence but their internal fluctuations are also traceable in both bioindicator and geochemical records (Figs. 5, 8-10).




Figure 12. The wheat cultivation phase (a) and Ti intensity variations (b) in Gahak record (this study) compared to a selection of regional and global proxy records of climate and solar activity. (c) Ti intensity variations in Lake Neor (Sharifi et al., 2015), (d) cereal pollen in Lake Almalou (Djamali et al., 2009a), (e) Lake Maharlou desiccation intervals (Brisset et al., 2019), (f) Caspian Sea level changes (Naderi-Beni et al., 2013 & Leroy et al., 2022), (g) Lake Urmia precipitation reconstruction (Sharifi et al., 2023), (h) Oxygen isotope record of Gejkar cave in Kurdistan (Flohr et al., 2017), (i) wet and dry events in Mesopotamia (Vogt et al., 2011), (j) spring/summer temperatures based on dendro-climatological data from Russian Altai, (k) Total solar irradiance (TSI) based on cosmonuclides production record (Bard et al., 2000). The Gahak agricultural phase during the MWP correlates with a warm but relatively wet period in the Iranian plateau as suggested by frequent desiccations of the Maharlou playa lake in SW Iran (e: Brisset et al., 2019) and more negative values of $\delta^{18}\text{O}$ of speleothem records in northern Zagros (h: Flohr et al., 2017) as well as the climate historical data from Mesopotamia (g: Vogt et al., 2011).

The MWP in Gahak is almost perfectly correlated with Lake Maharlou's (SW Iran) most important desiccation phase over the last four millennia (Fig. 12e; see also Brisset et al., 2019). Previously, Sharifi et al. (2015, 2018, and 2023) demonstrated how the downcore variation in Ti abundances during the Holocene can be related to the episodes of dry conditions over the interior of SW Asia. They concluded that the episodes of high Ti abundances in high-elevated Neor Lake (Sharifi et al. 2015, and 2018) are correlated to the episodes of high dust flux to the region during the dry periods. They also used lake sediment Ti/Al ratio from Lake Urmia in NW Iran along with the paleo-rainfall reconstruction (Fig. 12g) to show the correlation between periods of low precipitation-low lake levels with the dry episodes (Sharifi et al., 2023). The high values of XRF-measured Ti intensities of Gahak (Fig. 12b) during the MWP are also well-correlated with those from Lake Neor (NW Iran) (Fig. 12c) and low precipitation values over the Lake Urmia basin (Fig. 12g) suggesting relatively warm and dry conditions or at least less effective moisture over Western Asia. These trends are in line with the sudden shift in elemental z-scores (Fig. S2) from MWP to LIA indicating a significant change in elemental input and sedimentary regime of Gahak wetland corresponding to climatic shift from MWP to LIA. This climate shift is also correlated with the variation in total solar irradiance (TSI) where the observed dry episodes in Gahak, Neor Lake and Urmia Lake during the MWP are well correlated with higher solar irradiance values (Fig. 12k). The box plots and one-way ANOVA test for the Gahak Ti intensity variations further shows the significant difference between the MWP and LIA periods. The same hydroclimatic shift is characterized by dramatic rise of the Caspian Sea levels (Fig. 12f) after a lowstand during the MWP (Naderi Beni et al., 2013; Leroy et al., 2022). The non-pollen palynomorphs of Almalou peat bog (NW Iran) also suggest lower wetland moisture during the MWP compared to the LIA (not shown in Fig. 12) (Djamali et al., 2009). All these correlations suggest relatively low effective moisture during the MWP in central Iran. However, a speleothem record from northern Mesopotamia (Flohr et al., 2017) that shows a general drying trend starting from the beginning of the MWP onwards, indicates that the average effective moisture during the MWP was relatively higher and more stable compared to LIA (Fig. 12h). A historical climatological analysis of Arabic written sources of the Mediaeval Middle East also suggests wetter winters for 900-1100 CE although it shows drier winters for 1100-1250 CE (Vogt et al., 2011). It can thus be deduced that the Mediaeval Warm Period in the interior SW Asia was warm with relatively low effective moisture. However, our study shows that it was wet enough to maintain a sustainable agriculture lasting over three centuries. Indeed, almost all geochemical and biological proxies provide evidence

for a permanent wetland system during the MWP even if mean water levels were lower than the onset of the LIA. Water resources were thus available to perform irrigated agriculture during the MWP. In the next section, we try to infer the MWP climatic conditions based on agro-climatological requirements of wheat.




Figure 13. Box plots for Conover's all-pairs comparison test results applied to Ti intensity variations, farming-associated pollen, and pastoral pollen indicators corresponding to four main time intervals represented in Gahak record. The one-way ANOVA (Kruskal-Wallis H test) shows significant differences ($p < 0.001$) between the MWP and LIA periods for all tested variables.

Mobility of agro-pastoral communities during the MWP and LIA

The MWP has frequently been associated with a major agricultural transformation in continental Europe and the so-called 'cerealization' of northwestern Europe when the grain

production was largely increased and diversified, contributing to mediaeval economic growth (Schroeder, 2023). Bread wheat (*Triticum aestivum*), rye (*Secale cereale*), and oat (*Avena sativa*), became increasingly adopted and replaced spelt wheat (*T. spelta*) and barley. In the Alps and the Mediterranean mountains, higher temperatures of the MWP permitted the establishment of permanent settlements and expansion of agricultural landscapes in higher elevations (Mensing et al., 2016; Brandolini et al., 2023). Gahak record provides undisputed evidence of a three-century long intensive cultivation of cereals and possibly other crops in the uplands of the central Iranian plateau during the Mediaeval Warm Period (Fig. 8). Due to the quasi-absence of high-resolution pollen and archaeobotanical evidence from similar high-altitude contexts of the region, it is not currently possible to generalize this farming pattern into the whole region as an indicator of the beneficial effects of the MWP climatic change for mediaeval agriculture. Nonetheless, the beginning of the MWP in a high-resolution pollen record from Lake Almalou (NW Iran; 2400 m) is also characterized by a peak of cereal cultivation although it starts decreasing earlier than the Gahak sequence at around 1100 CE (Djamali et al., 2009).

Our findings suggest that sedentary farming communities settled the area from 950 CE for more than three centuries. Fossil pollen and entomological assemblages suggest that by the onset of the LIA, the sedentary agriculture and wheat cultivation became increasingly difficult to sustain and gave way to mobile seasonal agro-pastoralism which continues until today. This contrasted pattern of farming *versus* pastoralism during the MWP and the LIA is distinctly seen in the box plots of percentage values of anthropogenic pollen indicators (Fig. 13).

In general, the winter wheat is sown in autumn and harvested in summer of the next year. It needs moderately wet autumns, cold winters, and warm and wet springs and its yield particularly suffers from dry and cold springs and extended snow-cover, two conditions that were frequent during the LIA (Pfister, 2007; Pfister and Brázdil, 2006) but not during the MWP. In the highlands of the interior West Asia and Central Asia, wheat is grown as both irrigated and rain-fed grain crops with the highest percentage of dry-farmed wheat belonging to Iran in areas with mean annual precipitation exceeding 210-230 mm (Ballan and Bazin, 2012). When rain-fed, the wheat is sown in autumn to be harvested in June to October according to the elevation. When irrigated, it is also sown in autumn but will be harvested earlier from April (e.g., in lowlands of Khuzestan Province) to August. Irrigated winter bread wheat grows only on fertilized fields in rotation with summer crops (Ballan and Bazin, 2012). Bioclimatic characteristics of the winter bread wheat, predominantly *T. aestivum*, are helpful to estimate the seasonal patterns of temperature and precipitation of Gahak mountain site during MWP and LIA. A synthesis of many experimental works has estimated that lethal minimum and maximum temperatures for winter wheat are -17.2 and 47.5°C, and the optimal temperature range is 17-23°C (Porter and Gawith, 1999). These agro-meteorological data on winter wheat suggest several facts on the conditions of agroecosystems of the site during both MWP and LIA. First, they suggest that during the MWP, the seasonal pattern of precipitation was different. Although the annual precipitation amount could have been lower than today, the spring was most probably warm and wet and snow-cover extension was limited. The winter temperature minima stayed above the lethal temperatures of -17.2°C. These conditions changed dramatically at the onset of the LIA when dry and cold springs became more frequent with devastating biophysical effects on grain crops (Pfister, 2006). When snow cover lasts for a long

time in spring, there is not enough time to sow spring wheat because the needed time for grain-filling will be too short. The LIA was thus most probably characterized by an unsuitable combination of seasonal agro-meteorological factors for both irrigated and dry cereal farming. Nowadays, the annual precipitation (213 mm in a close weather station in Abyaneh; Fig. 1) seems not sufficiently high to sustain a rain-fed wheat cultivation even if the average of minimum temperatures of coldest month (-3.9°C) remains well above lethal temperature limit. It is thus more probable that farming communities of Gahak practiced irrigated wheat farming. The availability of water all year round in the Gahak basin as inferred by the permanent water-logged conditions, reinforces this possibility. The presence of modern qanat systems around the wetland (Fig. 2) suggests that this system could also have been used to supply irrigation water in the mediaeval time. The qanat system is an efficient groundwater exploitation system invented somewhere in the Iranian plateau and along the alluvial fans (Goblot, 1979). The transition from MWP to LIA is clearly indicating the decline of agricultural practices at the expense of the intensification of pastoralism and grazing pressure on mountain steppe ecosystems. Frequent extreme temperature events are considered the most important factor negatively impacting the wheat yield (Porter and Gawith, 1999). Such events were definitely more frequent during the LIA (Pfister and Brázdil, 2006) and made the high elevation cereal farming unsustainable. It can naturally be concluded that sedentary farming was replaced by the nomado-pastoral lifestyle in the middle of 13th century CE in Gahak. Judging from bioindicators, the mobile seasonal agro-pastoralism with limited subsistence agriculture would have continued through the LIA until today. Small peaks of cereal cultivation centered around 1580 CE and 1725 CE (Fig. 12a), may reflect short spells of the relative climate warming during the LIA.

Historical and geopolitical context of the Mediaeval Warm Period and Little Ice Age climatic changes in the central Iranian plateau

The agricultural expansion phase in Gahak corresponds to the reign of the Buyid, Ghaznavid, Seljuk, and Khwarazmian dynasties (Fig. 12), which successively took control of central Iran (Bosworth, 1968). The Buyid domination of the region (947 to 1029 CE) is particularly important because the Buyid era (934-1062 CE) marks a period of economic prosperity on the Iranian plateau (Busse, 1968). For instance, in the Fars region ('Persis') in SW Iran, large-scale hydraulic projects supported extensive agricultural activities (Saeidi Ghavi Andam et al., 2020). The Buyids, a Shi'i dynasty, gave particular favour to cities dominated by Muslim Shi'i populations, including Qum, Kashan, and surrounding areas (Amanat and Mottahedeh, 2018). Less is known about the history of the Kashan region under the Ghaznavids, who ruled from 1029 to 1051 CE. However, the region witnessed the peak of its economic prosperity during the Seljuk dynasty from 1051 CE and became a major centre for ceramic production (Amanat and Mottahedeh, 2018). At this time, increased demography and demand for food would have resulted in agricultural expansion in the highlands of Kashan and surrounding regions.

The end of the Mediaeval Warm Period agricultural phase in Gahak is remarkably well correlated with the Mongol invasion of western Asia (Fig. 12). The Mongols defeated the Khwarazmian army in 1219-1221 CE and invaded mainland Persia, including central Iran, in 1220-1240 CE (Boyle, 2007). The Mongol invasion also perfectly correlates with the beginning

of the Little Ice Age. There seems to be no conclusive evidence on the real impact of this invasion on the socio-economy of the Kashan area (Amanat and Mottahedeh, 2018). Nevertheless, a quote from the Persian historian Zakariya al-Qazvini, dated to 1283 CE, suggests that 50 years after the arrival of the Mongols and under the Ilkhanate rule, Kashan was still maintaining its important position as a major centre for the production and transportation of utensils and vessels (quoted in Amanat and Mottahedeh, 2018, pp. 406-407). A close look at wheat pollen curves shows that while pollen percentage values start decreasing from 1200 CE, concentration values remain relatively high for the next 150 years during the Ilkhanates and early Timurids (Fig. 12). Hence, it is most probably the severe climatic conditions of the LIA that ended the MWP agricultural phase and favoured a nomadic seasonal pastoral lifestyle for the following centuries. Similar to our conclusion, a recent study suggests that the decline of Central Asian irrigation-based urban societies was more related to a long-term drought regime rather than the mere Mongol invasion and destruction of cities in 1219 (Toonen et al., 2020).

For more than two centuries (1501-1736 CE) during the Little Ice Age, central parts of Persia experienced a period of political stability under the Safavid Empire. The near absence of agricultural activities in Gahak during the Safavid era is thus more logically attributed to severe climatic conditions rather than socio-political factors. The following quote from the book entitled *History of Kashan* (mid-18th century) suggests abundant snow and névés in the Karkas Mountains west of Kashan during the last decades of the Little Ice Age (early to mid-18th century):

“The winding road leads to the quaint village of Rahagh, nestled among the Sardsir (‘highland pasture’) Kashan hills to the west of Kashan, eight parasangs (approximately 48 km) away from the city. Beyond the village lies a majestic mountain with numerous valleys, each boasting its own water sources, lush trees, crimson barberries, and verdant fields. The mountain is two parasangs from the village (approximately 12 km), and its lofty peak remains snow-capped for years on its right flank, while on its left side, a road descends past the Gadok pass and joins the Wasaf river” (Zarrabi, 2000).

Conclusions

The alpine/subalpine wetlands of arid and semi-arid Southwest Asia are unique archives of hydroclimatic variations, ecosystem dynamics, and anthropogenic activities during the Holocene. The information obtained from these archives can be compared and contrasted with documentary and historical-climatological proxy data, especially for the pre-instrumental period. However, sources on mediaeval and pre-industrial human activities in Southwest Asian mountain areas are scarce compared to European mountain systems, highlighting the significance of sedimentary archives in uncovering the modality and intensity of these activities.

Our case study demonstrates that climatic shifts and their associated impacts on agro-pastoral practices, lifestyle changes, and the mobility of human communities have been clearly recorded

in these archives. Our data suggest that the Mediaeval Warm Period would have favoured high-elevation agricultural activities, while the Little Ice Age increased seasonal agro-pastoral practices in the highlands of Southwest Asia. Such information may be missing in written sources available to historians and ethno-archaeologists. These archives can refine our understanding of the role of small farming communities living in remote mountain areas in the economy of Mediaeval Persia and surrounding regions.

Many high-elevation wetlands and their sedimentary records are threatened by overgrazing, man-made drainage for land reclamation, and erosion. It is highly recommended that conservation measures be rapidly implemented to protect these high-elevation wetlands. The sedimentary sequences of the most endangered and eroding deposits should be systematically sampled, and the sediment cores deposited and kept in appropriate natural repositories for future research, similar to the current efforts for retreating mountain glaciers worldwide through the Ice Memory Initiative <https://www.ice-memory.org>.

Acknowledgement

We wish to thank Prof. Reza Mansouri (former director of the Iranian National Observatory - INO) for his encouragement and full support from the beginning of this study, and Dr. Habib Khosroshahi, the current director of INO, for continuing to support our project. A special thanks is due to Mr. Alireza Behnam, the technical staff member at INO, without whose involvement, devotion, and support in the field, this study would not have fulfilled its objectives. Laboratory analyses and radiocarbon measurements were supported by the “PALEO-PERSEPOLIS” ANR-DFG project (2015–2019) funded by the Agence Nationale de la Recherche (ANR-14-CE35-0026-01), the LIA project entitled “HAOMA” funded by the Centre National de la Recherche Scientifique (CNRS), and the MITRA project funded by the Swiss National Science Foundation. Field investigations were also partly supported by the Franco-Iranian Gundishapour joint project entitled “KECO-INO” (2015-2016).

References

- Amanat, M., Mottahedeh, R.P., 2018. Medieval Kashan: crossroads of commerce and culture. *Eurasian Studies* 16, 395-429.
- Arias, S.L., Mary, V.S., Velez, P.A., Rodriguez, M.G., Otaiza-González, S., Theumer, M.G., 2021. Where does the peanut smut pathogen, *Thecaphora frezii*, fit in the spectrum of smut diseases? *Plant Disease* 105, 2268-2280.
- Ballan, D., Bazin, M., 2012. Gandom. *Encyclopedia Iranica* 10, 270-276.
- Bard, E., Raisbeck, G., You, F., Jouzel, J., 2000. Solar irradiance during the last 1200 years based on cosmogenic nuclides. *Tellus B: Chemical and Physical Meteorology* 52, 985-992.
- Barlow, M., Zeitchik, B., Paz, S., Black, E., Evans, J., Hoel, A., 2016. A Review of Drought in the Middle East and Southwest Asia. *Journal of Climate* 29, 8547-8574.
- Bauer, R., Garnica, S., Oberwinkler, F., Reiss, K., Weiß, M., Begerow, D., 2015. Entorrhizomycota: a new fungal phylum reveals new perspectives on the evolution of fungi. *PLoS ONE* 10, e0128183.

Bayer Altin, T., Kaya, M., 2020. Climatic and social change during the Little Ice Age in Cappadocia Vicinity, Southern Central Anatolia, Turkey. *Regional Environmental Change* 20, 16. <https://doi.org/10.1007/s10113-020-01604-x>.

Bell, B.A., Fletcher, W.J., Hughes, P.D., Cornelissen, H.L., Fink, D., Rhoujjati, A., 2021. Palynological evidence from a sub-alpine marsh of enhanced Little Ice Age snowpack in the Marrakech High Atlas, North Africa. *Vegetation History and Archaeobotany* 31, 49-66.

Bertello L, Pevtsov AA, Ulrich RK (2020) 70 Years of Chromospheric Solar Activity and Dynamics. *Astrophys J* 897:181. <https://doi.org/10.3847/1538-4357/ab9746>

Beug, H.J., 2004. Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete. Verlag Dr Friedrich Pfeil, München.

Blaauw, M., 2010. Methods and code for “classical” age-modelling of radiocarbon sequences. *Quaternary Geochronology* 5, 512–518.

Bosworth, C.E., 1968. Chapter 1: The Political and Dynastic History of the Iranian World (A.D. 1000-1217). In: Boyle, J.A. (Ed.) *The Cambridge History of Iran, Volume V: The Saljuq and Mongol Periods*, Cambridge University Press, Cambridge, pp. 1-202 .

Boyle, J.A., Chapter 4: Dynastic and political history of the Il-Khans. In: Boyle, J.A. (Ed.) *The Cambridge History of Iran, Volume V: The Saljuq and Mongol Periods*, Cambridge University Press, Cambridge, pp. 303-421.

Brandolini, F., Kinnaird, T.C., Srivastava, A., Turner, S., 2023. Intensification of terrace agriculture on the Northern Apennines during the Medieval Climate Anomaly. *EGU General Assembly 2023*, Vienna, Austria, 24–28 Apr 2023, EGU23-5591, <https://doi.org/10.5194/egusphere-egu23-5591>.

Brisset, E., Djamali, M., Bard, E., Borschneck, D., Gandouin, E., Garcia, M., Stevens, L., Tachikawa, K., 2019. Late Holocene hydrology of Lake Maharlou, southwest Iran, inferred from high-resolution sedimentological and geochemical analyses. *Journal of Paleolimnology* 61, 111-128.

Bronk Ramsey, C., 1995. Radiocarbon calibration and analysis of stratigraphy: The OxCal program. *Radiocarbon*, 37, 425–430.

Büntgen, U., Myglan, V.S., Charpentier Ljugqvist, F., McCormick, M., Di Cosmo, ., Sigl, M., Jungclaus, J., Wagner, S., Krusic, P.J., Esper, J., Koplan, J.O., de Vaan, M.A., Luterbacher, J., Wacker, L., Tegel, W., Kirilyanov, A.V., 2016. Cooling and societal change during the Late Antique Little Ice Age from 536 to around 660 AD. *Nature Geoscience* 9, 231-236.

Büntgen, U., Allen, K., Anchukaitis, K.J., et al., 2021. The influence of decision-making in tree ring-based climate reconstructions. *Nature Communications* 12, 3411.

Burone, L., Ortega, L., Franco-Fraguas, P., et al., 2013. A multiproxy study between the Río de la Plata and the adjacent South-western Atlantic inner shelf to assess the sediment footprint of river vs. marine influence. *Continental Shelf Research* 55, 141–154.

Busse, H., 1975. Chapter 7: Iran under the Buyids. In: Fry, R. N. (Ed.) *The Cambridge history of Iran. Volume IV: The period from the Arab invasion to the Saljuqs*, Cambridge University Press, Cambridge, pp. 250-304.

Chater, A., Smith, P.A., 2018. Recent finds of *Entorrhiza* root smuts. *Field Mycology* 19, 55-60.

Cleveland, W.S., Grosse, E., Shyu, W.M., 1992. Chapter 8: Local regression models. In: Chambers, J.M., Hastie, T.J. (eds.), *Statistical Models in S*. Routledge, New York, 68 p.

Conover, W. J. , R. L. Iman, 1979. On multiple-comparisons procedures, Technical Reports LA-7677-MS, Los Alamos Scientific Laboratory.

Coope, G.R., 1986. Coleoptera analysis. In: Berglund, B.E. (Ed.) *Handbook of Holocene Palaeoecology and Palaeohydrology*. Wiley & Sons, Chichester. pp. 703–713.

Cooper, G.R.J., Cowan, D.R., 2008. Comparing time series using wavelet-based semblance analysis. *Computers and Geosciences* 34, 95–102. doi:10.1016/j.cageo.2007.03.009

Croudace, I.W., Rindby, A., and Rothwell, R.G., 2006. ITRAX: description and evaluation of a new multi-function X-ray core scanner. *Geological Society of London Special Publication* 267, 51– 63.

Cugny, C., 2011. Apport des microfossiles non-polliniques à l'histoire du pastoralisme sur le versant nord Pyrénéen: entre référentiels actuels et reconstitution du passé. Vol. 1 & 2. Unpublished PhD Dissertation, Université Toulouse 2, Le Mirail.

Dehghani, M., Djamali, M., Gandouin, E., Akhiani, H., 2017. A pollen rain - vegetation study along a 3600 m mountain - desert transect in the Irano-Turanian region; implications for the reliability of some pollen ratios as moisture indicators. *Review of Palaeobotany and Palynology* 247, 133-149.

Del Socorro Lozano-Garcia, M., Caballero, M., Ortega, B., Rodriguez, A., Sosa, S., 2007. Tracing the effects of the Little Ice Age in the tropical lowlands of eastern Mesoamerica. *PNAS* 104, 16200-16203.

Demory, F., Uehara, M., Quesnel, Y., Rochette, P., Romey, C., Tachikawa, K., Garcia, M., Borschneck, D., Pignol, L., Bard, E., & Andrieu-Ponel, V., 2019. A new high-resolution magnetic scanner for sedimentary sequences. *Geochemistry, Geophysics, Geosystems* 20, 3186–3200..

Dypvik H, Harris NB 2001. Geochemical facies analysis of fine-grained siliciclastics using Th/U, Zr/Rb and (Zr+Rb)/Sr ratios. *Chem Geol* 181:131–146. [https://doi.org/10.1016/S0009-2541\(01\)00278-9](https://doi.org/10.1016/S0009-2541(01)00278-9).

Djamali, M., de Beaulieu, J.-L., Miller, N., Andrieu-Ponel, V., Berberian, M., Gandouin, E., Lahijani, H., Ponel, P., Salimian, M., Guiter, F., 2009. A late Holocene pollen record from Lake Almalou in NW Iran: evidence for changing land-use in relation to some historical events during the last 3700 years. *Journal of Archaeological Science* 36, 1346-1375.

Djamali, M., Akhiani, H., Khoshravesh, R., Andrieu-Ponel, V., Ponel, P., Brewer, S., 2011. Application of the Global Bioclimatic Classification to Iran: implications for understanding the modern vegetation and biogeography. *Ecologia Mediterranea* 37, 91-114.

- Djamali, M., Gondet, S., Ashjari, J., Aubert, C., Brisset, E., Longerey, J., Marriner, N., Mashkour, M., Millet, N.F., Naderi-Beni, A., Pourkerman, M., Rashidian, E., Rigot, J.-B., Shidrang, S., Thiéry, A., Gandouin, E., 2018. Karstic-spring wetlands of the Persepolis Basin, SW Iran: unique sediment archives of Holocene change and human impact. *Canadian Journal of Earth Sciences* 55, 1158-1172.
- Djamali, M., Baumel, A., Brewer, S., Jackson, S.T., Kadereit, J.W., López-Vinyallonga, S., Mehregan, I., Shabanian, E., Simakova, A., 2012. Ecological implications of *Cousinia* Cass. (Asteraceae) persistence through the last two glacial-interglacial cycles in the continental Middle East for the Irano-Turanian flora. *Review of Palaeobotany and Palynology* 172, 10-20.
- Emadodin, I., Reinsch, T., Taube F., 2019. Drought and Desertification in Iran. *Hydrology* 6, 66.
- Eskandari, M., Assadi, M., Shirzadian, S., Mehregan, I., 2019. Ethnobotanical study and distribution of the *Solanum* Section *Solanum* species (Solanaceae) in Iran. *Journal of Medicinal Plants* 18, 85-98.
- Eskandari, M., Shirzadian, S., 2022. Introduction of important weeds of Solanaceae in Iran. Iranian Research Institute of Plant Protection technical report 61382 (in Persian).
- Flohr, P., Fleitmann, D., Zorita, E., Sadekov, A., Cheng, H., Bosomworth, M., Edwards, L., Matthews, W., Matthews, R., 2017. Late Holocene droughts in the Fertile Crescent recorded in a speleothem from northern Iraq. *Geophysical Research Letters* 44, 1528-1536.
- Gierlowski-Kordesch, E. H., 2010. Lacustrine carbonates: Developments in sedimentology, v. 61, p. 1-101.
- Gurjazkaite, K., Routh, J., Djamali, M., Vaezi, A., Poher, Y., Naderi-Beni, A., Tavakoli, V., Klyn, H., 2018. Vegetation history and human-environment interactions through the late Holocene in Konar Sandal, SE Iran. *Quaternary Science Reviews* 194, 143-155.
- Goblot, H., 1979. Les Qanats : une technique d'acquisition de l'eau. École des hautes études en sciences sociales, Paris.
- Govin, A., Holzwarth, U., Heslop, D., et al., 2012. Distribution of major elements in Atlantic surface sediments (36°N-49°S): Imprint of terrigenous input and continental weathering. *Geochemistry, Geophysics, Geosystems* 13, 1–23.
- Grimm, E.C., 1987. Coniss: a Fortran 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares. *Computers and Geosciences* 13, 13–35.
- Grimm, E.C., 2004/2005. TILIA and TGView Software. Version 2.0.2. Illinois State University, Illinois.
- Haghani, S., Leroy, S.A.G., Khdir, S., Naderi Beni, A., Lahijani, H.A.K., 2015. An early ‘Little Ice Age’ brackish water invasion along the south coast of the Caspian Sea (sediment of Langarud wetland) and its wider impacts on environment and people. *The Holocene* 26, 3-16.
- Hernández-Restrepo, M., Groenwald, J.Z., Elliott, M.L., Canning, G., McMillan, V.E., Crous, P.W., 2016. Take-all or nothing. *Studies in Mycology* 83, 19-48.

Hammer, Ø., Harper, D.A.T., Ryan, P.D. 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4.

Hughes, P.D. 2014. Little Ice Age glaciers in the Mediterranean mountains. *Mediterranée* 122, 63-79.

Juggins, S., 2022. rioja: Analysis of Quaternary Science Data, R package version (1.0-5). (<https://cran.r-project.org/package=rioja>).

Kjær, K.H., Bjørk, A.A., Kjeldsen, K.K., Hansen, E.S., Andresen, C.S., Siggaard-Andersen, M.-L., Shfaqat A. Khan e, Anne Sofie Søndergaard f, Colgan, W., Schomacker, A., Woodroffe, S., Funder, S., Rouillard, A., Jensen, J.F., Larsen, N.K., 2022. Glacier response to the Little Ice Age during the Neoglacial cooling in Greenland. *Earth-Science Reviews* 227, 103984.

Koch, K., 1989a. Die Käfer Mitteleuropas, Ökologie 1. Krefeld: Goecke and Evers.

Koch, K., 1989b. Die Käfer Mitteleuropas, Ökologie 2. Krefeld: Goecke and Evers.

Koch, K., 1992. Die Käfer Mitteleuropas, Ökologie 3. Krefeld: Goecke and Evers.

Koch, J., Clague, J.J., 2011. Extensive glaciers in northwest North America during Medieval time. *Climatic Change* 107, 593–613.

Koch, J., 2015. Little Ice Age and recent glacier advances in the Cordillera Darwin, Tierra del Fuego, Chile. *Anales Instituto Patagonia* 43, 127-136.

Kroonenberg, S.B., Abdurakhmanov, G.M., Badyukova, E.N., van der Borg, K., Kalashnikov, A., Kasimov, N.S., Rychagov, G.I., Svitoch, A.A., Vonhof, H.B., Wesselingh, F.P., 2007. Solar-forced 2600 BP and Little Ice Age highstands of the Caspian Sea. *Quaternary International* 173–174, 137–143.

Leroy, S.A.G., Lahijani, H., Djamali, M., Vahabi-Moghadam, Naqinezhad, A., Shah-Hosseini, M., Miller, C.S., Tavakoli, V., 2011. Late Little Ice Age palaeoenvironmental records from the Anzali and Amirkola Lagoons (south Caspian Sea): Vegetation and sea level changes. *Palaeogeography Palaeoclimatology Palaeoecology* 302, 415-434.

Lo, F.L., Chen, H.F., Fang, J.N., 2017. Discussion of suitable chemical weathering proxies in sediments by comparing the dissolution rates of minerals in different rocks. *Journal of Geology* 125, 83–99.

Lomb, N. R., 1976: Least-squares frequency analysis of unequally spaced data. *Astrophysics and Space Science* Vol. 39, pp. 447-462.

Lupker, M., Aciego, S.M., Bourdon, B., et al., 2010. Isotopic tracing (Sr, Nd, U and Hf) of continental and marine aerosols in an 18th century section of the Dye-3 ice core (Greenland). *Earth Planetary Science Letters* 295, 277–286.

Ma, L., Vaquero, J.M., 2020. New evidence of the Suess/de Vries cycle existing in historical naked-eye observations of sunspots. *Open Astron* 29, 28–31.

Magnan, G., van Bellen, S., Davies, L., Froese, D., Garneau, M., Mullan-Boudreau, G., Zacccone, C., Shotyky, W., 2018. Impact of the Little Ice Age cooling and 20th century climate change on peatland

vegetation dynamics in central and northern Alberta using a multi-proxy approach and high-resolution peat chronologies. *Quaternary Science Reviews* 185, 230-243.

Makri, S., Wienhues, G., Bigalkea, M., Gilli, A., Reyc, F., Tinner, W., Vogel, H., Grosjean, G., Variations of sedimentary Fe and Mn fractions under changing lake mixing regimes, oxygenation and land surface processes during Late-glacial and Holocene times. *Science of the Total Environment*, 755, 143418. <https://doi.org/10.1016/j.scitotenv.2020.143418>.

Martins, V., Dubert, J., Jouanneau, J.M., et al., 2007. A multiproxy approach of the Holocene evolution of shelf-slope circulation on the NW Iberian Continental Shelf. *Marine Geology* 239, 1–18.

Moore, P.D., Webb, J.A., Collinson, M.E., 1991. *Pollen Analysis*, 2nd ed. Blackwell, Oxford.

Matthews, J.A. and Briffa, K.R., 2005. The ‘Little Ice Age’: re- evaluation of an evolving concept. *Geographiska Annaler* 87A, 17–36.

McGregor, H., Evans, M., Goosse, H. et al., 2015. Robust global ocean cooling trend for the pre-industrial Common Era. *Nature Geosci* 8, 671–677.

Mensing, S., Tunno, I., Cifani, G., Passigli, S., Noble, P., Archer, C., Piovesan, G., 2016. Human and climatically induced environmental change in the Mediterranean during the Medieval Climate Anomaly and Little Ice Age: A case from central Italy. *Anthropocene* 15, 49-59.

Miehe, G., Miehe, S., Kaiser, K., Reudenbach, C., Behrendes, L., Duo, L., Schlütz, F., 2009. How old is pastoralism in Tibet? An ecological approach to the making of a Tibetan landscape. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 276, 130–147.

Mosaferi, S., Sheidai, M., Keshavarzi, M., Noormohammadi, Z., 2015. *Polygonum aviculare* (Polygonaceae) subspecies, new records for the flora of Iran. *Modern Phytomorphology* 8, 31-36.

Mulitza, S., Prange, M., Stuut, J.B., et al., 2008. Sahel megadroughts triggered by glacial slowdowns of Atlantic meridional overturning. *Paleoceanography* 23, 1–11.

Mursula K, Usoskin IG, Kovaltsov GA (2002) A 22-year cycle in sunspot activity. *Adv Sp Res* 29:1979–1984. [https://doi.org/10.1016/S0273-1177\(02\)00244-2](https://doi.org/10.1016/S0273-1177(02)00244-2)

Naqinezhad, A., Jalili, A., Attar, F., Ghahreman, A., Wheeler, B.D., Hodgson, J.G., Shaw, S.C. and Maassoumi, A.A., 2009. Floristic characteristics of the wetland sites on dry southern slopes of the Alborz Mts., N. Iran: The role of altitude in floristic composition. *Flora* 204, 254-269.

Naqinezhad, A., Nowak, A., Świerszcz, S., Jalili, A., Kamrani, A., Wheeler, B. D., ... & Hájek, M., 2021. Syntaxonomy and biogeography of the Irano-Turanian mires and springs. *Applied Vegetation Science* 24, e12571.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. *vegan: Community Ecology Package*. R package version 2.5-6. <https://CRAN.R-project.org/package=vegan>

PAGES2k Consortium, 2017. A global multiproxy database for temperature recon- structions of the Common Era. *Sci. Data* 4:170088.

- Parnell, A. C., Haslett, J., Allen, J. R. M., Buck, C. E., Huntley, B., 2008. A flexible approach to assessing synchronicity of past events using Bayesian reconstructions of sedimentation history. *Quaternary Science Reviews* 27, 1872–1885.
- Peregrine, P.,N., 2020. Climate and social change at the start of the Late Antique Little Ice Age. *The Holocene* 30, 1643-1648.
- Perez, L., García-Rodríguez, F., Hanebuth, T.J.J., 2016. Variability in terrigenous sediment supply offshore of the Río de la Plata (Uruguay) recording the continental climatic history over the past 1200 years. *Climate of the Past* 12, 623–634.
- Pfeffer, A., 1995. Zentral- und westpaläarktische Borken- und Kernkäfer. Pro Entomologia, c/o Naturhistorisches Museum Basel.
- Peristykh, A.N., Damon, P.E., 2003. Persistence of the Gleissberg 88-year solar cycle over the last ~12,000 years Evidence from cosmogenic isotopes. *Journal of Geophysical Research Sp Phys* 108:1–15. <https://doi.org/10.1029/2002JA009390>
- Pfister, C., 1981. An analysis of the Little Ice Age climate in Switzerland and its consequences for agricultural production. In: Wigley, T.M.L., Ingram, M.J., Farmer, G. (Eds.) *Climate and history: studies in past climates and their impact on Man*. Cambridge University Press, Cambridge, pp. 214-248.
- Pfister, C., Brázdil, R., 2006. Social vulnerability to climate change in the “Little Ice Age”: an example from Central Europe in the early 1770’s. *Climate of the Paast* 2, 115-129.
- Piątek, M., Lutz, M., Wang, Y., Wang, S., Kellner, R., 2021. *Thecaphora dahuangis*, a new species causing leaf smut disease of the traditional medicinal plant *dahuang* (*Rheum palmatum*) in China. *Plant pathology* 70, 1292-1299.
- Ponel P., Court-Picon M., Badura M., Guiter F., Beaulieu J.-L. de, Andrieu-Ponel V., Djamali M., Leydet M., Gandouin E. & Buttler A., 2011. Holocene history of Lac des Lauzons (2180 m a.s.l.), reconstructed from multiproxy analyses of Coleoptera, plant macro-remains and pollen (Hautes-Alpes, France). *The Holocene* 21, 565-582.
- Porter, J.R., Gawith, M., 1999. Temperatures and the growth and development of wheat: a review. *European Journal of Agronomy* 10, 23-36.
- Quinn, J.A., 2009. *Desert biomes*. Greenwood Press, Westport.
- Radfar, J., 1993. 1:100.000 geological map of Kashan. Geological Survey of Iran.
- Ram, B., Singh, K.P., 2004. Smuts of wheat: a review. *Indian Phytopathology* 57, 125-134.
- Ramezani, E., Marvie Mohadjer, M.R., Knapp, H.-D., Ahmadi, H., Joosten, H., 2008. The late-Holocene vegetation history of the Central Caspian (Hyrcanian) forests of northern Iran. *Holocene* 18, 305–319.

Ramezani, E., 2013. Palynological reconstruction of late-Holocene vegetation, climate, and human impact in Kelardasht (Mazandaran province, N Iran). *Iranian Journal of Forest and Poplar Research* 21, 48-62.

Sharifi, A., Pourmand, A., Canel, E.A., Ferer-Tyler, E., Peterson, L.C., Aichner, B., Feakins, S.J., Daryaee, T., Djamali, M., Naderi Beni, A., Lahijani, H.A.K., Swart, P.K., 2015. Abrupt climate variability since the last deglaciation based on a high-resolution, multi-proxy peat record from NW Iran: The hand that rocked the Cradle of Civilization? *Quaternary Science Reviews* 123, 215-230.

Sharifi, A., Murphy, L.N., Pourmand, A., Clement, A.C., Canuel, E.A., Naderi Beni, A., A.K. Lahijani, H., Delanghe, D., Ahmady-Birgani, H., 2018. Early-Holocene greening of the Afro-Asian dust belt changed sources of mineral dust in West Asia. *Earth and Planetary Science Letters* 481, 30-40.

Sharifi, A., Djamali, M., Peterson, L.C., Swart, P.K., Guadalupe Pulido Ávila, M., Esfahanejad, M., de Beaulieu, J.-L., Lahijani, H.A.K., Pourmand, A., 2023. The Rise and Demise of Iran's Urmia Lake during the Holocene and the Anthropocene: "What's past is prologue". *Regional Environmental Change*, 23, 121.

Reille, M., 1992. Pollen et spores d'Europe et d'Afrique du Nord. *Laboratoire de botanique historique et de palynologie*, Marseille.

Reille, M., 1995. Pollen et spores d'Europe et d'Afrique du Nord. *Laboratoire de botanique historique et de palynologie-Supplément 1*, Marseille.

Reille, M., 1998. Pollen et spores d'Europe et d'Afrique du Nord. *Laboratoire de botanique historique et de palynologie-Supplément 2*, Marseille.

Reimer, P., Austin, W., Bard, E., Bayliss, A., Blackwell, P., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R., Friedrich, M., Grootes, P., Guilderson, T., Hajdas, I., Heaton, T., Hogg, A., Hughen, K., Kromer, B., Manning, S., Muscheler, R., ... Talamo, S., 2020. The IntCal20 northern hemisphere radiocarbon age calibration curve (0–55 cal kBP). *Radiocarbon* 62, 725–757.

Reiss, K., Schön, M.E., Ziegler, R., R., Lutz, M., Shivas, R.G., Piatek, M., Garnica, S., 2019. The origin and diversification of the Entorrhizales: deep evolutionary roots but recent speciation with phylogenetic and phylotypic split between associates of the Cyperaceae and Juncaceae. *Organisms Diversity and Evolution* 19, 13-30.

Ritz, K., 2005. Fungi. In: Hillel, D. (Ed.) *Encyclopedia of soils in the Environment*, pp. 110-119.

Rowan, A.V., 2017. The 'Little Ice Age' in the Himalaya: A review of glacier advance driven by Northern Hemisphere temperature change. *The Holocene* 27, 292-308.

Scargle, J. D., 1989: Studies in astronomical time series analysis. II - Statistical aspects of spectral analysis of unevenly spaced data. *Astrophysical Journal* 1:263 pp. 835-853.

Schroeder, N., 2023. The 'Cerealization' of continental north-west Europe, c. 800-1200. In: Hamerow, H., McKerracher, M. (Eds.) *New perspectives on the medieval agricultural revolution; crop, stock and furrow*. Liverpool University Press, Liverpool. pp. 199-210.

- Shumilovskikh, L.S., van Geel, B., 2020. Non-pollen palynomorphs. In: Henry, A.G. (Ed.) Handbook for the Analysis of Micro-Particles in Archaeological Samples. Springer, pp. 65–94.
- Steinhilber F, Abreu J a, Beer J, et al (2012) 9,400 Years of Cosmic Radiation and Solar Activity From Ice Cores and Tree Rings. *Proceedings of the National Academy of Science of the USA* 109, 5967–71.
- Stockmarr, J., 1971. Tablets with spores used in absolute pollen analysis. *Pollen et Spores* 13, 615–621.
- Thomson, M.J., MacDonald, G.M., 2020. Climate and growing season variability impacted the intensity and distribution of Fremont maize farmers during and after the Medieval Climate Anomaly based on a statistically downscaled climate model. *Environmental Research Letters* 15, 105002.
- Toonen, W.H.J., Macklin, M.G., Dawkes, Giles, Durcan, J.A., Leman, M., Nikolayev, Y., Yegorov, A., 2020. A hydromorphic reevaluation of the forgotten river civilizations of Central Asia. *Proceedings of the National Academy of Sciences* 117, 32982-32988.
- van Geel, B., Coope, G.R., van der Hammen, T., 1989. Palaeoecology and stratigraphy of the Lateglacial type section at Usselo (The Netherlands). *Review of Palaeobotany and Palynology* 60, 25-129.
- Van Geel, B., Aptroot, A., 2006. Fossil ascomycetes in Quaternary deposits. *Nova Hedwigia* 89, 313–329.
- Van Zeist, W., Bottema, S., 1977. Palynological investigations in western Iran. *Palaeohistoria* 19, 19–85.
- Vasighzadeh, A., Zafari, D., Selçuk, F., Hüseyin, E., Kurşat, M., Lutz, M., Piątek, M., 2014. Discovery of *Thecaphora schwarzmaniana* on *Rheum ribes* in Iran and Turkey: implications for the diversity and phylogeny of leaf smuts on rhubarbs. *Mycological Progress* 13, 881–892.
- Vogt, S., Glaser, R., Luterbacher, J., Riemann, D., Al Dyab, Gh., Schoenbein, J., Garcia-Bustmante, E., 2011. *PAGES* 19, 28-29.
- Waldinger, M., 2022. Medieval Warm Period (MWP) and Little Ice Age (LIA). *Journal of Political Economy* 130, 2275-2314.
- Walker, J., 1972. Type studies on *Gaeumannomyces graminis* and related fungi. *Transactions of the British Mycological Society* 58, 427–457.
- Zarrabi, A. K., 2000, *Tarikh i Kashan* (The History of Kashan), Amir Kabir, Tehran.